

## Physics 471 – Fall 2024

### Homework #2 – due Wednesday, September 11

Point values for each problem are in square brackets

1. [1] Given 5 consecutive flips of a coin: What is the probability of first getting two heads in a row, followed by three consecutive tails?

2. [2] Sketch a graph of  $f(x) = 3e^{-(x^2/25)}$  versus  $x$ . (Clearly indicate the  $x$  and  $y$  scales you choose. “Sketch” means hand-drawn; don’t use a computer package to do this for you.)

3. [5] **Practice with Statistics.** Feel free to use a calculator (or spreadsheet) to add/divide/etc, but please don’t use built-in statistical functions.

You measure the lengths of 14 objects in a box, and get the following results (in cm):

1; 3; 3; 4; 4; 5; 5; 5; 5; 7; 7; 7; 8; 9

a) What is the probability that an object chosen at random from this box has length 4 cm? (Here and below, assume equal probability for selecting any one object)

b) What’s the average length  $\langle L \rangle$  (also called the expectation value of length) of the objects? What is the probability that an object chosen at random from the box has length =  $\langle L \rangle$  ?

c) What is the average value of the square of the lengths,  $\langle L^2 \rangle$  ?

d) Use your results to find the standard deviation  $\sigma$  of lengths of objects in this box

e) What is the probability that the length of an object chosen randomly from this box is in the range  $\langle L \rangle \pm \sigma$  ? (Given what you know about stand. deviation, does this seem reasonable?)

4. [4] **Orthogonality and inner products.** Consider the following 3 state vectors:

$$\begin{aligned} |\psi_1\rangle &= \frac{2}{\sqrt{5}} |+\rangle + \frac{1}{\sqrt{5}} |-\rangle \\ |\psi_2\rangle &= \frac{\sqrt{2}}{\sqrt{3}} |+\rangle + \frac{1}{\sqrt{3}} i |-\rangle \\ |\psi_3\rangle &= \frac{1}{\sqrt{2}} |+\rangle + \frac{1}{\sqrt{2}} e^{i\pi/4} |-\rangle \end{aligned}$$

Use bra-ket notation in your calculations (*not* matrix notation, please)

Use orthogonality and normalization of our basis:  $\langle +|+\rangle = \langle -|-\rangle = 1$ ,  $\langle +|-\rangle = \langle -|+\rangle = 0$ .

a) [3] For each of the 3 states above, find some vector that is orthogonal to it. (Use our convention that we keep the coefficient of the  $|+\rangle$  basis ket positive and real.)

b) [1] Calculate the inner products  $\langle \psi_2 | \psi_3 \rangle$  and  $\langle \psi_3 | \psi_2 \rangle$ . How are these two results related to one another?

**5. [5] Kets (state vectors).** Consider the following three (candidate) state vectors:

$$\begin{aligned} |\psi_1\rangle &= 3|+\rangle - 4|-\rangle \\ |\psi_2\rangle &= 2|+\rangle + i|-\rangle \\ |\psi_3\rangle &= |+\rangle - 2e^{\frac{-i\pi}{4}}|-\rangle \end{aligned}$$

- a) [1] Normalize each of the above states (following our convention that the coefficient of the  $|+\rangle$  basis ket is always positive and real.)
- b) [1] For *each of these three states*, find the probability that the spin component will be “up” along the Z-direction. Use bra-ket notation in your calculations!
- c) [1] For JUST the second state,  $|\psi_2\rangle$ , find the probability that the spin component will be “up” along the X-direction. Use bra-ket notation in your calculations. (*Hint: you will need McIntyre eq 1.36 for this one. You will need McIntyre 1.60 for the next part!*)
- d) [2] For JUST state  $|\psi_3\rangle$ , find the probability that the spin component will be “up” along the Y-direction. Use bra-ket notation in your calculations. *Be careful, there is some nasty complex-number arithmetic required on this one that is very important, and people frequently make mistakes on it that change the answer significantly!*

**6. [2] The physics of quantum states.**

Using postulate 4, show that the probability to obtain a particular measurement of spin is unaffected by changing a state  $|\psi\rangle$  to a state  $e^{i\beta}|\psi\rangle$ . Briefly discuss the significance of your result, what does it say to you? Also, how is it connected to the convention you imposed in Q3 and 4 above (setting the coefficient of the  $|+\rangle$  basis ket to be positive and real?)

**7. [1] Interpreting the coefficients in kets.**

Construct a quantum state for which the probability that it is spin up in the Z-direction is *three times* the probability that it is spin down. Is your answer unique?