

ASEN 5050 SPACEFLIGHT DYNAMICS Two-Body Motion

Prof. R. S Nerem
University of Colorado – Boulder

Lecture 3: The Two Body Problem

1

Announcements

- Homework #2 is due a week from today!
 - Either handed in or uploaded to D2L
 - Late policy is 10% per school day, where a “day” starts at 9:00 am.

Lecture 3: The Two Body Problem

2

Space News

Russian extends travel time to International Space Station

Posted August 26



MOSCOW — The Russian Federal Space Agency says the next manned trip to the International Space Station will be extended from the usual six hours to two days.

Roscosmos said in a statement released on Wednesday that the decision was made due to security concerns after the space lab had to adjust its orbit last month, dodging space junk.

The roll-out of a new Soyuz rocket in March 2013 allowed Russia to cut travel time to the orbiting lab from two days travel in cramped quarters to just six hours.

The next manned mission to the ISS is due to blast off from the Russia-leased launch pad in Kazakhstan on September 2, carrying a Russian, a Kazakh and a Dane.

Lecture 3: The Two Body Problem

3

Homework #1

- Darn orbital dynamics. The ISS is not cooperative at the moment. Here are the next passes (viewed from Boulder):

ISS - Visible Passes

Search period start: 03 September 2015 00:00

Search period end: 13 September 2015 00:00



Orbit: 400 x 403 km, 51.6° (Epoch: 01 September)

Passes to include: ☒ visible only ☐ all

Click on the date to get a star chart and other pass details.

Date	Brightness (mag)	Start			Highest point			End			Pass type
		Time	Alt.	Az.	Time	Alt.	Az.	Time	Alt.	Az.	
06 Sep	-0.4	05:49:25	10°	SSE	05:50:34	12°	SE	05:51:50	10°	ESE	visible
08 Sep	-1.5	05:37:22	10°	SSW	05:40:10	27°	SE	05:43:00	10°	ENE	visible
09 Sep	-0.7	04:46:09	12°	SSE	04:47:03	14°	SE	04:48:51	10°	E	visible
10 Sep	-3.1	05:28:00	23°	SW	05:29:53	67°	SE	05:33:07	10°	ENE	visible
11 Sep	-1.7	04:37:08	30°	ESE	04:37:08	30°	ESE	04:39:39	10°	ENE	visible
11 Sep	-2.1	06:10:10	10°	W	06:13:06	30°	NNW	06:16:02	10°	NE	visible
12 Sep	-3.0	05:18:48	37°	W	05:19:42	51°	NW	05:22:55	10°	NE	visible

Lecture 3: The Two Body Problem

4

Homework #1

ISS - Visible Passes

Search period start: 03 October 2015 00:00

Search period end: 13 October 2015 00:00

Orbit: 400 x 403 km, 51.6° (Epoch: 01 September)

Passes to include: ☒ visible only ☐ all

Click on the date to get a star chart and other pass details.

Date	Brightness (mag)	Start			Highest point			End			Pass type
		Time	Alt.	Az.	Time	Alt.	Az.	Time	Alt.	Az.	
03 Oct	-2.1	19:23:19	10°	SSW	19:26:03	25°	SE	19:28:03	15°	E	visible
03 Oct	-0.7	20:59:21	10°	W	21:00:41	20°	W	21:00:41	20°	W	visible
04 Oct	-3.0	20:05:48	10°	WSW	20:09:01	64°	NW	20:10:01	39°	NNE	visible
05 Oct	-3.3	19:12:33	10°	SW	19:15:46	63°	SE	19:19:00	10°	ENE	visible
05 Oct	-0.8	20:49:52	10°	WNW	20:51:50	20°	NW	20:51:50	20°	NW	visible
06 Oct	-1.7	19:56:02	10°	W	19:58:59	31°	NNW	20:00:54	17°	NNE	visible
07 Oct	-2.7	19:02:25	10°	WSW	19:05:37	54°	NNW	19:08:50	10°	NE	visible
07 Oct	-0.6	20:40:40	10°	NW	20:42:28	14°	NNW	20:42:28	14°	NNW	visible
08 Oct	-1.0	19:46:37	10°	WNW	19:49:04	19°	NNW	19:51:22	11°	NNE	visible
09 Oct	-1.6	18:52:43	10°	W	18:55:36	28°	NNW	18:58:28	10°	NE	visible
09 Oct	-0.4	20:31:35	10°	NNW	20:32:47	12°	N	20:32:47	12°	N	visible
10 Oct	-0.7	19:37:27	10°	NW	19:39:15	14°	NNW	19:41:01	10°	NNE	visible
11 Oct	-0.5	20:22:12	10°	NNW	20:22:56	11°	N	20:22:56	11°	N	visible
12 Oct	-0.6	19:28:22	10°	NNW	19:29:31	11°	N	19:30:39	10°	NNE	visible

Lecture 3: The Two Body Problem

5

Today's Lecture Topics

- Converting between the anomalies
- Then: More two-body orbital element computations

Our toolbox:

- Newton's law of gravitation
- Specific Energy
- Vis-Viva Equation

Most everything else comes straight from these.

Lecture 3: The Two Body Problem

6

Properties of Conic Sections

	\mathbf{v}	$\boldsymbol{\varepsilon}$	\mathbf{e}	\mathbf{a}
Ellipses	$< \sqrt{\frac{2\mu}{r}}$	< 0	$0 \leq e < 1$	$a > 0$
Parabolas	$= \sqrt{\frac{2\mu}{r}}$	$= 0$	$e = 1$	$a = \infty$
Hyperbolas	$> \sqrt{\frac{2\mu}{r}}$	> 0	$e > 1$	$a < 0$

\curvearrowright Since $\frac{h^2}{\mu} = a(1 - e^2)$ is positive

Lecture 3: The Two Body Problem

7

Orbital Period

- Last time we proved Kepler's 2nd and 3rd laws and arrived at an expression for the orbital period of an ellipse:

$$P = 2\pi \sqrt{\frac{a^3}{\mu}} \quad \text{units} = \sqrt{\frac{\text{km}^3}{\text{km}^3/\text{s}^2}} = \text{seconds}$$

Lecture 3: The Two Body Problem

8

Mean Motion

- We also arrived at an expression for the *mean motion*:

$$n = \frac{2\pi \text{ radians}}{P \text{ sec}} \qquad P = 2\pi \sqrt{\frac{a^3}{\mu}}$$

$$n = \frac{2\pi}{P} = \frac{2\pi}{2\pi \sqrt{\frac{a^3}{\mu}}}$$


$$= \sqrt{\frac{\mu}{a^3}} \text{ rad/s}$$

Lecture 3: The Two Body Problem

9

Where are we in an orbit?

- We have:
 - Our position in an orbit relative to Earth
 - The time
 - The true anomaly, v
- We want to know:
 - How long it will take to get somewhere
 - The time profile of $v(t)$
- We pose the answer by determining how much area is swept out in some amount of time.




Lecture 3: The Two Body Problem

10

The Anomalies

- The true anomaly, v
 - The actual, measured angle
 - Notice that this does not advance at a constant rate in an elliptical orbit
- Mean anomaly, M
 - An angle that *does* advance at a constant rate in an elliptical orbit.
$$M = n(t - t_p)$$
- Eccentric anomaly, E
 - An angle that helps translate from the true anomaly to the mean anomaly




Lecture 3: The Two Body Problem

11

The Anomalies

- The true anomaly, v
 - The actual, measured angle
 - Notice that this does not advance at a constant rate in an elliptical orbit
- Mean anomaly, M
 - An angle that *does* advance at a constant rate in an elliptical orbit.
$$M = n(t - t_p)$$
- Eccentric anomaly, E
 - An angle that helps translate from the true anomaly to the mean anomaly




Lecture 3: The Two Body Problem

12

The Anomalies


- Some quick mental exercises:



Lecture 3: The Two Body Problem

13

The Anomalies



Satellite is at periapse

True Anomaly = **0 deg**


Mean Anomaly = **0 deg**

Eccentric Anomaly = **0 deg**

Lecture 3: The Two Body Problem

14

The Anomalies



Satellite is at $v = 80$ deg $e = 0.3$

True Anomaly = **80 deg**


Mean Anomaly = **48 deg**

Eccentric Anomaly = **63 deg**

Lecture 3: The Two Body Problem

15

The Anomalies



Satellite is at $v = 120$ deg $e = 0.3$

True Anomaly = **120 deg**


Mean Anomaly = **87 deg**

Eccentric Anomaly = **104 deg**

Lecture 3: The Two Body Problem

16

The Anomalies



Satellite is at apoapse

True Anomaly = **180 deg**

Mean Anomaly = **180 deg**

Eccentric Anomaly = **180 deg**

Lecture 3: The Two Body Problem

17

The Anomalies

- The true anomaly advances quickly away from periapse
 - The mean anomaly advances steadily
 - The eccentric anomaly is in between
- At apoapse, they all catch up.
- Beyond apoapse, it's all in reverse (i.e., symmetric)
 - True anomaly advances the slowest away from apoapse
 - Mean anomaly advances steadily
 - Eccentric anomaly is in between

Lecture 3: The Two Body Problem

18

Let's see
some math!





Figure 2-2. Geometry of Kepler's Equation. The eccentric anomaly uses an auxiliary circle as shown. The ultimate goal is to determine the area, A_1 , which allows us to calculate the time. (Vallado, 2013)

Lecture 3: The Two Body Problem

19

True Anomaly
Mean Anomaly
Eccentric Anomaly

Note: $c = a - r_p$
 $= a - a(1-e)$
 $= ae$



For a coordinate system centered on Earth, write the location of the satellite in terms of E

$$X_{SAT} = a \cos E - ae$$

Eq. of Ellipse $\frac{X_{SAT}^2}{a^2} + \frac{Y_{SAT}^2}{b^2} = 1 \Rightarrow Y_{SAT} = b \sin E$

Lecture 3: The Two Body Problem

20

Derivation of Kepler's Equation

Now,

$$\begin{aligned}
 r^2 &= X_{\text{SAT}}^2 + Y_{\text{SAT}}^2 \\
 &= (a \cos E - ae)^2 + (b \sin E)^2 \\
 &= a^2 [\cos^2 E - 2e \cos E + e^2 + (1 - e^2) \sin^2 E] \\
 &= a^2 [1 - 2e \cos E + e^2 (1 - \sin^2 E)] \\
 &= a^2 [1 - 2e \cos E + e^2 \cos^2 E] \\
 &= a^2 (1 - e \cos E)^2 \quad \Rightarrow \quad r = a(1 - e \cos E)
 \end{aligned}$$

Now we will derive Kepler's Equation

Derivation of Kepler's Equation

Remember $r = \frac{p}{1 + e \cos \nu}$

$$\dot{r} = \frac{-p(-e \sin \nu) \dot{\nu}}{(1 + e \cos \nu)^2} = \frac{p(e \sin \nu) \dot{\nu} r^2}{p^2} = \frac{h e \sin \nu}{p}$$

But also $r = a(1 - e \cos E)$

$$\dot{r} = ae \sin E \dot{E}$$

So, $ae \dot{E} \sin E = \frac{h e \sin \nu}{p}$

$$\cancel{ae \dot{E} \sin E} = \frac{h \cancel{e}}{p} \left(\frac{b}{r} \cancel{\sin E} \right)$$

Note $b \sin E = r \sin \nu$

$$\sin \nu = \frac{b}{r} \sin E$$

Derivation of Kepler's Equation

$$\text{Thus, } r\dot{E} = \frac{hb}{pa} = \frac{\sqrt{\mu a(1-e^2)}a\sqrt{1-e^2}}{a(1-e^2)a} \quad \frac{h^2}{\mu} = p$$

$$r\dot{E} = \frac{\sqrt{\mu a a(1-e^2)}}{a^2(1-e^2)} = \sqrt{\mu} \frac{a^{3/2}}{a^{4/2}} = \sqrt{\frac{\mu}{a}}$$

$$\text{Thus, } a(1 - e \cos E)\dot{E} = \sqrt{\frac{\mu}{a}}$$

$$\dot{E} - e \cos E \dot{E} = \frac{\mu^{1/2}}{a^{3/2}} = n$$

$$\text{Integrating, } E - e \sin E = nt + \text{constant}$$

Derivation of Kepler's Equation

If reference time is perigee, $E = 0$,

$$\cancel{E}_{t_p} - e \sin \cancel{E}_{t_p} = nt_p + \text{constant} \Rightarrow \text{constant} = -nt_p$$

$$E - e \sin E = n(t - t_p) \equiv M \quad \text{Kepler's Equation}$$

*Angle swept out at the
mean angular velocity,
n, the mean motion*

$M = \text{Mean Anomaly}$

$$M = n(t - t_p) = E - e \sin E$$

Kepler's Equation relates the Mean
Anomaly to the Eccentric Anomaly

Derivation of Kepler's Equation

Can we relate either the Mean or Eccentric Anomaly to the True Anomaly?

Yes! $r = a(1 - e \cos E)$

$$r = \frac{p}{1 + e \cos \nu}$$

Useful Formula:

$$r = a(1 - e \cos E) = \frac{p}{1 + e \cos \nu} = \frac{a(1 - e^2)}{1 + e \cos \nu}$$

Lecture 3: The Two Body Problem

25

Conversions

- Any issues with these relationships?

$$M = n(t - t_p) = E - e \sin E$$

$$r = a(1 - e \cos E) = \frac{p}{1 + e \cos \nu} = \frac{a(1 - e^2)}{1 + e \cos \nu}$$

- Quadrants! Two satellites with $\nu = 10$ deg and $\nu = -10$ deg will be impossible to distinguish.

Lecture 3: The Two Body Problem

26

Quadrant Checks

- True Anomaly \rightarrow Eccentric Anomaly (Eq 2-9)

$$\sin(E) = \frac{\sin(\nu)\sqrt{1-e^2}}{1+e\cos(\nu)} \quad \cos(E) = \frac{e+\cos(\nu)}{1+e\cos(\nu)}$$

- Eccentric Anomaly \rightarrow True Anomaly

$$\sin(\nu) = \frac{\sin(E)\sqrt{1-e^2}}{1-e\cos(E)} \quad \cos(\nu) = \frac{\cos(E)-e}{1-e\cos(E)}$$

(both ν and E are either 0 – 180 deg or 180 – 360 deg)

Lecture 3: The Two Body Problem

27

More Quadrant Checks

- True Anomaly \rightarrow Eccentric Anomaly (Eq 2-14)

$$\tan\left(\frac{E}{2}\right) = \sqrt{\frac{1-e}{1+e}} \tan\left(\frac{\nu}{2}\right)$$

- Eccentric Anomaly \rightarrow True Anomaly (Eq 2-13)

$$\tan\left(\frac{\nu}{2}\right) = \sqrt{\frac{1+e}{1-e}} \tan\left(\frac{E}{2}\right)$$

Lecture 3: The Two Body Problem

28

Derivation of Kepler's Equation

$$\text{So, } \cos v = \frac{\cos E - e}{1 + e \cos E} = \frac{a(\cos E - e)}{r}$$

$$\sin v = \sqrt{1 - \cos^2 v} = \frac{\sqrt{1 - e^2} \sin E}{1 + e \cos E} = \frac{a\sqrt{1 - e^2} \sin E}{r}$$

$$\tan^2 \frac{v}{2} = \frac{1 - \cos v}{1 + \cos v} = \frac{1 - \frac{\cos E - e}{1 + e \cos E}}{1 + \frac{\cos E - e}{1 + e \cos E}} = \frac{1 - e \cos E - \cos E + e}{1 - e \cos E + \cos E - e}$$

$$= \frac{(1 + e)(1 - \cos E)}{(1 - e)(1 + \cos E)} = \frac{1 + e}{1 - e} \tan^2 \frac{E}{2}$$

$$\tan \frac{v}{2} = \sqrt{\frac{1 + e}{1 - e}} \tan \frac{E}{2}$$

Lecture 3: The Two Body Problem

29

If we have v and need more:

- If v is given:

$$r = \frac{p}{1 + e \cos v}$$

$$\cos E = \frac{r \cos v + ae}{a}$$

$$\sin E = \frac{r \sin v}{b}$$

Solve $E - e \sin E = M$ for t (or M)

Lecture 3: The Two Body Problem

30

If we have t and need more:

- If t is given:

$$M = n(t - t_p)$$

$$\text{Solve } E - e \sin E = M \quad \text{for } E$$

$$r = a(1 - e \cos E)$$

$$\text{and } \cos v = \frac{a \cos E - ae}{r}$$

$$\begin{aligned} \sin v &= \frac{b}{r} \sin E \\ &= \frac{a\sqrt{1-e^2}}{r} \sin E \end{aligned}$$

Lecture 3: The Two Body Problem

31

Derivation of Kepler's Equation

If t is given:

$$M = n(t - t_p)$$

$$\text{Solve } E - e \sin E = M \quad \text{for } E$$

$$r = a(1 - e \cos E)$$

$$\text{and } \cos v = \frac{a \cos E - ae}{r}$$

$$\begin{aligned} \sin v &= \frac{b}{r} \sin E \\ &= \frac{a\sqrt{1-e^2}}{r} \sin E \end{aligned}$$

If v is given:

$$r = \frac{p}{1 + e \cos v}$$

$$\cos E = \frac{r \cos v + ae}{a}$$

$$\sin E = \frac{r \sin v}{b}$$

$$\text{Solve } E - e \sin E = M \quad \text{for } t \text{ (or } M)$$

Another Useful Relation:

$$\tan \frac{v}{2} = \sqrt{\frac{1+e}{1-e}} \tan \frac{E}{2}$$

Lecture 3: The Two Body Problem

32

Solving Kepler's Equation

Want to use Newton-Raphson Iteration.


Assume we want to solve $f(y) = 0$ for y .

Assume $y = x + \delta$, where x is an approximate guess, and δ is a small correction. Expanding in a Taylor Series:

$$0 = f(y) = f(x + \delta) \approx f(x) + f'(x)\delta + \frac{f''(x)}{2!}\delta^2 + \dots$$

Neglecting 2nd order terms and higher, we can solve for δ :

$$\delta = -\frac{f(x)}{f'(x)}$$



Lecture 3: The Two Body Problem

33

Solving Kepler's Equation

We need to iterate this since we have neglected higher order terms:

$$X_{n+1} = X_n + \delta_n = X_n - \frac{f(X_n)}{f'(X_n)} \quad \begin{array}{l} \text{Iterate until } \delta_n \text{ is} \\ \text{acceptably small} \\ (\delta = y - x) \end{array}$$

We want to solve $E - e \sin E = M \Rightarrow f(E) = M - E + e \sin E = 0$
 $f'(E) = -1 + e \cos E$

So

$$E_{n+1} = E_n + \frac{M - E_n + e \sin E_n}{1 - e \cos E_n}$$

Careful with $e = 1$, i.e., don't use for high e .

Lecture 3: The Two Body Problem

34