

# saopaulo

May 7, 2021

## 1 Multilayer food webs

### 1.0.1 backward or forward?

### 1.1 ---

### 1.2 VI Fritz Müller Seminar Series

#### 1.2.1 Rodrigo Cogni,

#### 1.2.2 Paulo R. Guimarães Jr,

#### 1.2.3 Tiago B. Quental,

#### 1.2.4 Patricia P. Coltri

### 1.3 Thanks! Obrigado! Gracias!

#### 1.3.1 References

##### Biogeography

- Hutchinson, G. (1957). Concluding remarks. Cold Spring Harb. Symp. Quant. Biol., 22: 415-427.
- Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: ecological and evolutionary perspectives. PNAS, 106:19659-19665.
- Blonder, B. et al (2014). The N-dimensional hypervolume. Global Ecology and Biogeography, 23:595-609.
- Blonder, B. (2017). Hypervolume concepts in niche and traitbased ecology. Ecography, 41:1441-1455.
- Barros, C. et al. 2016. Ndimensional hypervolumes to study stability of complex ecosystems. Ecol. Lett., 19:729-742.

**Multivariate selection** \* Lande (1980). The genetic covariance between characters maintained by pleiotropic mutations. Genetics, 94:203-215. \* Melo and Marroig (2015). Directional selection can drive the evolution of modularity in complex traits. PNAS, 112:470-475. \* Goswami et al (2015). The fossil record of phenotypic integration and modularity: A deep-time perspective on developmental and evolutionary dynamics. PNAS, 112:4891-4896. \* Assis et al. (2020). Genetic correlations and ecological networks shape coevolving mutualisms. Ecology Letters 23:1789-1799.