Thermal Change, Network Fragmentation and Biotic Structure in a Mediterranean Fish Metacommunity

Final integrated report for the Guadalquivir River Basin 2026–2045 simulation horizon

Corrected October 2026 release

October 2026

Contents

1 Abstract **3**

2 Introduction **4**
2.1 Metacommunities in dendritic river networks 4
2.2 The Guadalquivir Basin as a model system 4
2.3 Objectives 5

3 Materials and Methods **5**
3.1 Study domain and assemblage 5
3.2 Local population dynamics 6
3.3 Dendritic dispersal and obstacles 7
3.4 Temperature forcing and initial conditions 7
3.5 Experimental design 8
3.6 Response metrics and analysis 8

4 Results **9**
4.1 Climate forcing and aggregate community response 9
4.2 Post-spin-up validation against the 2006–2009 survey 10
4.3 Brown trout as a thermal–hydraulic sentinel 11
4.4 Hydraulic controls and the passability effect 12
4.5 Interaction structure and native–invasive redistribution 12
4.6 Other species and aggregate indicators 13
4.7 Phase 2 re-run and structural sensitivities 13

5 Discussion **15**
5.1 Warming is weak at the aggregate scale and dominated by the transient 15
5.2 The trout signal is real but mostly transient 15
5.3 Connectivity is context dependent 15
5.4 Interaction structure is a modest structural uncertainty 16
5.5 What the structural sensitivities do and do not change 16
5.6 Monitoring and management implications 16

6 Limitations and Scope of Inference **17**

7 Conclusions **19**

8 Acknowledgements **20**

9 Data and Code Availability **20**

10 Figures **22**

11 Summary Tables **38**

12 Data Sources **39**

1 Abstract

Mediterranean river fish communities experience warming, dendritic habitat fragmentation and biological invasion simultaneously. Their responses cannot be inferred from climate forcing alone because persistence and thermal tracking depend on network connectivity and biotic interactions. We analysed a deterministic, spatially explicit metacommunity model of the Guadalquivir River Basin, southern Spain. The model represented 775 sites, 24 fish species, daily site-level water temperature, species-specific thermal performance, asymmetric interactions and directional dispersal through a reconstructed river-network tree. We synthesised a climate ensemble of 44 combinations of 11 general circulation models and four Shared Socioeconomic Pathways, a no-warming control, a 32-run obstacle factorial experiment and a 48-run interaction-structure experiment. All results in this release are recomputed from the Phase 2 post-fix model outputs (corrections C1–C7, E1–E22, applied to the regenerated interaction table); they supersede the September 2026 values, and the parser-fix, competition-structure and passability-floor sensitivities are reported in Section 4.7.

The corrected basin-mean realised warming over 2026–2045 averaged 0.54 °C under SSP1-2.6, 0.64 °C under SSP2-4.5, 0.62 °C under SSP3-7.0 and 0.70 °C under SSP5-8.5, with individual climate-model combinations spanning 0.21–1.09 °C; the 2036–2045 mean warmed 0.60–0.81 °C by pathway (0.18–1.29 °C across runs). Aggregate community metrics responded only weakly to this forcing. Over the 44 scenario runs, end-of-2045 total biomass changed by +0.075 model units per °C of realised warming ($R^2 = 0.14$), native richness by −0.0053 species per site per °C ($R^2 = 0.06$) and the deterministic realised richness loss by −0.00067 per °C ($R^2 = 0.01$). A large common transient dominates the raw trajectories: in the no-warming control, basin total biomass fell 35.5% and native biomass 40.8% between the 2026 observed initial state and end-of-2045, while native richness rose slightly. Climate effects are therefore reported as scenario-minus-control differences.

Brown trout (*Salmo trutta*) declined strongly in absolute terms under the interim projection route (control −37.3%; scenarios −38.4 to −38.8%), but only a small part of that decline is warming-attributable: scenario-minus-control losses were 1.1–1.5 percentage points, and trout relative biomass declined by 1.76 percentage points of baseline per °C ($R^2 = 0.63$). Exposure above the 20 °C upper limit at the 72 baseline-established trout sites had a baseline-climatology maximum of 90 days per year, while the 2045 scenario maxima were 80–135 days (mean of the per-run maxima 109 days) and the mean exposure rose from 23 to 45 days. In the obstacle experiment, upstream movement cost was the leading control of total and native biomass and of native richness (balanced within-design sums of squares 0.666, 0.729 and 0.935 respectively; the descriptive mean rank was 1.88 of seven factors), whereas passability dominated invasive biomass (SS = 0.741) and trout biomass (SS = 0.951); the climate endpoint was a minor factor (SS $\leq$ 0.015). The passability sign reversal reported in the September release is *not* recovered: with the corrected interaction direction and river-network graph, blocking passability raised basin-mean native biomass at every upstream cost and both endpoints, and improving passability lowered it. The invasive-favouring interaction matrix produced only a small redistribution relative to the original matrix (native biomass −1.7%, invasive biomass +3.6%, trout −1.6%).

Over this early-warming, twenty-year horizon the community indicators are controlled much more strongly by network hydraulics and by the uncalibrated transient than by the roughly 0.6 °C of realised warming. The results are a conditional, relative, deterministic sensitivity assessment rather than an absolute forecast, and are bounded by the interim and uncalibrated projection route, the symmetric distributional thermal niches, the uniform barrier-passability assumption, the same-day air–water calibration and the competition structure.

2 Introduction

2.1 Metacommunities in dendritic river networks

Riverine metacommunities are structured by the interaction of local environmental filtering, demographic regulation, biotic interactions and dispersal through branching networks. The metacommunity framework motivates this synthesis [Leibold et al., 2004].

Dendritic topology constrains movement to connected channels and makes directional movement important [Altermatt, 2013]. Barriers can alter both the amount of movement and the spatial distribution of demographic rescue and colonisation [Perkin and Gido, 2012]. The effect of a barrier is therefore expected to depend on its position in the network, the background cost of upstream movement and the species using the corridor.

Mediterranean rivers are especially informative systems for examining these interactions. They contain range-restricted endemic fishes and strong spatial gradients in temperature, habitat quality and flow regulation. Warming can move species away from or towards their thermal optima, while fragmentation can prevent populations from tracking suitable reaches. Invasive species can change the direction and magnitude of these responses through asymmetric competition and predation.

2.2 The Guadalquivir Basin as a model system

The Guadalquivir River Basin is a large, strongly regulated Mediterranean basin in southern Spain with high Iberian freshwater endemism and a long history of dam-associated fish-community change [Herrera et al., 2022, Ramos-Merchante and Prenda, 2018]. The model domain contains 775 sampling sites and 289 water-body reporting units (288 assigned water bodies plus one unassigned site, 1.2.2) grouped into 77 sub-catchments. The network connecting the sites is the corrected on-path river-network tree (correction C4): it has 774 edges (a tree on the 775 sites), a total length of 13,871.5 km and four parent-elevation inversions, against 31,206.3 km and 238 inversions in the legacy graph. The 774-edge link count is a network quantity and must not be read as a water-body count. The modelled assemblage contains 10 native residents, 11 exotic/invasive species and three migratory or diadromous taxa. Migratory taxa remain in the dynamical system and contribute to total biomass, but are excluded from native-richness reporting because their estuarine or marine life histories are not represented by the modelled freshwater corridor.

The root-join reconstruction is quantified in the corrected geometry. Of the 774 edges, 508 are on-path parent edges (8,636.9 km, 62.3% of total length) and 266 are minimum-spanning-tree root joins (5,234.5 km, 37.7%). The root joins connect 267 roots (34.5% of the 775 sites) and have a median length of 15.8 km and a maximum of 206.6 km. Of the 1,036 matched obstacles, 341 matched (245 applied) lie on a root-join edge; a broader "incident to a root site" count gives 531 matched (392 applied), but that also includes parent edges entering a root. Compounded passability is extreme on a few links: 330 links carry at least one obstacle, one link carries 33 obstacles and therefore an upstream factor of 1.0×10^{-33}, 28 links carry five or more obstacles, and 16 links fall below 10^{-6}. The root joins add 41 elevation inversions to the 4 on parent edges. The root count is sensitive to the on-path tolerance: 440 roots at $(0.01, 100$ m), 267 at $(0.02, 200$ m) and 119 at $(0.05, 500$ m); the simulations use the middle setting. These values are recomputed by `scripts/network_root_join_diagnostics.jl`.

The basin contains a dense inventory of transverse structures. Of the 1,658 non-completely-passable obstacles in the 2026 inventory, 973 were located within 2,000 m of the original river-network graph and 1,036 within 2,000 m of the corrected on-path tree used for the simulations (Fig. 15); the 63-obstacle difference is attributable to the C4 graph reconstruction, and the simulations use the corrected count. We use a 2,000 m crosswalk radius rather than assigning every structure to its nearest segment. The obstacle inventory and the 775-site network have

independent positional uncertainties: the obstacles are field points whose coordinates may carry coordinate-reference, GPS or digitisation error, the sampling sites are point locations that need not coincide with a vertex of the simplified network, and the reconstructed tree is a 775-vertex skeleton rather than a surveyed centreline. A 2 km corridor absorbs the resulting offsets—including coordinate-reference differences, GPS and digitisation error, and the offset between a site and the network segment that represents its reach—while reducing the risk that a structure is snapped to a non-adjacent tributary or across a sub-catchment boundary. A nearest-segment rule with no cap would assign every structure somewhere, including structures tens of kilometres from the modelled corridor, and would increasingly attach them to the wrong edge. The radius was fixed a priori and not tuned to the results. Figure 15 shows that 2 km sits at the knee of the distance distribution: the marginal inclusion rate falls from about 240–290 obstacles per km between 1 and 2 km to about 84–104 per km between 2 and 5 km. Beyond 2 km the curve flattens further, with 1,285–1,288 obstacles within 5 km, 1,488–1,491 within 10 km, and all 1,658 only at 227 km (on-path) or 295 km (legacy). Obstacles outside the radius are not treated as absent from the basin: they remain in the inventory but are not assigned a passability effect on the modelled network. Matched links have base upstream passability of 0.1 and downstream passability of 0.5, and barrier passabilities accumulate multiplicatively along a link (correction E12). A global upstream movement cost additionally reduces dispersal as elevation gain increases. These features make the basin a useful system for examining thermal–hydraulic coupling in a regulated Mediterranean river network [Perkin and Gido, 2012, Larsen et al., 2021].

2.3 Objectives

This report addresses three questions:

1. How strongly does projected warming between 2026 and 2045 alter aggregate biomass and richness relative to a stationary seasonal control?

2. How do upstream movement cost and local obstacle passability influence native, invasive and total biomass across the reconstructed river network?

3. Does the structure of biotic interactions alter apparent sensitivity to climate and connectivity, and which species provides the earliest indication of thermal change?

The analysis uses three complementary experiments: a multi-model climate ensemble, a balanced obstacle factorial and a deliberately contrasting interaction-structure sweep. The distinction between robust mechanistic results and effects conditional on perturbed model structure is maintained throughout.

3 Materials and Methods

3.1 Study domain and assemblage

The model represents one Guadalquivir basin with 775 sites, 289 water-body reporting units (288 assigned water bodies plus one unassigned site, 1.2.2) and 77 sub-catchments, covering 24 fish species. Native residents are *Aphanius baeticus*, *Anaecypris hispanica*, *Squalius pyrenaicus*, *Pseudochondrostoma willkommii*, *Luciobarbus sclateri*, *Squalius alburnoides*, *Iberochondrostoma lemmingii*, *Cobitis paludica*, *Iberochondrostoma oretanum* and *Salmo trutta*. Exotic/invasive species are *Gambusia holbrooki*, *Micropterus salmoides*, *Lepomis gibbosus*, *Cyprinus carpio*, *Carassius gibelio*, *Ameiurus melas*, *Oncorhynchus mykiss*, *Esox lucius*, *Gobio lozanoi*, *Tinca tinca* and *Alburnus alburnus*. Migratory or diadromous taxa are *Anguilla anguilla*, *Liza ramada* and *Mugil cephalus*. The 774 network edges are a separate quantity from the 289 water-body reporting units.

Alburnus alburnus (code AAL) is an exotic (EX) species and is grouped with the invasive/exotic taxa, not with the migratory taxa. Bleak does move upstream to spawn, but these movements are more limited than those of the native barbel (*Luciobarbus sclateri*), which is retained as a native resident; AAL is therefore not treated as diadromous. Reclassifying AAL raised the end-of-2045 invasive richness by 0.036 species per site and invasive biomass by 0.044 model units relative to the former classification in the corrected state (Section 4.6), and the Phase 2 post-fix values are 0.761 species per site and 4.94 model units; native, trout and total-biomass results are unchanged because the species groups affect reporting, not the population dynamics. The interaction and dispersal trait data for AAL remain incomplete, and its dispersal scalar is still a placeholder; this affects the absolute level of its density but not the native or hydraulic rankings.

3.2 Local population dynamics

For site i and species s, density $N_{i,s}$ follows the corrected competitive Lotka–Volterra form (corrections C2), with the interaction term inside the intrinsic-growth bracket:

$$\frac{dN_{i,s}}{dt} = N_{i,s}\left[r_s W_{i,s}\left(1 - \frac{\sum_j N_{i,j}}{K_i} + \sum_j c_{sj}\frac{N_{i,j}}{K_i}\right) - m_{i,s}^{\mathrm{heat}}(t)\right] + \sum_{j\in \mathrm{nb}(i)}\left(m_{ji}^{s} N_{j,s} - m_{ij}^{s} N_{i,s}\right), \tag{1}$$

where $c_{sj} = 1 - \alpha_{sj}$ and α_{sj} is the signed effect of species j on species s (negative α denotes a competitive or predatory effect), read in the correct direction from the interaction table (correction C1). K_i is site carrying capacity, r_s is the intrinsic growth rate and $W_{i,s}$ is environmental suitability. Carrying capacity was scaled to 1 times observed total density with a minimum-capacity floor. Annual literature-derived rates were converted to daily rates. The six team biological options were set to the corrected re-run values: absent species grow at the full rate (`absence_growth_fraction` = 1.0), isolated-pool capacity is capped at the non-pool median (`pool_capacity_mode` = cap, 84 sites), fishless *DIFICIL* sites are transit-only (`fishless_dificil_capacity` = near_zero, 56 sites), non-reproducing diadromous taxa have zero local growth (`nonreproducing_local_growth` = zero), the four Guadiato fish-farm eel records are excluded, and the salinity envelope is applied (correction E18); see `parameters_climate_scenarios_`

Environmental suitability was defined as

$$W_{i,s}(t) = \exp\left[-\frac{(T_i(t) - T_s^{*})^2}{2\sigma_s^2}\right] h_i, \tag{2}$$

where $T_i(t)$ is daily site temperature, T_s^{*} is the species thermal optimum, σ_s is thermal breadth and h_i is a static habitat-suitability index bounded between 0.1 and 1.0. The habitat index is a rescaled bank-stability index derived by an inverse, normalised transform of the field variable IET (*Índice de estabilidad del talud*), for which higher values mean a less stable bank. The model uses IET only as a monotone habitat proxy, not as a trophic-state or water-quality index, and the index is held constant. Thermal optima were set to the midpoint of each empirical temperature range and thermal breadth to one-sixth of that range ($T^{*} = 12.0$ °C and $\sigma = 2.67$ °C for brown trout, whose empirical range is 4–20 °C). The model therefore uses a fixed, symmetric Gaussian niche with no acclimation, plasticity or evolution (see C6 in the Limitations).

Heat stress was represented separately:

$$m_{i,s}^{\mathrm{heat}}(t) = k\,\max\left(0, T_i(t) - T_s^{\mathrm{up}}\right)^2, \tag{3}$$

where T_s^{up} is the empirical upper thermal limit and the shared coefficient k was recalibrated from the corrected baseline (correction C5) so that the most exposed species–site combination

under baseline climate loses no more than 5% of annual survival. No cold-stress mortality term was applied.

3.3 Dendritic dispersal and obstacles

Movement from site i to site j was defined as

$$m_{ij} = D_{\text{median}} \frac{x_{ij} p_{ij}}{d_{ij}}, \tag{4}$$

where $D_{\text{median}} \approx 0.0274$ km day^{-1}, d_{ij} is hydrological distance along the reconstructed tree, x_{ij} is the elevation-dependent movement factor and p_{ij} is directional passability. Species-specific rates were obtained by multiplying this base rate by literature-derived relative dispersal scalars.

Downstream or level movement was assigned $x_{ij} = 1$. For upstream movement,

$$x_{ij} = \frac{1}{1 + c\Delta e}, \tag{5}$$

where c is global upstream movement cost and Δe is elevation gain. The corrected network is an on-path river-network tree reconstructed from the network-distance matrix (on-path parent rule plus an explicit minimum-spanning-tree root join; correction C4). Obstacles are snapped to this tree and their passabilities accumulate multiplicatively along a link rather than taking the element-wise minimum (correction E12); non-operational and ambiguous structures are classified and excluded or annotated. A sparse dispersal matrix was recomputed for each upstream-cost and passability combination.

The four passability states were baseline (multiplier 1.0), improved (1.5), reduced (0.5) and blocked (0.1), with passability capped at 1.0. No exploitation-pressure scenario was applied in the experiments reported here.

3.4 Temperature forcing and initial conditions

Climate simulations used daily site-level water temperature for 11 general circulation models under SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5. Site temperature is the corrected bias-corrected `guadex_tw` daily water-temperature series (corrections C5, E1, E22): the modelled level is the corrected baseline `tw_baseline_mean` for 1986–2005 and the daily anomaly is added to it, so the model temperature equals the corrected daily series. The corrected basin-mean baseline site temperature across 775 sites is 15.82 °C (2026 annual means; range 9.79–18.06 °C). The no-warming control repeated the baseline seasonal climatology.

The projections follow the C3 interim route: they start from the observed 2006–2009 community and advance through a fixed three-year baseline spin-up, with no calibration of growth, capacity, interaction or dispersal parameters. The system was integrated with the adaptive fifth-order Runge–Kutta method `Tsit5` at relative and absolute tolerances of 10^{-6}, with one time unit equal to one day and a positivity callback. Reporting is end-of-year (correction E5), so the value labelled 2045 is the state at $t = 7300$ days.

Two warming quantities are kept distinct. The dose–response analysis uses the *realised* applied warming of each run (`realised_warming_2026_2045_mean_degc` and `realised_warming_2036_2045_mean` mean over sites; correction C7), not the imposed basin warming curve. The obstacle experiment used the coolest (SSP1-2.6/IITM-ESM) and warmest (SSP2-4.5/UKESM1-0-LL) runs of the corrected ensemble. Endpoints were selected on the **2036–2045** realised window, where they are 0.180 and 1.292 °C; their 2026–2045 means over the same runs are 0.208 and 1.091 °C. Both windows are named explicitly so a value is never quoted without its window.

3.5 Experimental design

Table 1: Corrected simulation design.

Component	Specification
Spatial domain	Guadalquivir basin; 775 sites, 289 water-body reporting units (288 assigned + 1 unassigned) and 77 sub-catchments; on-path river-network tree with 774 edges
Assemblage	24 species: 10 native residents, 11 exotic/invasive and 3 migratory/diadromous
Climate ensemble	11 climate models × 4 SSPs = 44 scenarios plus one no-warming control (45 runs)
Obstacle experiment	2 climate endpoints × 4 upstream costs × 4 passability states = 32 runs
Interaction experiment	3 interaction matrices × 2 climate endpoints × 2 upstream costs × 4 passability states = 48 runs
Upstream costs	0.01, 0.05, 0.10 and 0.50 in the obstacle experiment; 0.01 and 0.50 in the interaction experiment
Interaction matrices	Original qualitative matrix; random placebo with off-diagonal values in $[-1, 0]$; invasive-favouring worst-case matrix
Projection route	C3 interim: observed start + 3-year baseline spin-up, no calibration; reported as scenario − control
Simulation horizon	2026–2045 with daily temperature forcing and end-of-year reporting

The original interaction matrix maps qualitative categories to coefficients (-1.0 no coexistence, -0.8 displacement, -0.5 predation, -0.3 competition or interference, -0.2 weak effect, 0.0 neutral), now parsed in the correct direction and with undirected competition made symmetric (C1). The random placebo used a fixed seed of 42. All alternative-interaction runs used a thermal-breadth multiplier of 0.3, so interaction structure is confounded with thermal breadth in that sweep; this is noted in the Limitations.

3.6 Response metrics and analysis

Richness was the number of species exceeding 0.1 density units, averaged over sites or reporting units. Native, invasive and total biomass were sums of the relevant species densities. Realised native richness loss was calculated as

$$L(t) = \max\left(0, 1 - \frac{R(t)}{R_0}\right), \tag{6}$$

where R_0 is richness in the spun-up baseline. This is a deterministic realised richness contraction; it is not an extinction probability (the model is deterministic and has no stochastic replicates; correction E8). The quasi-extinction diagnostic flags a site when a species remained below max(0.1, 0.1× baseline density) for three consecutive annual observations, evaluated only over sites at which the species was baseline-established (correction E6); it is not used as a primary climate outcome.

All headline numbers in this report were recomputed from the Phase 2 post-fix output files on 2026-11-01 with the following estimators:

- **Warming statistics**: mean/min/max of `realised_warming_*_mean_degc` by pathway and across all 44 scenario runs (`results/climate_scenarios_phase2/runs_index.csv`).
- **Dose–response**: ordinary least squares of the end-of-2045 basin metric on realised 2026–2045 mean warming across the 44 scenario runs, excluding the control; R^2 is the coefficient

of determination of that fit. Final-year basin metrics are from each run's `export/levels/level_basin.csv` (year 2045) and are reproduced in `runs_index.csv`.

- **Brown trout**: per-run relative biomass change and baseline/final biomass from each run's `export/levels/quasi_extinction_summary.csv` (species ST); the "per °C" slope is OLS on relative change. Scenario-minus-control differences use the shared no-warming control.
- **Exposure**: `mean/median/max_exposure_days` for ST over the `baseline_established` site set (72 sites) in each run's `export/levels/exposure_summary.csv`, year 2045.
- **Obstacle sweep**: factor ranking from `results/sensitivity_obstacles_phase2/report_plots/parame` (median within-block span in run-set standard-deviation units; rank 1 = largest span, averaged over the eight response metrics); sums of squares from `report_plots/variance_decomposition.csv` (balanced within-sweep fractions). Passability effects are blocked/improved/reduced minus baseline at the same cost and endpoint.
- **Interaction sweep**: per-matrix means and passability spans from `results/alt_interactions_phase2/r` sums of squares from the alternative-sweep rows of the shared `variance_decomposition.csv`.

No formal significance testing was performed and no stochastic replicate simulations were run. Biomass is expressed in model units and is not calibrated field biomass.

4 Results

4.1 Climate forcing and aggregate community response

Added basin-mean realised warming over 2026–2045 averaged 0.542 °C under SSP1-2.6, 0.643 °C under SSP2-4.5, 0.619 °C under SSP3-7.0 and 0.701 °C under SSP5-8.5 (Table 3). Across individual climate-model and pathway combinations the 2026–2045 mean spanned 0.208–1.091 °C. For the later decade 2036–2045 the pathway means were 0.598, 0.729, 0.714 and 0.814 °C, and the run span was 0.180–1.292 °C. Climate-model spread overlapped pathway differences: in the 2026–2045 window, for example, the warmest SSP1-2.6 run (0.809 °C) exceeded the coolest SSP5-8.5 run (0.443 °C).

The control was not stationary in the raw trajectories. Between the 2026 observed initial state and end-of-2045 the no-warming control lost 35.5% of total biomass and 40.8% of native biomass while native richness rose 3.0% and invasive biomass rose 13.5%; these changes are the residual transient of the interim route and are common to scenario and control. Scenario effects are therefore read as scenario-minus-control differences, which are small: pathway-mean end-of-2045 total biomass deltas relative to the control were +0.008 (SSP1-2.6), +0.035 (SSP2-4.5), +0.016 (SSP3-7.0) and +0.030 (SSP5-8.5) model units per site, and pathway-mean native-richness deltas were +0.0026, +0.0002, +0.0019 and +0.0034 species per site.

Across the 44 scenario runs, the end-of-2045 aggregate responses to realised warming were weak (Table 3). Total biomass *increased* with warming, at +0.075 model units per °C ($R^2 = 0.14$); native richness declined at −0.0053 species per site per °C ($R^2 = 0.06$); and the deterministic realised richness loss declined at −0.00067 per °C ($R^2 = 0.01$). The correction therefore reverses the sign and weakens the total-biomass slope reported in September and removes the previously reported positive richness-loss slope. The reasons are the corrected bias-corrected water temperature, the corrected interaction form and, above all, the interim route: the aggregate trajectories are dominated by a transient common to all runs, leaving only a small differential climate signal at 2045. Relative to the pre-fix corrected outputs, the parser fix moves the native-richness slope from −0.0041 to −0.0053 and the richness-loss slope from −0.00065 to −0.00067, and leaves the total-biomass slope unchanged at the reported precision; the interpretation is unaffected (Section 4.7).

Most dominant native cyprinids have optima of 19–20 °C against a corrected basin-mean baseline of 15.82 °C, so modest warming moves them towards their optima. Brown trout has an optimum of 12.0 °C; the mean corrected baseline water temperature at the 72 baseline-established trout sites is 13.72 °C, about 0.65 thermal standard deviations above the optimum. Aggregate metrics consequently conceal opposing species responses.

E21 warming uncertainty band. The primary air–water warming relationship is a same-day calibration (slope ≈ 0.510 across Spain, 0.436 for the Guadalquivir). Lagged air fits (3-, 7- and 30-day means) yield higher slopes, up to ≈ 0.65 Spain-wide, but rest on only about three in-sample lag-eligible sites ($n = 104$, one year of data), so a lagged refit cannot serve as a primary calibration. We therefore report the lagged relationship as a sensitivity band on the projected warming, spanning $\times 0.63$ to $\times 1.28$ of the central value (`guadex_tw/outputs/tables/air_water_slope_uncertainty_ban`). On the GuadeX network for the 2026–2045 horizon the central/ low/ high ensemble-median water warming by pathway is 0.580/0.368/0.741 °C (SSP1-2.6), 0.677/0.429/0.864 °C (SSP2-4.5), 0.647/0.411/0.826 °C (SSP3-7.0) and 0.730/0.463/0.931 °C (SSP5-8.5). Because the lagged slopes are lower than the same-day slope within the Guadalquivir subset, the upper end of this band may not apply to the basin, and the band should be treated as a conservative sensitivity envelope rather than a calibrated range (see Limitations 3).

4.2 Post-spin-up validation against the 2006–2009 survey

The interim route is only defensible if the three-year spun-up state resembles the survey it starts from. We compared the saved post-spin-up state of the no-warming control with the observed 2006–2009 density matrix, species by species, over all 775 sites (Table 2). Observed presence is a positive survey density; modelled presence is a spun-up density above the 0.1-unit reporting threshold; the true skill statistic is sensitivity + specificity -1. The abundance rank correlation is Spearman's ρ over the sites where the species was observed, and the last column is the change in that species' total density in the control between 2026 and 2045. Across the 24 species the pooled sensitivity is 0.975 and the pooled specificity 0.920.

The comparison is a mixed but interpretable result. Recall is high everywhere, but the model is systematically more widespread than the survey: for most species modelled occupancy exceeds observed occupancy, and the TSS of the widespread *Luciobarbus sclateri* is only 0.42 because 247 of its 424 survey-unoccupied sites are predicted as occupied. The high TSS of the rare species is therefore driven mainly by their many correctly empty sites, not by a good abundance fit. This over-prediction, and the weak control trajectory of several invasive species, is the acceptance baseline that full calibration must improve; it is reported as a finding, not treated as a validation success.[1]

The thresholds that will define a successful future calibration campaign are fixed in advance, and they are explicitly *future-calibration criteria*, not acceptance gates for this release: per-species TSS ≥ 0.40 where at least 10 sites are occupied, and $|\text{modelled} - \text{observed occupancy}| \leq 0.10$ for LS, SA, SP, PW and CP. Against these criteria the current spun-up control passes the TSS test where it applies (LS TSS = 0.417) but fails the occupancy test for every one of the five widespread natives, with LS over-predicted by 31.9 percentage points (45.3% observed against 77.2% modelled). The values in Table 2 are identical at the reported precision to the pre-fix validation, so the calibration gap is a property of the model structure, not of the parser repair.

[1] Reproduce with `julia -project=. scripts/validate_initial_state.jl`; the table is written to `results/climate_scenarios_phase2/validation/initial_state_validation.csv`.

Table 2: Post-spin-up validation against the 2006–2009 survey (no-warming control). Occupancies are percentages of the 775 sites; TSS is sensitivity + specificity −1; ρ is the abundance rank correlation over observed sites (- where fewer than three sites were occupied); ΔN is the control change in total density, 2026–2045.

Species	Obs. occ. (%)	Mod. occ. (%)	TSS	ρ	ΔN (%)
AA	1.2	1.3	0.55	0.80	0.0
AAL	1.4	4.3	0.97	0.89	−64.4
AB	0.1	0.4	1.00	–	4.2
AH	1.4	1.7	1.00	0.55	142.2
AM	0.1	0.3	1.00	–	−64.8
CC	5.2	12.6	0.74	0.61	−73.7
CG	1.8	5.8	0.89	0.48	−80.9
CP	15.7	30.3	0.82	0.89	17.9
EL	0.3	0.4	1.00	–	−75.6
GH	8.4	20.9	0.76	0.51	241.0
GL	0.5	1.9	0.99	1.00	15.3
IL	9.0	15.7	0.93	0.89	−38.9
IO	0.1	0.3	1.00	–	204.1
LG	5.8	18.6	0.82	0.60	−84.5
LR	0.6	3.4	0.97	0.90	0.0
LS	45.3	77.2	0.42	0.60	−45.9
MC	0.5	3.5	0.97	1.00	0.0
MS	2.5	7.1	0.84	0.75	−87.9
OM	1.5	4.0	0.89	0.88	13.7
PW	14.5	38.8	0.68	0.55	−66.9
SA	20.5	40.0	0.75	0.73	−44.2
SP	20.8	38.2	0.77	0.85	−38.1
ST	4.6	9.3	0.89	0.48	−32.4
TT	0.5	0.8	0.49	1.00	−63.2

4.3 Brown trout as a thermal–hydraulic sentinel

Under the interim route, basin brown-trout biomass declined from a spun-up baseline of 570.7 model units in every run. The decline is dominated by the common transient: the no-warming control fell to 358.0 units (−37.3%), and the scenario pathway means fell to −38.4% (SSP1-2.6), −38.5% (SSP2-4.5), −38.5% (SSP3-7.0) and −38.8% (SSP5-8.5) (Table 3). The warming-attributable component, taken as the scenario-minus-control difference, is small but consistently negative: −1.10 percentage points of baseline (SSP1-2.6), −1.23 (SSP2-4.5), −1.22 (SSP3-7.0) and −1.48 (SSP5-8.5). Across the 44 runs, trout relative biomass declined by 1.76 percentage points of baseline per °C of realised warming ($R^2 = 0.63$), and trout final biomass declined by 10.08 model units per °C ($R^2 = 0.63$).

Thermal exposure increased with warming. Over the 72 baseline-established trout sites, the maximum number of days per year above the 20 °C upper limit was 90 under the baseline climatology (control), but reached 80–135 across the 2045 scenario runs (mean of the per-run maxima 109.5). The per-run mean exposure rose from 23.4 days in the baseline climatology to a scenario mean of 45.4 days (range 30.2–65.7), and the median from 17.0 to 48.1 days (range 24.5–81.5). The first scenario year (2026) already shows elevated exposure (mean 42.7, max 110 days), a consequence of the pre-2026 offset between the projection products and the 1986–2005 baseline rather than of the simulated warming itself (Limitations 11). Trout occupies only about 9% of sites, so its large relative decline is diluted in basin-wide richness and biomass. It is an early-warning indicator of thermal–hydraulic coupling, not a proxy for whole-community vulnerability.

In the obstacle experiment, trout was controlled mainly by passability rather than by back-

ground climate: passability accounted for SS = 0.951 of final trout biomass, upstream cost SS = 0.031 and the climate endpoint SS = 0.015. At the cool endpoint, final trout biomass at upstream cost 0.05 was 355.4 units under baseline passability and 427.3 units under blocked passability (+71.9 units); at cost 0.50 the corresponding change was 353.3 to 423.4 units (+70.1). Blocked passability left more trout biomass in the basin at every upstream cost and both endpoints.

4.4 Hydraulic controls and the passability effect

The primary ranking is the balanced within-sweep sum-of-squares decomposition, because it is identified within matched factorial blocks. Upstream cost dominated total biomass (SS = 0.666), native biomass (SS = 0.729), native richness (SS = 0.935) and realised richness loss (SS = 0.970); passability dominated invasive biomass (SS = 0.741) and trout biomass (SS = 0.951); and the climate endpoint was minor for every metric (SS $\leq$ 0.015) (Table 4). As a descriptive cross-metric summary, the seven factors also rank by mean span (mean rank, 1 = strongest) as upstream cost 1.88 and passability 1.88 (tied), thermal breadth 2.50, interaction matrix 4.38, passability×upstream cost 4.75, climate endpoint 6.12 and passability×temperature 6.50. That descriptive ordering places the two hydraulic factors first, the thermal-breadth term third (but that term is confounded with the alternative-interaction sweep and is not independently identified), the interaction-matrix term fourth, the passability–cost interaction fifth, and the climate endpoint and passability–temperature term last.

The passability sign reversal described in the September release is not reproduced in the corrected sweep. For basin native biomass, blocked-minus-baseline differences at the cool endpoint were +1.672, +1.667, +1.724 and +1.912 model units at upstream costs 0.01, 0.05, 0.10 and 0.50, and at the warm endpoint +1.612, +1.586, +1.632 and +1.808. Blocking therefore raised basin-mean native biomass at every upstream cost and both endpoints, with no change of sign. Reduced passability behaved in the same direction (+0.85 to +1.01 units), and improved passability lowered basin native biomass at every cost and endpoint (−0.66 to −0.74 units). The same pattern held for total biomass (blocked-minus-baseline +1.78 to +2.08 units). The previously proposed explanation — that blocking restricted competitively effective invasive taxa — is also unsupported, because in the corrected matrix invasive species do not dominate native biomass and the invasive-favouring contrast is small (Section 4.5).

This is a context-dependent redistribution rather than a uniform benefit of blocking. The positive basin-level effect is a net of gains and losses among sub-catchments and water bodies, and the outputs cannot uniquely separate reduced native immigration, reduced emigration and suppressed invasive dispersal. The robust finding is that, under this model and this corrected network, a reduction in passability did not reduce aggregate native biomass over the 2026–2045 horizon.

4.5 Interaction structure and native–invasive redistribution

The corrected alternative-interaction sweep produced a much smaller contrast than the September release. Averaged over the 16 runs of each matrix, the original matrix gave basin native biomass 32.27, invasive biomass 2.97, trout biomass 445.7, native richness 2.61 and total biomass 35.39 model units; the random placebo gave 30.70, 2.63, 432.5, 2.56 and 33.48; and the invasive-favouring matrix gave 31.73, 3.08, 438.4, 2.59 and 34.95. Relative to the original matrix, the invasive-favouring matrix changed native biomass by −1.7%, invasive biomass by +3.6%, trout biomass by −1.6%, native richness by −0.5% and total biomass by −1.2%. In the matched cool-end, low-cost, baseline-passability contrast, native biomass changed from 33.07 to 32.42 units and invasive biomass from 2.76 to 2.89 units.

The interaction matrix accounted for SS = 0.116 of native biomass, = 0.161 of invasive biomass and = 0.034 of trout biomass in the alternative sweep, in each case well below upstream

cost (native biomass SS = 0.668; trout SS = 0.814). Passability spans were similar across matrices for both trout (mean span 29.1 units under original, 29.2 under random, 29.4 under invasive-favouring) and native biomass (1.40, 1.27, 1.30 units). The September statement that passability spans were large for trout only under the invasive-favouring matrix, and small for native biomass only under that matrix, is not reproduced: the spans are essentially matrix-independent. Because all alternative-interaction runs used a thermal-breadth multiplier of 0.3 and the three-year interim spin-up, the sweep identifies a structural sensitivity but the magnitudes are confounded and should not be read as an isolated interaction effect. The invasive-favouring matrix remains a bounding scenario, not an empirical uncertainty distribution.

4.6 Other species and aggregate indicators

Warm-adapted native cyprinids and most invasive and migratory taxa changed comparatively little over the study horizon in aggregate. With AAL now included in the exotic/invasive group, invasive richness at end-of-2045 was 0.761 species per site on average across scenarios (range 0.755–0.765) against 0.763 in the control, and invasive biomass was 4.94 model units on average (range 4.84–5.04) against 5.05 in the control. These values are far from the flat 0.217 reported in September: the corrected initial state and interaction form allow a higher invasive richness, although the value is dominated by a small number of established populations. The incomplete AAL trait data still make its absolute density provisional, but the values are no longer lower bounds on classification grounds. Native richness (2.64 species per site) and total richness were effectively invariant across scenarios and the control.

The brown-trout quasi-extinction fraction, now computed over the 72 baseline-established sites (E6), was 0.278–0.306 (mean 0.290) across the climate runs, 0.042–0.472 in the obstacle sweep and 0.056–0.444 in the alternative sweep. The near-saturation values (0.94–0.97) reported in September were an artefact of counting all 775 sites rather than only established sites; the corrected fractions are well below saturation and vary with passability and upstream cost rather than with climate.

4.7 Phase 2 re-run and structural sensitivities

The parser repair was quantified, not asserted. The interaction table was rebuilt after the corrected run. It still has 227 directed rows, but 34 directed rows change: 16 reverse-direction entries that had wrongly inherited -0.5 predation become -0.3 competition, and nine undirected "competition and predation" cells are newly flagged ambiguous on both directions (18 directed rows; the table now carries 31 ambiguous unordered pairs and 62 directed rows, against 22 and 44 before). The named direction of every pair is unchanged at -0.5. All three experiments were then re-run against the regenerated table (45 climate, 32 obstacle and 48 interaction runs), with the realised forcing verified bit-identical in all three, so each comparison isolates the change in the α matrix.

In the climate ensemble the parser fix moves the scenario-minus-control metric by at most 0.0026 native species of basin mean richness (2 of 775 sites), 0.0013 of realised richness loss and 0.0011 biomass units on a $\approx$ 30-unit mean ($\approx$ 0.004%). The delta keeps its sign in 42 of 45 richness runs, 42 of 45 richness-loss runs and all 45 biomass runs. In the obstacle sweep the largest changes are 0.0013 species, 0.0063 biomass units and 0.057 units of trout biomass, with zero sign changes and all 40 signed factor contrasts unchanged. In the interaction sweep the largest changes are 0.0013 species (one site, in 8 of 48 runs), 0.033 biomass units and 0.51 trout biomass units; the trout quasi-extinct fraction is bit-identical in all 48 runs, and the random-placebo runs are numerically identical to 10^{-15}, which shows that the intervening code changes are inert and that the empirical-matrix differences are the parser effect. The larger drift in the constructed invasive-favouring matrix is the $099a87d$ AAL reclassification, not the parser repair. The matrix ordering is unchanged in all 16 design blocks and the obstacle factor ranking has

Spearman $\rho = 1.00$. The repair is therefore material for the auditability of the α matrix and the implied $c = 1 - \alpha$ coefficients, and immaterial to every reported conclusion.

Competition structure. The primary specification keeps a shared per-site carrying capacity, with $c_{sj} = 1 - \alpha_{sj}$:

$$\frac{dN_{i,s}}{dt} = N_{i,s}\left[r_s W_{i,s}\left(1 - \sum_j \frac{N_{i,j}}{K_i} + \sum_j \alpha_{sj}\frac{N_{i,j}}{K_i}\right) - m_{i,s}^{\text{heat}}(t)\right], \qquad c_{sj} = 1 - \alpha_{sj}, \quad (7)$$

so every non-zero interaction gives $c_{sj} \in [1.3, 1.8]$, above self-limitation. The sensitivity replaces this with species-specific self-limitation and $c_{sj} = -\alpha_{sj}$ $(j \neq s)$, with no shared capacity:

$$\frac{dN_{i,s}}{dt} = N_{i,s}\left[r_s W_{i,s}\left(1 - \frac{N_{i,s}}{K_i} + \sum_{j \neq s} \alpha_{sj}\frac{N_{i,j}}{K_i}\right) - m_{i,s}^{\text{heat}}(t)\right], \qquad c_{ss} = 1, \quad c_{sj} = -\alpha_{sj}, \quad (8)$$

which gives $c_{sj} \in [0.3, 0.8]$, below self-limitation. The 45-run alternative was run over the same 44 scenarios and shared control. The *magnitude* of the weak-warming result is robust: the basin native-richness delta is at most 0.0129 species under either structure (≤ 10 of 775 sites), and the warm-endpoint delta is numerically identical at -0.006452 species. The *sign* of the ensemble-mean richness response, however, depends on the choice: shared-K gives a mixed, slightly positive mean (+0.0020; 26 positive, 4 zero, 14 negative), whereas the species-specific structure gives a uniformly negative response (44/44; mean -0.0071), and 30 of 44 runs flip sign. The small shared-K realised richness-loss signal collapses to approximately zero (mean -0.001551 to -0.000037; 31 of 44 alternative deltas exactly zero), while the total-biomass and trout responses amplify (mean absolute delta $0.036 \rightarrow 0.173$, about 4.8×; trout relative change $-0.0126 \rightarrow -0.0984$, about 7.8×; trout final biomass $-7.2 \rightarrow -59.4$, about 8.3×). The trout direction never flips (44/44 negative under both structures). The control states themselves differ between structures, so only the deltas are compared. All primary numbers in this report remain the shared-K results.

Passability floor and on-path-only obstacles. Two variants of the 32-run obstacle sweep test the reconstructed-network overlay. The first imposes a compounded-passability floor of 10^{-3} (the passability of three 0.1 barriers in series), which changes 56 unordered links in the baseline passability scenario and 112 under the reduced and blocked scenarios, and raises the extreme 10^{-33} link by 31 orders of magnitude. The second restricts obstacle snapping to on-path parent edges, so reconstructed root joins carry no barriers: of the 245 applied obstacles on 111 root-join links, 197 become unmatched and 48 re-snap onto parent edges (matched obstacles fall from 1,036 to 759, applied from 796 to 599). The factor order is identical in the baseline and both variants for all eight metrics (Spearman $\rho = 1.00$, top two factors unchanged in 8/8, no sign flips). The floor is a small perturbation: at most +0.0297 native species of basin richness (1.1%), 0.045 biomass units on ≈ 33, 0.50 trout biomass units on ≈ 427, and one trout quasi-extinction site (1/72). Parent-only is a larger magnitude perturbation — mean native richness +8.5% (up to +0.2968), total biomass −2.3%, invasive biomass −7.1% and trout quasi-extinction up to +0.1142 absolute — but leaves the ranking and the direction of every factor effect unchanged. The extreme link is 1.33.2–1.33.58, a *parent* edge with 33 obstacles (combined upstream dam factor 10^{-34}); parent-only leaves it untouched, while the floor raises it to 10^{-3}, and it contributes only 0.42–1.12% of the dispersal rate added by the floor. It is therefore an extreme value, not the driver of the passability conclusion.

Validation and calibration criteria. The validation table is regenerated from the Phase 2 shared control and written to `results/climate_scenarios_phase2/validation/initial_state_validation.` its values are identical at the reported precision to the pre-fix table (Table 2). The fixed acceptance thresholds for the future calibration campaign — per-species TSS ≥ 0.40 where at least 10 sites are occupied, and $|\text{modelled} - \text{observed occupancy}| \leq 0.10$ for LS, SA, SP, PW and CP

— are calibration targets, not gates for this release; the current control passes the TSS criterion where it applies (LS 0.417) but fails the occupancy criterion for all five widespread natives.

The per-GCM matched controls (11 runs driven by each GCM's own historical series) and the coded-value refit of the national air–water model remain deferred by the Phase 2 decisions and are stated in the Limitations; they are the two remaining structural caveats on the climate deltas.

5 Discussion

5.1 Warming is weak at the aggregate scale and dominated by the transient

The corrected result changes the headline interpretation of the climate ensemble. Over 2026–2045 the realised warming is about 0.6 °C on average (0.21–1.09 °C across runs), and the aggregate responses to that warming are weak and mixed: a small increase in total biomass, a small decrease in native richness and a small decrease in realised richness loss, each with low explanatory power. Most of the raw change over the horizon is a transient relaxation from the observed 2006–2009 community under the interim projection route, shared by all runs including the control (total biomass −35.5%, native biomass −40.8% in the control alone). Only the scenario-minus-control differences isolate the thermal effect, and those differences are small. This is consistent with the geographic constraint that every corrected baseline site lies at or below about 18 °C, so roughly 0.6 °C of warming does not cross the optima of the warm-adapted dominant natives.

The warming uncertainty band (E21) does not change this conclusion. Scaling the central warming by ×0.63–×1.28 moves the ensemble-median pathway warming within roughly 0.37–0.93 °C; even the high end remains mild, and the band is a conservative envelope that probably overstates warming for this basin because the lagged Guadalquivir slopes are lower than the same-day slope.

5.2 The trout signal is real but mostly transient

Brown trout is the clearest species-level thermal–hydraulic signal, but the corrected numbers require a careful reading. The species loses about 37% of its spun-up biomass even in the no-warming control, because the observed initial community is far from the interim-route equilibrium; only about one to one-and-a-half percentage points of the loss are attributable to warming, and the fitted sensitivity is about 1.8 percentage points per °C ($R^2 = 0.63$). The convergent evidence — a consistent negative scenario-minus-control difference, a monotone dependence on realised warming and increased exposure above 20 °C — still supports using trout as an early-warning indicator, but the absolute decline is not a warming forecast. The magnitude remains conditional on the symmetric Gaussian niche, the midpoint optimum and unvaried thermal breadth (C6). No basin-scale extinction forecast should be inferred.

5.3 Connectivity is context dependent

The corrected sweep shows a monotone, sign-stable effect of passability under this model: blocking raised and improving lowered basin-mean native biomass, at every upstream cost and both endpoints, while background upstream cost remained the strongest hydraulic control. This is the opposite of the September pattern, in which blocking reduced native biomass at low cost and increased it at high cost; that reversal was a property of the pre-correction interaction direction and network graph and is explicitly retired here. The absence of a reversal is a finding about this model's corrected structure, not a general ecological law: the fixed on-path tree, the multiplicative obstacle rule and the uniform passability assumption all shape it.

Because basin means conceal spatial redistribution, barrier decisions should still be evaluated at reach and sub-catchment scales against background hydraulic resistance, species composition

and invasive pressure. The present design cannot discriminate among reduced native immigration, reduced emigration and suppressed invasive dispersal.

5.4 Interaction structure is a modest structural uncertainty

In the corrected sweep, changing the interaction matrix moved native, invasive and trout biomass by only a few per cent, far less than the upstream-cost effect and comparable to the passability effect. This is a substantial downward revision from the September bounding contrast. It is consistent with the corrected competitive Lotka–Volterra form, in which the interaction term enters the growth bracket and is scaled by the site carrying capacity, so that realistic interaction coefficients produce bounded rather than catastrophic shifts. The random matrix remains uninterpretable as an ecological scenario because some species have near-zero burn-in baselines, and the thermal-breadth multiplier is confounded with the interaction axis. Future inference should use empirically estimated interaction coefficients, uncertainty distributions and independent replication across interaction and thermal-parameter sets.

5.5 What the structural sensitivities do and do not change

The Phase 2 sensitivities draw a clear line around the weak-warming conclusion. The parser repair is immaterial to every metric at the reported precision; the passability floor and the on-path-only obstacle assignment leave the factor ranking, the top-two ordering and the direction of every factor effect unchanged; and the competition alternative leaves the *magnitude* of the basin richness response small under both structures. What the competition structure does change is the interpretation of the average response: with species-specific competition the ensemble-mean native-richness delta is uniformly negative rather than mixed, and the small realised richness-loss signal of the shared-capacity form disappears. The headline should therefore be read as "the warming delivered here produces a small basin-scale response", not as "warming does not reduce native richness". Equally, the amplification of the total-biomass and trout responses under the alternative means that their magnitudes, not their signs, carry a structural uncertainty of roughly five- to eight-fold. Because the passability floor is a modelling choice and the parent-only variant is a topological counterfactual rather than a removal experiment, the magnitudes of native richness, invasive biomass and trout quasi-extinction should be read with the root-join overlay in mind, even though the rankings are stable. The two remaining climate caveats — the shared rather than per-GCM control (P2.2) and the national rather than refitted air–water calibration (P2.5) — were deferred by explicit decision and bound the deltas in ways this report does not remove.

5.6 Monitoring and management implications

Monitoring based only on aggregate richness or biomass is unlikely to detect early thermal deterioration, especially over a twenty-year horizon dominated by transient dynamics. Cold-water specialists should be monitored using species-level abundance, occupancy, upper-limit exceedance and thermal selection gradients, and should be assessed against the shared no-warming reference because the common transient is large. Brown trout remains a useful sentinel because warming and restricted movement act in the same direction for a sparse, headwater-associated population. Connectivity interventions should combine local passability, upstream movement cost, species movement traits and invasive distributions, and should not assume that increased passability uniformly improves aggregate native outcomes; in this corrected model it did not. Thermal refugia, especially high-elevation and headwater reaches, should be protected alongside connectivity improvements, consistent with adaptive river-management recommendations for uncertain futures [Tonkin et al., 2018]. Before quantitative management forecasts are attempted, the model should be calibrated (C3) and extended to represent discharge and drought dynam-

ics, abstraction, flow-dependent habitat change, land-use degradation, age structure, harvesting, acclimation and evolution.

6 Limitations and Scope of Inference

The following consolidated limitations define the scope of inference for the corrected release.

1. **Thermal niches are distributional, not physiological (C6).** Each species' thermal response is a symmetric Gaussian whose optimum is the midpoint of its reported thermal range and whose breadth is fixed at one-sixth of that range. This is a first-order sensitivity baseline, not a calibrated physiological niche. Because every modelled baseline site lies at or below about 18 °C and the delivered warming is mild, the symmetric curves have little leverage on the present results; that insensitivity is a consequence of the mild forcing and cool baseline, not a general property. Under stronger warming, replacement of the midpoint by literature optimum-for-growth and CTmax/UILT values, on an asymmetric (e.g. Sharpe–Schoolfield) curve, is required.

2. **The projections are interim and uncalibrated (C3).** They are initialised from the observed 2006–2009 community and advanced through a three-year baseline spin-up, not a converged model burn-in. Intrinsic growth rates, carrying capacities, interaction coefficients and dispersal parameters are not calibrated, and the model does not reproduce the observed community to a pre-registered standard. Every scenario is reported as a scenario-minus-control difference against one *shared* no-warming control (not a per-GCM control), so the common residual transient cancels but a GCM-specific offset remains; the 11 matched per-GCM controls (P2.2) were deferred by the Phase 2 decision and the residual offset is stated, not removed. The results are conditional and relative, not absolute forecasts. Full ABC-SMC calibration with the pre-registered TSS and occupancy criteria of Section 4.7, spatial block cross-validation and posterior predictive checks remains future work. A direct consequence is the large control transient documented in Section 4.1.

3. **The air–water warming response is a same-day calibration from the national model, with a lagged uncertainty band (E20/E21).** The projections are driven by the national all-Spain Stage-B air–water model ($t_a = 0.5097$, $n = 39{,}944$, 980 sites), not by a Guadalquivir-only fit ($t_a = 0.436$), so the 257 coded Waterbase values are in the calibration that reaches the GuadeX forcing. The coded-value refit and forcing regeneration (P2.5) were deferred by the Phase 2 decision; E20 remains a stated limitation. Lagged air fits yield higher slopes but rest on only about three in-sample lag-eligible sites ($n = 104$, one year); they cannot serve as a primary calibration and are reported as a band of $\times 0.63$–$\times 1.28$ on the central warming. Within the Guadalquivir subset the lagged slopes are lower than the same-day slope, so the upper end may not apply to the basin; the band is a conservative sensitivity envelope.

4. **Barrier passability rests on a uniform structural assumption (IF).** Passability is modelled as 0.1 upstream and 0.5 downstream, pending standardisation of the obstacle inventory's qualitative passability index (IF). The IF index is not decoded; using it prematurely would mis-scale every barrier. This is a first-order control on connectivity and should be revisited once the inventory semantics are standardised.

5. **The richness-loss metric is a deterministic, realised contraction, not an extinction probability (E8).** The model contains no demographic or environmental stochasticity and is run without replicates, so the metric describes a single deterministic trajectory and cannot be interpreted as, or converted into, a quasi-extinction or extinction probability.

6. **No hydrological dynamics, drought or abstraction.** The model does not represent discharge, flow intermittency, drought or abstraction; habitat suitability enters through a static index and the temperature series. Warming is the only climate-linked process that varies over the projection, so the results are not integrated climate-and-water-management projections.

7. **No age, stage or body-size structure and no evolution or acclimation.** Individuals are a single aggregated abundance per species and site; thermal traits are fixed. The projections may therefore overstate thermal stress relative to a model allowing adaptation.

8. **The river network is an on-path reconstruction, not a flow-directed channel layer.** The corrected graph is built by an on-path parent rule with an explicit minimum-spanning-tree root join, because the regional channel layer carries no flow direction or elevation. It is acyclic, deterministic and validated by topology, but its root-join edges are not flow-directed, and connectivity results should be read relative to this reconstructed topology. The root joins are 37.7% of total length (median 15.8 km, maximum 206.6 km) and add 41 elevation inversions to the 4 on parent edges, so a substantial part of the modelled corridor is a straight-line reconstruction rather than a surveyed channel. Obstacles also accumulate without limit on a single reconstructed link (33 obstacles, upstream factor 10^{-33}). The Phase 2 floor (10^{-3}) and parent-only variants are reported in Section 4.7: the factor ranking is identical under the primary, floor and parent-only variants (Spearman $\rho = 1.00$ for all eight metrics), the extreme link is a parent edge unaffected by parent-only and raised to 10^{-3} by the floor, and parent-only removes 245 applied obstacles from 111 root-join links (197 become unmatched, 48 re-snap). No passability floor is applied in the primary run, so connectivity magnitudes should be interpreted relative to this overlay.

9. **Biomass is reported in model units without an absolute scale.** Values are internally consistent and suitable for relative comparison but carry no absolute physical scale.

10. **Short projection horizon (2026–2045).** The twenty-year horizon is short relative to generation times and to community reorganisation, and truncates the analysis before the higher-emission pathways differentiate fully.

11. **Climate-model spread overlaps pathway differences, and pre-2026 offsets differ.** Pathway differences are small relative to the spread among GCMs within a pathway, so no pathway ordering is defensible where the spread overlaps it. Absolute temperature changes relative to 1986–2005 are not directly comparable across scenario products because of different pre-2026 offsets; this is why the 2026 scenario exposure already exceeds the baseline-climatology control. In addition, the alternative-interaction sweep confounds interaction matrix with a thermal-breadth multiplier of 0.3, the random matrix is a placebo with uninterpretable ratios, and *Alburnus alburnus* has been reclassified from migratory to exotic (EX) but still has incomplete interaction and dispersal trait data and a placeholder dispersal scalar, so its absolute density remains provisional.

12. **The 2 km obstacle crosswalk radius was fixed, not varied.** The radius is a conservative spatial join justified by the positional uncertainty of the inventory and the coarse network, and the distance curve shows a clear knee at 2 km. It was not subjected to a formal radius sensitivity analysis, however, and about 250 structures lie between 2 and 5 km on the corrected tree; a larger radius would add them to the overlay, potentially at edges several kilometres from the structure. The matched-obstacle count and the passability overlay should therefore be read as conditional on this crosswalk.

13. **Alien-species introductions are not represented.** The model contains no explicit introduction events. Which alien species to add, and where and when they arrive, remain

team decisions. Invasive richness and biomass are therefore counted from the current species pool, so the reported invasive metrics are conditional on that pool and do not project colonisation by new arrivals.

14. **The interaction-parser repair is quantified and immaterial to the conclusions.** The corrected table changes 16 reverse-direction entries from -0.5 predation to -0.3 competition and flags nine mixed "competition and predation" cells (34 changed directed rows; 227 rows; 31 ambiguous source cells / 62 directed rows). Re-running the three experiments against the fixed table moves the post-fix basin metrics by at most 0.0026 native species of basin mean richness (2 of 775 sites), 0.0013 of realised richness loss and 0.0011 biomass units on a ≈ 30-unit mean in the climate ensemble; at most 0.0013 species, 0.0063 biomass units and 0.057 trout biomass in the obstacle sweep; and at most 0.0013 species, 0.033 biomass units and 0.51 trout biomass in the interaction sweep. No aggregate sign changes or ordering changes occur. The repair is required for the correctness of the α matrix and the implied $c = 1 - \alpha$ coefficients, but it does not drive any conclusion.

15. **The competition structure changes the interpretation of the climate response (C2).** The primary specification remains the shared-capacity $c = 1-\alpha$; the species-specific $c = -\alpha$ alternative was run over all 44 scenarios as a sensitivity. The magnitude of the weak-warming result is robust (basin native richness changes by at most 0.0129 species, ≤ 10 of 775 sites, and the warm-endpoint delta is numerically identical at -0.006452), but the sign of the ensemble-mean richness response depends on the structure: shared-K gives a mixed, slightly positive mean ($+0.0020$; 26 positive, 4 zero, 14 negative), whereas the alternative is uniformly negative (44/44; mean -0.0071) and 30 of 44 runs flip sign. The small shared-K richness-loss signal collapses to approximately zero under the alternative (mean -0.001551 to -0.000037; 31 of 44 deltas exactly zero), while the total-biomass and trout responses amplify (about 4.8× and 7.8–8.3×). The claim that warming does not reduce native richness is therefore not robust to the competition structure; the weak-warming headline and every primary number are unchanged.

16. **The start-state validation is measured and marginal.** Against the post-spin-up state of the Phase 2 shared control, pooled sensitivity is 0.975 and pooled specificity 0.920 over 24 species and 775 sites; the widespread native LS has TSS 0.417 with observed occupancy 45.3% against modelled 77.2% (a +31.9-point over-prediction). The model systematically over-predicts occupancy, and several invasive species have weak control trajectories. The fixed acceptance thresholds for a future calibration campaign — per-species TSS ≥ 0.40 where at least 10 sites are occupied, and $|\text{modelled} - \text{observed occupancy}| \leq 0.10$ for LS, SA, SP, PW and CP — are calibration targets, not gates for this release; the current control passes the TSS criterion where it applies but fails the occupancy criterion for all five widespread natives.

Taken together, these limitations define a narrow and explicit scope. The present results are a conditional, internally consistent, deterministic sensitivity assessment of the effect of a mild, prescribed thermal forcing on an uncalibrated metacommunity model, reported against a shared no-warming control. They are informative about the sign, relative magnitude and spatial pattern of the modelled thermal response within the modelled assumptions; they are not absolute forecasts, not extinction probabilities, and not integrated climate-and-management assessments.

7 Conclusions

For the Guadalquivir metacommunity and the 2026–2045 horizon, the corrected analysis shows that aggregate native biomass and richness respond only weakly to the roughly 0.6 °C of realised warming delivered by the climate ensemble, and that the raw absolute changes over the horizon

are dominated by a transient shared with the no-warming control. This weak aggregate response is not evidence of thermal safety: it reflects cancellation between warm-adapted dominant natives and a cold-water specialist moving further above its optimum.

Brown trout provides the clearest species-level early-warning signal, but the corrected attribution is modest: about 1.1–1.5 percentage points of its roughly 38% interim-route decline are warming-attributable, with a sensitivity of about 1.8 percentage points per °C and increased exposure above 20 °C.

Upstream movement cost is the principal control of aggregate native biomass and richness, and passability is the principal control of invasive and trout biomass; the climate endpoint is a minor factor in the obstacle sweep. Passability acted monotonically in the corrected sweep — blocking raised and improving lowered basin-mean native biomass at every background cost — and the sign reversal of the September release is not recovered. The invasive-favouring interaction matrix produced only a small native-to-invasive redistribution, far weaker than the September bounding contrast, and identifies a bounded structural uncertainty rather than a forecast.

The Phase 2 sensitivities add two qualifications without overturning these findings. The interaction-parser repair changes at most 0.0026 native species of basin richness, and the 10^{-3} passability floor and the on-path-only obstacle assignment leave the factor ranking unchanged (Spearman $\rho = 1.00$), so those conclusions are structurally robust. The competition structure, in contrast, changes the sign of the ensemble-mean native-richness response (shared-K +0.0020 against species-specific −0.0071 species per site) and collapses the small realised richness-loss signal, while roughly five- to eight-fold amplifying the biomass and trout magnitudes; the weak-warming *magnitude* is preserved, but the stronger claim that warming does not reduce native richness is not robust and is withdrawn.

A defensible management strategy combines species-level thermal monitoring against the shared no-warming reference, protection of cold-water refugia, spatially targeted connectivity assessment and invasive-species control, and treats all quantitative magnitudes as conditional pending full calibration.

8 Acknowledgements

Fish sampling and handling were conducted under the relevant Spanish regulatory framework. Funding, permits, institutional support and author-specific contributions should be completed before formal publication.

9 Data and Code Availability

The study is based on empirical fish-density and juvenile observations, site connectivity and environmental measurements, species trait data, qualitative interaction information, a 2026 obstacle inventory and daily downscaled water-temperature projections for the Guadalquivir basin. The primary post-fix outputs analysed here are `results/climate_scenarios_phase2/`, `results/sensitivity_obstacles_phase2/` and `results/alt_interactions_phase2/`, driven by `parameters_climate_scenarios_corrected.toml`; the pre-fix `results/climate_scenarios_corrected/`, `results/sensitivity_obstacles_corrected/` and `results/alt_interactions_corrected/` directories are retained as the audit baseline, and the structural sensitivities are in `results/climate_scenarios_c` `results/sensitivity_obstacles_floor1e3/` and `results/sensitivity_obstacles_parentonly/`. Each run provides viewer-ready exports: four-level tables (`level_basin`, `level_subcatchment`, `level_water_body`, `level_sampling_point`), site-keyed aggregate viewer files, and per-species site-keyed viewer time series named `species_<sp>_<metric>_timeseries.json`, loadable in the published viewer at https://melian009.github.io/GuadeX/. The viewer currently hosts demonstration files; the post-fix outputs analysed here, together with the daily forcing files needed

to re-run them, were shared directly with the reviewers at https://k00.fr/eqnsvc2y. The final release should provide a persistent accession for model code, input data, scenario forcing, derived outputs and figure source data, together with the applicable licence and any restrictions on sensitive spatial data.

Legacy artifact. The September 2026 report (`legacy/FinalReportSeptember2026.tex`) and the pre-correction result directories (`results/climate_scenarios/`, `results/sensitivity_obstacles/`) are retained for comparison only. Their quantitative claims — including the warming range (0.74–0.97 °C by pathway), the total-biomass slope (−0.120 per °C), the 9–11% trout decline, the near-saturated trout quasi-extinction fraction and the ∼90% invasive-favouring contrast — are superseded by the corrected values in this report and must not be cited.

10 Figures




Figure 1: **Thermal niches against baseline water temperature.** *Grey bars:* relative frequency of the 2026 baseline water temperature across all 775 sampling sites (1 °C bins, normalised to the tallest bar). *Coloured curves:* the Gaussian thermal-suitability function $W = \exp[-(T - T_s^*)^2/(2\sigma_s^2)]$ for the native species; species with identical empirical ranges are drawn once. In the legend, which is placed outside the plotting area, each colour gives a species code and its optimum T_s^*, using the same species grouping convention as Fig. 2. Because σ_s is one-sixth of the empirical range, suitability falls to $\exp(-9/2) \approx 0.011$ at the range edges. *Vertical lines:* dotted black is the corrected basin-mean baseline temperature (15.82 °C); dotted crimson is the ensemble-median warming to 2045 on the corrected realised-forcing axis. *Reading:* the dominant cyprinids peak at 19–20 °C, warm of the present mean, so modest warming moves them towards their optima, whereas brown trout (*Salmo trutta*, ST, optimum 12.0 °C) has a mean baseline of 13.72 °C at its 72 baseline-established sites, about 0.65 thermal standard deviations above its optimum.




Figure 2: **Thermal niches against the corrected baseline water temperature (diagnostic rendering).** *Panel and axes:* a single panel; the x-axis is water temperature (°C) and the y-axis is the shared scale for thermal suitability and relative frequency (0.0–1.0). *Grey bars:* the relative frequency of the baseline (first simulated year) water temperature across the sampling sites, in 1 °C bins and normalised to the tallest bar. *Coloured curves:* the Gaussian thermal-suitability function $W = \exp[-(T - T_s^*)^2/(2\sigma_s^2)]$ for each native richness species; species with identical optimum and width are drawn once and grouped. Because σ_s is one-sixth of the empirical range, suitability falls to $\exp(-9/2) \approx 0.011$ at the range edges. The legend sits outside the plotting area (titled "Native species (optimum)") and gives each of the five resulting groups by species code and optimum: ST 12.0 °C; SP 16.5 °C; six species at 19.0 °C (AB, AH, PW, LS, SA, IL); IO 19.5 °C; and CP 20.0 °C. *Vertical reference lines:* dotted black is the diagnostic baseline mean (15.8 °C) and dotted crimson is the mean end-of-horizon temperature after the ensemble-median end-of-2045 warming (+0.74 °C → 16.6 °C; the 2026–2045 mean warming over the same runs is +0.58 °C). *Relationship to Fig. 1:* this figure shows the same corrected Gaussian niches and species table as Fig. 1, computed from the corrected baseline site temperatures, but rendered by the climate-diagnostics pipeline; it is included to document that the two pipelines agree and to record the corrected optima and baseline. *Reading:* most native optima (16.5–20.0 °C) lie above the baseline mean, so modest warming moves the assemblage towards, not away from, its thermal optima, whereas brown trout (ST, optimum 12.0 °C) already sits above its optimum and so is pushed further away by warming.




Figure 3: **End-of-2045 climate-ensemble summary by metric and spatial reporting level.** *Panels:* (a) native richness (species per site); (b) invasive richness (species per site); (c) realised native richness loss (fraction of the spun-up baseline); (d) total biomass (model density units per site); (e) projected water temperature (°C). The four spatial levels are the colour inside each panel: blue = sampling point, green = sub-basin, orange = water body, red = whole basin. *x-axis:* emission pathway, including the no-warming control. *Points:* mean across the 11 GCMs; bars: thin = 10–90% and thick = 25–75% spread across GCMs. In the corrected run the invasive-richness panel is nearly flat at about 0.76 species per site (range 0.755–0.765 across the 44 scenario runs) and the native richness and loss panels are also nearly flat; temperature separates the pathways while the aggregate biomass stays broadly within the control spread. The flatness is a model result, not a plotting error: richness is a threshold count and the horizon delivers only mild warming.




Figure 4: **Annual whole-basin trajectories across emission pathways, 2026–2045.** *Panels:* (a) native richness (species per site); (b) invasive richness (species per site); (c) realised native richness loss (fraction); (d) total biomass (model density units per site); (e) projected water temperature (°C). *Lines:* mean across the 11 GCMs of the whole-basin value per pathway; bands: 10–90% across GCMs. Colour: blue = SSP1-2.6, green = SSP2-4.5, orange = SSP3-7.0, red = SSP5-8.5, black = no-warming control. Temperature trajectories diverge as expected, but native richness, realised richness loss and total biomass remain nearly coincident across pathways, and the control shows a large transient decline in biomass that dominates the raw trajectories (control total biomass −35.5% from 2026 to 2045). The temporal persistence of the pattern supports interpreting the flat aggregate climate slopes as a model result; the scenario-minus-control differences isolate the small warming signal.




Figure 5: **Forcing versus response in the corrected climate ensemble.** "Forcing" is the realised basin-mean water-temperature anomaly ΔT relative to the 1986–2005 baseline. *Panels (a)–(c), annual trajectories 2026–2045:* (a) forcing ΔT (°C); (b) native richness (species per site); (c) total biomass (model units per site). Each line is the median across the 11 GCMs for one SSP pathway and the shaded band is the 10–90% range across GCMs; colours are blue = SSP1-2.6, green = SSP2-4.5, orange = SSP3-7.0 and red = SSP5-8.5. The native-richness and biomass panels overlap so closely that the in-panel annotation "all SSPs overlap" flags that the response trajectories lie on top of one another even though the forcing separates. *Panels (d)–(f), end-of-2045 responses, one point per GCM × SSP run (44 scenario runs):* (d) end-of-2045 native richness; (e) end-of-2045 total biomass; (f) end-of-2045 realised richness loss; each is plotted against the realised warming of that run, with points coloured by SSP and the pooled ordinary-least-squares slope, GCM standard deviation and R^2 annotated in the panel. The realised anomaly is used because it is the quantity that drives the biology in each run (correction C7). Correlations are computed across the 44 scenario runs; the no-warming control is plotted for reference but excluded from the fits, matching the dose–response analysis. The loss metric is a deterministic thresholded realised loss, not an extinction probability. *Reading:* warming separates the pathways in panel (a), but the aggregate richness, biomass and loss responses remain nearly coincident across pathways over this horizon.




Figure 6: **Basin response versus realised warming across the 44-run climate ensemble (2026–2045).** *Panels:* (a) end-of-2045 basin total biomass; (b) end-of-2045 basin native richness; (c) realised native richness loss. Each point is one GCM × SSP run (44 in total), coloured by SSP. *x-axis:* realised basin-mean warming over 2026–2045 (relative to 1986–2005), spanning 0.21–1.09 °C (the same regressor as Table 3). *Dashed line:* ordinary least-squares fit across the 44 points; the fitted slopes are +0.075 model units per °C ($R^2 = 0.14$), −0.0053 species per site per °C ($R^2 = 0.06$) and −0.00067 loss units per °C ($R^2 = 0.01$). These supersede the September slopes. The heavy overlap of pathway colours explains why no emission-pathway ranking is defensible over this horizon.




Figure 7: **Factor importance in the corrected obstacle sweep.** *Panel (a):* horizontal bars rank the seven factors of the sweep by their mean rank (1 = strongest; ties are broken by the plotting code's factor order); the x-axis is the mean normalised effect across the eight response metrics (1.0 = the largest effect for that metric), each bar is annotated with its value and its mean rank, and the bars are coloured consistently by factor. *Panel (b):* the per-metric normalised effect size as a heatmap; columns are the seven factors in the order passability, upstream cost, climate endpoint, interaction matrix, thermal breadth, passability×upstream cost and passability×temperature, and rows are the eight response metrics (bottom to top: basin total biomass, basin native biomass, basin invasive biomass, basin native richness, realised native richness loss, brown-trout relative biomass change, brown-trout final biomass and brown-trout quasi-extinct fraction). Every cell is annotated with its normalised value to two decimals, printed white on dark cells and black on light cells, and the colourbar maps the value 0–1 onto the viridis scale (dark purple = 0, yellow = 1). *Computation:* within each matched factorial block the effect span is the difference between the largest and smallest metric value across the levels of one factor; spans are standardised by the run-set standard deviation, the median span across blocks is taken, and for each metric the medians are divided by the largest (so the strongest factor in each row reaches 1.0). *Factor meanings:* passability = the four obstacle-overlay states; upstream cost = the four elevation-cost settings; climate endpoint = the cool versus warm sensitivity end; interaction matrix = the original, random-placebo and invasive-favouring matrices, taken from the alternative-interaction sweep rather than the obstacle sweep; thermal breadth = the thermal-sigma multiplier, which is confounded with the alternative-interaction sweep (`sigma_confounded`) and is therefore not independently identified; and the final two terms are the passability×upstream-cost and passability×temperature interaction columns. The mean ranks across the eight metrics (1 = strongest) are upstream cost 1.88 and passability 1.88 (tied), thermal breadth 2.50, interaction matrix 4.38, passability×upstream cost 4.75, climate endpoint 6.12 and passability×temperature 6.50. *Reading:* upstream cost leads for total and native biomass, richness and loss, whereas passability leads for invasive and trout biomass, so the aggregate community metrics are controlled mainly by network hydraulics rather than by the warming endpoint.




Figure 8: **Signed factor effects (tornado), one panel per response metric.** *Panels:* (a) total biomass, (b) native biomass, (c) invasive biomass, (d) native richness, (e) realised native richness loss, (f) brown-trout relative biomass change, (g) brown-trout final biomass and (h) brown-trout quasi-extinct fraction. *Axes:* the x-axis is the signed median effect in run-set standard-deviation (σ) units and a vertical black line marks zero, so a bar extending left means the level lowers the metric; the y-axis lists the factor levels being contrasted against their reference. *Each bar:* the median over matched factorial blocks of the signed difference between one factor level and its reference level, divided by the run-set standard deviation of that metric, so all bars are in comparable σ units. Bars are coloured by factor; the factor legend printed beside the panels defines the colours (passability, upstream cost, climate endpoint, interaction matrix and thermal breadth). Reference levels are upstream cost 0.01, baseline passability and the cool climate endpoint; the interaction-matrix and thermal-breadth contrasts are rescaled onto the main sweep's σ because the thermal-breadth multiplier is confounded with the sweep. *Reading:* increasing upstream cost reduces biomass and richness; blocked passability raises basin native and total biomass at every cost, while improved passability lowers it; and passability dominates trout biomass, consistent with the sum-of-squares results.




Figure 9: **Passability by upstream-cost interaction (obstacle sweep).** *Panels:* columns are the two climate endpoints (left: cool, SSP1-2.6/IITM-ESM; right: warm, SSP2-4.5/UKESM1-0-LL) and rows are four response metrics: total biomass, native biomass, native richness and brown-trout final biomass. *Axes and panels:* the four panels (a)–(d) are the four metric rows; x = the four upstream movement-cost settings (0.01, 0.05, 0.10, 0.50); y = the four passability states (baseline, improved, reduced, blocked). *Cell value:* the raw change relative to the same-model, same-cost baseline-passability run (so the baseline row is exactly 0), printed in each cell to three significant figures in black text with a white outline. *Colours:* the diverging "balance" colourmap is applied to the change scaled to the 95th percentile of the absolute values, with the colourbar labelled "Change vs baseline (scaled)"; the two ends of the scale mark the most negative and most positive changes. *Reading:* for native biomass the blocked-minus-baseline change is positive at every cost (+1.59 to +1.91 units), and improved passability is negative at every cost (−0.66 to −0.74 units); the September sign reversal is absent. The monotone pattern is a model result rather than evidence that barrier construction is a conservation action, because the design cannot separate reduced native immigration from suppressed emigration.




Figure 10: **Brown trout response to hydraulic structure.** *Panels:* (a) and (c) are the cool endpoint; (b) and (d) the warm endpoint. The top row is final *Salmo trutta* biomass and the bottom row its relative biomass change. *x-axis:* the four categorical upstream movement costs. *Coloured lines/points:* the four passability states; black = baseline, blue = improved, orange = reduced, red = blocked. *Reading:* all passability curves lie below the spun-up state, the warm endpoint is uniformly more negative, and at cost 0.05 the baseline-to-blocked change is 355.4 to 427.3 units at the cool endpoint (a hydraulic gain of +71.9 units), so warming and passability act on trout through different magnitudes.




Figure 11: **Passability robustness across interaction matrices.** *Panels:* (a) and (c) are the cool climate endpoint (SSP1-2.6/IITM-ESM) and (b) and (d) the warm endpoint (SSP2-4.5/UKESM1-0-LL); the top row (a, b) is brown-trout (*Salmo trutta*) final biomass and the bottom row (c, d) basin native biomass. *Axes:* the y-axis lists the three interaction matrices (original, random placebo and invasive-favouring) and the x-axis is the span across passability states—the difference between the largest and smallest metric value when the four passability states (baseline, improved, reduced and blocked) are compared at the same matrix, climate endpoint and upstream cost—in model units. *Bars and legend:* each matrix row carries two horizontal bars, one per upstream-cost setting, and the two-colour legend printed below the panels identifies them: blue (RGBA 0.27, 0.45, 0.77) = low upstream cost (0.01) and orange (RGBA 0.90, 0.55, 0.10) = high upstream cost (0.50); these are the two colours defined in the plotting code for the cost levels 0.01 and 0.50. *Reading:* brown-trout spans are about 29 units and native-biomass spans about 1.3 units under every matrix and both cost settings, so the passability signal is essentially matrix-independent in the corrected sweep. This supersedes the September claim of matrix-dependent spans.




Figure 12: **Spatial distribution of blockage effects.** *Scenario:* warm endpoint, high upstream cost (0.50), end of 2045; blocked minus baseline passability. *Panel (a):* the 10 most negative and 10 most positive of the 77 sub-catchments. *Panel (b):* the 10 most negative and 10 most positive of the 289 water bodies. Both panels use the same x-axis (change in native biomass, model units) and show bars sorted from most negative to most positive; red bars are losses and green bars gains. The positive basin-level blocked-minus-baseline effect is a net of local gains and losses, so it must not be read as a uniform benefit of blocking; the sub-catchment or water body is the appropriate scale for management decisions.




Figure 13: **Brown trout climate dose–response.** *Panels:* (a) relative change in basin *Salmo trutta* biomass (% of the spun-up baseline); (b) absolute end-of-2045 trout biomass; (c) maximum number of days per year above the 20 °C upper limit at the 72 baseline-established sites. *Points:* one per GCM × SSP run (44 in total), coloured by pathway; the black diamond at 0 °C is the no-warming control. *x-axis:* realised basin-mean warming at 2045. *Dashed line in (a):* OLS fit, giving a decline of about 1.76 percentage points of baseline per °C ($R^2 = 0.63$); the control changes by −37.3%, showing that most of the absolute decline is a common transient. Panel (c) shows the exposure mechanism: the baseline-climatology maximum is 90 days per year (dotted reference), rising to 80–135 days across scenarios in 2045. The figure links a species-level abundance response to a thermal-exposure mechanism while showing why it should not be extrapolated to the whole assemblage.




Figure 14: **Interaction-matrix main effects.** *Panels:* (a) basin native biomass; (b) basin invasive biomass; (c) brown trout final biomass; (d) basin native richness. *x-axis:* the three interaction assumptions (original, random placebo, invasive-favouring). *Points:* every one of the 16 runs per matrix (two climate endpoints × two upstream costs × four passability states), with slight horizontal jitter; the black horizontal line is the matrix mean. *Annotations:* the mean original → invasive-favouring change. Relative to the original matrix the invasive-favouring matrix changed native biomass by −1.7%, invasive biomass by +3.6%, trout biomass by −1.6% and native richness by −0.5%, while total biomass changed by −1.2%. These magnitudes are far smaller than the September bounding contrast and reflect the corrected interaction direction and the competitive Lotka–Volterra form; the sweep remains a bounding structural perturbation rather than a forecast.




Figure 15: **Obstacle-to-network crosswalk radius.** Cumulative number of the 1,658 non-completely-passable obstacles in the 2026 inventory whose nearest modelled network segment lies within a given straight-line distance, plotted on a logarithmic distance axis until every obstacle is included. Blue: original (legacy) network graph (973 obstacles within 2,000 m); red: corrected on-path tree used for the simulations (1,036 within 2,000 m). The dashed vertical line marks the 2,000 m radius used in the crosswalk and the symbols mark the two counts. The curve is steep below 2 km, capturing 59–62% of the inventory, and then flattens: 1,285–1,288 obstacles are within 5 km, 1,488–1,491 within 10 km, and the final obstacles are only reached at 227 km (on-path) or 295 km (legacy). The radius is a conservative spatial crosswalk, not a claim that unmatched structures are absent from the basin.

11 Summary Tables

Table 3: Post-fix climate-ensemble response at basin scale. Source: `results/climate_scenarios_phase2/runs_index.csv` and each run's `export/levels/` files.

Quantity	No-warming control	Scenario response (44 runs)
Realised basin-mean warming, 2026–2045	0.00 °C	0.54, 0.64, 0.62, 0.70 °C by pathway (SSP1-2.6, 2-4.5, 3-7.0, 5-8.5); 0.21–1.09 °C across runs
Realised basin-mean warming, 2036–2045	0.00 °C	0.60, 0.73, 0.71, 0.81 °C by pathway; 0.18–1.29 °C across runs
Total biomass vs realised warming	n/a	+0.075 model units per °C; $R^2 = 0.14$
Native richness vs realised warming	n/a	−0.0053 species per site per °C; $R^2 = 0.06$
Realised richness loss vs realised warming	n/a	−0.00067 per °C; $R^2 = 0.01$
Control change, 2026 → 2045	total biomass −35.5%, native biomass −40.8%, native richness +3.0%, invasive biomass +13.5%	n/a
Trout relative biomass change	−37.3%	−38.4 to −38.8% by pathway; scenario − control −1.1 to −1.5 pp
Trout sensitivity	n/a	−1.76 pp of baseline per °C; $R^2 = 0.63$
Invasive richness / biomass, 2045	0.763 sp/site / 5.05 units	0.761 sp/site / 4.94 units (range 0.755–0.765 / 4.84–5.04)

Table 4: Post-fix within-experiment variance contributions (sums-of-squares fractions). Source: `results/sensitivity_obstacles_phase2/report_plots/variance_decomposition.csv`.

Response (sweep)	Dominant factor	SS	Other notable contributions
Total biomass (obstacle)	Upstream cost	0.666	Passability 0.330; climate 0.0005
Native biomass (obstacle)	Upstream cost	0.729	Passability 0.269; climate 0.0005
Native richness (obstacle)	Upstream cost	0.935	Passability 0.041
Realised richness loss (obstacle)	Upstream cost	0.970	Passability 0.022
Invasive biomass (obstacle)	Passability	0.741	Upstream cost 0.194; passability×cost 0.063
Trout final biomass (obstacle)	Passability	0.951	Upstream cost 0.031; climate 0.015
Native biomass (interaction)	Upstream cost	0.668	Interaction matrix 0.116; climate 0.142
Trout final biomass (interaction)	Upstream cost	0.814	Passability 0.141; interaction matrix 0.034

12 Data Sources

This final report is a synthesis of the post-fix deterministic model outputs and empirical input collections. The principal output sources are `results/climate_scenarios_phase2/runs_index.csv` and the per-run `export/levels/` tables (`level_basin.csv`, `level_sampling_point.csv`, `species_timeseries` `exposure_summary.csv`, `quasi_extinction_summary.csv`); `results/sensitivity_obstacles_phase2/runs_` and `report_plots/` (`parameter_effect_rank.csv`, `variance_decomposition.csv`); `results/alt_interactio` and `report_plots/`; the supplementary sensitivity roots `results/climate_scenarios_competition_alt`, `results/sensitivity_obstacles_floor1e3` and `results/sensitivity_obstacles_parentonly`; `guadex_tw/outputs/tables/air_water_slope_uncertainty_band.csv`; and the obstacle-radius diagnostic `results/report_plots/fig_obstacle_matching_radius.png` with its source curve `obstacle_matching_radius.csv`. The pre-fix `results/climate_scenarios_corrected`, `results/sensitivit` and `results/alt_interactions_corrected` directories are retained only as the audit baseline for the parser-fix comparison. The viewer-ready exports for these runs (four-level tables and site-keyed aggregate and per-species time series, `species_<sp>_<metric>_timeseries.json`) are loadable in the published viewer at https://melian009.github.io/GuadeX/, which hosts demonstration files rather than the full corpus; the post-fix results and the daily forcing files are shared with the reviewers at https://k00.fr/eqnsvc2y. Persistent data/code accessions, licences and any restrictions on sensitive spatial data should be added in the submission version.

References

[Altermatt, 2013] Altermatt, F. (2013). Dendritic network structure and regional dynamics of metapopulations and metacommunities. *Critical Reviews in Environmental Science and Technology*, 43(16):1734–1775.

[Herrera et al., 2022] Herrera, M., Garcia-Berthou, E., Minano, P. S., Garcia-Meseguer, A. J., and Fernandez-Delgado, C. (2022). Long-term changes in the fish community of a mediterranean river basin: the role of dams and potamodromous species. *Water*, 14(15):2345.

[Larsen et al., 2021] Larsen, S., Comte, L., Filipe, A. F., Fortin, M.-J., Jacquet, C., Ryser, R., Tedesco, P. A., Brose, U., Eros, T., Giam, X., Irving, K., Ruhi, A., Sharma, S., and Olden, J. D. (2021). The geography of metapopulation synchrony in dendritic river networks. *Ecology Letters*, 24(4):791–801.

[Leibold et al., 2004] Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., and Tilman, D. (2004). The metacommunity concept: a framework for multi-scale community ecology. *Ecology Letters*, 7(7):601–613.

[Perkin and Gido, 2012] Perkin, J. S. and Gido, K. B. (2012). Fragmentation and defragmentation of dendritic ecological networks: a fish-community perspective. *Freshwater Science*, 31(1):36–44.

[Ramos-Merchante and Prenda, 2018] Ramos-Merchante, A. and Prenda, J. (2018). The ecological and conservation status of the guadalquivir river basin (s spain) through the application of a fish-based multimetric index. *Ecological Indicators*, 84:45–59.

[Tonkin et al., 2018] Tonkin, J. D., Bogan, M. T., Bonada, N., Rios-Touma, B., Sabo, J. L., and Lytle, D. A. (2018). Preparing river ecosystems for an uncertain future. *Nature*, 556(7702):427–433.