

# **Metacommunity capacity of complex life-cycles in disturbed landscapes**

**Authors**

**Addresses**

**Running title:**

**ABSTRACT**

**Keywords:** Metacommunities, patch dynamics, dispersal dynamics, individual-based model, approximate Bayesian computing,

## INTRODUCTION

Metapopulation models predict the survival of highly-disperse populations regardless local depletion of species ([Hanski, 1999](#); [Akçakaya et al. 2007](#)). However, habitat loss and fragmentation may be so extensive that result in a massive species extinction ([Montoya, 2008](#); [Rybicki & Hanksi, 2013](#); [Haddad et al. 2015](#)). Model predictions have shown that a decrease in connectivity among assemblages, from continuous to sparsely-distributed populations is accompanied by species loss ([Metzger et al. 2009](#); [Niebuhr et al. 2015](#)). The role of natural (e.g. stochastic events) and human-induced (e.g. pollution, harvesting) perturbations have been extensively studies in the last decades ([Dornelas, 2010](#) and references therein) and it has been frequently studied in ecological theory (e.g. [Volkov et al. 2007](#); [Gardner & Engelhardt, 2008](#)). The degree of anthropogenic pressure may be a capital factor for landscape connectivity, since directly affects the persistence or decrease of assemblages ([Supp & Ernest, 2014](#)). Besides perturbations, dispersal rates in disturbed landscapes need to be high in order to maintain viable populations ([Provan et al. 2009](#)). However, individual-based models (IBM) are needed to predict dispersal rates in species where small specimens, i.e. juveniles, are not reproductively active and even larger-sized adults harbor the highest reproductive potential (e.g. [Hendricks & Mulder, 2008](#); [Werner & Griebeler, 2011](#)).

IBM simulate populations as being composed of discrete individual organisms ([DeAngelis & Grimm, 2014](#); [Van der Väärt et al. 2016](#)). In IBMs the actions of single individuals are simulated and they interact with other and the landscape they live in ([DeAngelis & Mooij, 2005](#)). They incorporate attributes vary among the individuals and can change through time such as, growth, foraging, dispersal and reproduction, among others ([Martin et al. 2013](#); [DeAngelis & Grimm, 2014](#)). These models have been used as size-structured methods to integrate a high variety of data which output are pivotal for

management purposes ([Punt et al. 2013](#)) and conservation strategies ([Nabe-Nielsen et al. 2014](#)).

We herein develop metacommunity models based on individuals of two intertidal species. The first model assumes that dispersal rates between patches are distance-dependent, with low rates between highly-separated assemblages. The second model assumes that dispersal rates are positively correlated to individual density. The third model assumes that larger individuals have larger reproductive potential. The fourth model considers a low probability of dispersal to peripheral assemblages relative to central ones.

We confront the model with long-time series data (1994-2014) of two commercial limpet species (*Patella candei crenata* and *P. aspera*) in an overpopulated island (>500 inhab km<sup>-2</sup>) with a high coastal pressure ([Riera et al. 2016](#)). **TEXT ABOUT RESULTS AND CONCLUSIONS!!**

## REFERENCES

- Akçakaya, HR, G. Mills, C. Doncaster & C. Patrick. 2007. The role of metapopulations in conservation. In, McDonald, DW. & S. Katrina (eds.). Key topics in Conservation Biology. Oxford, UK, Blackwell publishing, 64-84.
- DeAngelis, DL. & V. Grimm. 2014. Individual-based models in ecology after four decades. *F1000Prime Rep*, 6: 39.
- DeAngelis, DL. & WM. Mooij. 2005. Individual-based modelling of ecological and evolutionary processes. *Annual Review of Ecology and Evolution Systematics*, 36: 147-168.
- Dornelas, M. 2010. Disturbance and change in biodiversity. *Philosophical Transactions of the Royal Society B*, 365: 3719-3727.
- Gardner, RH. & KAM. Engelhardt. 2008. Spatial processes that maintain biodiversity in plant communities. *Perspectives in Plant Ecology and Evolution Systematics*, 9. 211-218
- Haddad, N.M. et al.2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, 1: e1500052.
- Hanski, I. 1999. Habitat connectivity, habitat continuity, and metapopulations in dynamic landscapes. *Oikos*, 87: 209-219.
- Hendricks, AJ. & C. Mulder. 2008. Scaling of offspring number and mass to plant and animal size: model and meta-analysis. *Oecologia*, 155: 705-716.
- Martin BT., T. Jager, RM. Nisbet, TG. Preuss, M. Hammers-Wirtz. & V. Grimm. 2013. Extrapolating ecotoxicological effects from individuals to populations: a generic approach based on Dynamic Energy Budget Theory and individual-based modelling. *Ecotoxicology*, 22: 574-583.
- Montoya, D. 2008. Habitat loss, dispersal and the probability of extinction of tree species. *Community Integrative Biology*, 1: 146-147.
- Nabe-Nielsen J., RM. Sibly, J. Tougaard, J. Teilmann. & S. Sveegard. 2014. Effects of noise and by-catch on a Danish harbor porpoise population. *Ecological Modelling*, 272: 242-251.
- Niebuhr, BS, Wosniack, ME, Santos, MC. Raposo, EP, Viswanathan, GM. Da Luz, GE. & MR. Pie. 2015. Survival in patchy landscapes: the interplay between dispersal, habitat loss and fragmentation. *Scientific Reports*, 5: 11898.
- Provan, J. GE. Beatty, SL. Keating, CA. Maggs & G. Savidge. 2009. High dispersal potential has maintained long-term population stability in the North Atlantic copepod *Calanus finmarchicus*. *Proceedings of the Royal Society B*, 276: 301-307.

Punt, AE, T. Huang & M.N. Maunder. 2013. Review of integrated size-structured models for stock assessment of hard-to-age crustacean and mollusk species. *ICES Journal of Marine Science*, 70: 16-33.

Riera R, Ó. Pérez, O. Álvarez, D. Simón, D. Díaz, Ó. Monterroso & J. Núñez. 2016. Clear regression of harvested intertidal mollusks. A 20-year (1994-2014) comparative study. *Marine Environmental Research*, 113: 56-61.

Supp, Sr. & SKM. Ernest. 2014. Species level and community level responses to disturbance: a cross-community analysis. *Ecology*, 95: 1717-1723.

Van de Väärt, E. A. Johnston & RM. Sibly. 2016. Predicting how animals will de where: How to build, calibrate and evaluate individual-based models. *Ecological Modelling*, 326: 113-123.

Volkov, I. JR. Banavar, SP. Hubbell & A. Maritan. 2007. Patterns of relative species abundance in rainforests and coral reefs. *Nature*, 450: 45-49.

Werner, J. & EM, Griebeler. 2011. Reproductive biology and its impact on body size: comparative analysis of mammalian, avian and dinosaurian reproduction. *PLoS ONE*, 6: e28442.