

Training and nutrition factors
associated with body fat percentage in
males and females

Data and analytical approach for female

Analyzed the training aspect for female gender and following through this analysis steps :

1. Workout type by fat percentage
2. Workout type by workout frequency
3. Workout type by session duration
4. Workout type by calorie burned

Analyzed the nutrition aspect for female gender and following through this analysis steps :

1. Diet type by fat percentage
2. Protein per kg by fat percentage
3. Daily meals frequency by fat percentage

Data and analytical approach for male

Analyzed the training aspect for male gender and following through this analysis steps :

1. Workout type by fat percentage
2. Workout type by workout frequency
3. Workout type by session duration
4. Workout type by calorie burned

Analyzed the nutrition aspect for male gender and following through this analysis steps :

1. Diet type by fat percentage
2. Protein per kg by fat percentage
3. Daily meals frequency by fat percentage

Main question

What exercise and nutrition patterns are most associated with healthy body fat percentage ranges across demographic groups?

Female participants aged 40–55 who has 23% - 27% body fat percentage doing strength training 3–4 times per week....

**They exercised for
approximately 1.3 hours
per session**


They followed paleo diet

**They consumed
approximately 0.7
g/kg/day of protein**

**They consumed 2–4 meals
per day**

Training analysis for female

Workout type by fat percentage



Workout type by workout frequency



Workout type by session duration (in hours)




Workout type by calorie burned




Nutrition analysis for Female


Diet type by fat percentage



Protein per kg by fat percentage



Daily meals frequency by fat percentage



Male participants aged 40–55 who has 18% - 21% body fat percentage doing cardio training 3 times per week....

**They exercised for
approximately 1.2 hours
per session**


**They followed low-
carbohydrates diet**

**They consumed
approximately 0.7
g/kg/day of protein**


**They consumed 2–4 meals
per day**

Training analysis for male


Workout type by fat percentage




Workout type by workout frequency



Workout type by session duration in hours



Workout type by calorie burned




Nutrition analysis for male


Diet type by fat percentage



Protein per kg by fat percentage



Daily meals frequency by fat percentage



Limitation

This analysis is observational and identifies associations rather than causal relationships.

The dataset was filtered to participants with healthy body fat percentage ranges and may not generalize to all populations.

Lifestyle factors such as sleep quality, genetics, and overall caloric intake were not included.

Protein intake findings should not be interpreted as dietary recommendations.