

Density-dependent network structuring within and across wild animal systems

Gregory F Albery^{*1,2}, Daniel J Becker³, Josh A Firth^{4,5}, Matthew Silk^{6,7}, Amy R Sweeny^{8,9}, Eric Vander Wal¹⁰, Quinn Webber¹¹, Bryony Allen^{12,13}, Simon A Babayan¹⁴, Sahas Barve^{15,16}, Mike Begon¹², Richard J Birtles¹⁷, Theadora A Block^{18,19}, Barbara A Block²⁰, Janette E Bradley²¹, Sarah Budischak²², Cfhristina Buesching^{23,24}, Sarah J Burthe²⁵, Aaron B Carlisle²⁶, Jennifer E Caselle²⁷, Ciro Cattuto²⁸, Alexis S Chaine²⁹, Taylor Chapple³⁰, Barbara J Cheney³¹, Timothy Clutton-Brock³², Melissa Collier¹, David J Curnick³³, Richard J Delahay³⁴, Damien R Farine^{35,36,37,38}, Andy Fenton¹², Francesco Ferretti³⁹, Helen Fielding^{40,41}, Vivienne Foroughirad^{1,42}, Celine Frere⁴³, Michael G Gardner⁴⁴, Eli Geffen⁴⁵, Stephanie S Godfrey⁴⁶, Andrea L Graham⁴⁷, Phil S Hammond⁴⁸, Maik Henrich⁴⁹, Marco Heurich^{49,50,51}, Paul Hopwood⁷, Amiyaal Ilany⁵², Joseph A Jackson¹⁷, Nicola Jackson⁴³, David Jacoby⁵³, Ann-Marie Jacoby⁵⁴, Miloš Ježek⁵⁵, Lucinda Kirkpatrick⁵⁶, Alisa Klamm⁵⁷, James A Klarevas-Irby^{35,36,58,59}, Sarah Knowles⁴, Lee Koren⁵², Ewa Krzyszczyk⁶⁰, Jillian M Kusch^{61,62}, Xavier Lambin⁶³, Jeffrey E Lane⁶², Herwig Leirs⁵⁶, Stephan T Leu⁶⁴, Bruce E Lyon¹⁸, David W MacDonald²³, Anastasia E Madsen^{65,66}, Janet Mann¹, Marta Manser⁶⁷, Joachim Mariën⁵⁶, Apia Massawe⁶⁸, Robbie A McDonald⁴¹, Kevin Morelle^{55,69}, Johann Mourier^{70,71,72}, Chris Newman²³, Kenneth Nussear⁷³, Brendah Nyaguthii^{37,38,58,74}, Mina Ogino^{35,36,58}, Laura Ozella⁷⁵, Yannis Papastamatiou⁷⁶, Steve Paterson⁷⁷, Eric Payne⁷⁸, Amy B Pedersen⁹, Josephine M Pemberton⁹, Noa Pinter-Wollman⁷⁹, Serge Planes^{71,72}, Aura Raulo⁴, Rolando Rodríguez-Muñoz⁷, Christopher Sabuni⁶⁸, Pratha Sah¹, Robbie J Schallert⁸⁰, Ben C Sheldon⁴, Daizaburo Shizuka⁸¹, Andrew Sih⁷⁸, David L Sinn⁸², Vincent Sluydts⁵⁶, Orr Spiegel⁴⁵, Sandra Telfer⁶³, Courtney A Thomason^{39,83}, David M Tickler⁸⁴, Tom Tregenza⁷, Kimberley VanderWaal⁸⁵, Eric L Walters¹⁵, Klara M Wanelik^{77,86}, Elodie Wielgus⁴⁹, Jared Wilson-Aggarwal^{7,41}, Caroline Wohlfeil⁴⁴, Shweta Bansal¹

*Corresponding author: gfalbery@gmail.com. Affiliations are at the end of the document.

Abstract

High population density should drive individuals to more frequently share space and interact, producing better-connected spatial and social networks. Despite this widely-held assumption, it remains unconfirmed how local density generally drives individuals' positions within wild animal networks. We analysed 34 datasets of simultaneous spatial and social behaviour in >55,000 individual animals, spanning 28 species of fish, reptiles, birds, mammals, and insects. >80% of systems exhibited strongly positive relationships between local density and network centrality, providing broad empirical evidence that local density increases connectedness at the individual level. However, >75% of density-connectedness relationships were nonlinear, and density's importance declined at higher values in >70% of systems, signifying saturating effects. Density's effect was much stronger and less saturating for spatial than social networks, suggesting population density drives individuals to become disproportionately spatially connected rather than socially. These findings reveal fundamental trends underlying societal structuring, with widespread behavioural, ecological, and evolutionary implications.

Keywords: Behavioural ecology, Spatial ecology, Disease ecology, Epidemiology, Population dynamics, Social network structure, Network analysis, Spatial analysis

Introduction

The number of individuals occupying a given space – i.e., population density – is a central factor governing social systems. At higher densities, individuals are expected to more frequently share space, associate, and interact, producing more-connected spatial and social networks and thereby influencing downstream processes such as mating, learning, and competition. In particular, density-driven increases in network connectedness should provide more opportunities for parasites to spread between hosts [1–5]. Despite the fundamental nature of such density-dependent processes, evidence is relatively limited that individuals inhabiting higher-density areas have more spatial and social connections. Furthermore, density's effects should differ for asynchronous space sharing (e.g. home range overlap) *versus* social associations (e.g. den sharing or grouping) or interactions (e.g. mating or fighting). While several studies have compared animal populations at different densities to demonstrate variation in social association rates among populations (e.g., [6,7]) or groups (e.g., [8–10]), attempts to identify such density effects *within* continuous populations of individuals are rarer (but see [11–14]), and their findings have never been synthesised or compared. We therefore have an incomplete understanding of how density, as a fundamental ecological parameter, determines socio-spatial dynamics within and across systems. This inhibits our ability to identify and predict how changes in density – e.g. through culling, natural mortality, dispersal, or population booms – influence downstream processes that depend on shared space and social interactions.

The rate at which an individual interacts with conspecifics depends on its spatial and social behaviour within the context of the surrounding environment and population. Adding more individuals into the same space should cause them to more frequently spatially overlap and socially associate or interact (Figure 1). Often, individuals are modelled as randomly moving and interacting molecules (“mass action” or “mean field”). In this conceptualisation, direct contact between two molecules is analogous to a social interaction or association; rates of such interactions are often assumed to increase with density (“density-dependent”; e.g., [15]), and/or to be homogenous in space (e.g., [10]). In reality individuals are unlikely to behave and interact randomly in space, and instead will be influenced by spatially varying factors including local density [16]. Changes in density may cause individuals to alter their foraging behaviour [17–19], dispersal [20,21], social preference or avoidance [13,22], mating behaviour [23], or preferred group size [7]; in some cases, interaction rates may change to remain constant in the face of changing densities [24]. These and related processes might produce strong nonlinearities in density-interaction relationships, which can complicate the predictions of density dependence models of pathogen transmission, for example [2,4,5]. Nevertheless, these nonlinearities are poorly understood and rarely considered.

Several wild animal studies have suggested saturating nonlinear relationships between density and social association rates [8–10,13]. Such relationships imply that association rates do not increase passively with density, and that behavioural or demographic processes likely change as density increases, with the ultimate consequence of slowing association rates. However, these nonlinearities are difficult to examine between populations or between species because they introduce a range of confounders and have few replicates along the

density axis [2]. Characterising gradients of density across individuals within a population offers a workaround to this problem, and facilitates an appreciation of the fact that interactions occur between individuals rather than at the population level. Examining between-individual variation is one reason that social network analysis – which allows characterisation and analysis of individual-level social traits, amongst other things – has become so popular in animal ecology in recent years [25–29]. Additionally, recent years have seen a substantial growth in understanding of socio-spatial behaviours, including harmonising the concepts of spatial and social density [2,16,30]. Applying network analyses coupled with this socio-spatial understanding of density could provide an individual-level picture of density's effects on spatial and social connectedness, offering far higher resolution and statistical power and greater ability to detect nonlinearities and between-system differences [2]. By providing new understanding of the correlates and emergent consequences of variation in density, this expansion could help to identify general rules underlying social structuring and network scaling in space.

Critically, different types of interactions or associations should show different relationships with density: for example, the need to compete for food at higher densities could drive a disproportionate increase in aggression [31], but this is unlikely to be true of mating interactions. This rationale is well-understood in disease ecology, as differences in density-contact relationships are thought to drive differences in density dependence of infection – where “contact” is defined as an interaction or association that could spread a pathogen (Figure 1). “Contacts” then form the basis of spatial and social networks used to investigate pathogen transmission dynamics, which should likewise diverge with density just as contacts do. For example, density should drive greater transmission of respiratory pathogens but not sexually transmitted pathogens [1,32]. Establishing these density-contact relationships is integral to understanding disease dynamics and developing control measures [1,33], but we still have a poor understanding of how different interactions (and therefore contact events for different pathogens) are driven by density. This direct/indirect interaction dichotomy is most fundamental to disease ecology [30], but given building interest in the spatial-social interface and relationships between spatial and social networks in behavioural ecology [16], the framework is readily related to other fields. Established density-interaction relationships are diverse and include feral dog bites [11], ant antennations [34] and trophallaxis [24], ungulate group memberships [12,17], rodent co-trapping [8,35], and agamid association patterns [13,14], but no study has yet synthesised how the rates of multiple interaction types relate to density, within or across systems.

Identifying the general rules underlying density dependence requires quantifying density's relationship with proxies of interaction rates at fine scales across a diversity of systems, then identifying the factors determining their slope and shape. To this end, we collate a meta-dataset of over 55,000 individual animals across 34 wildlife systems globally to ask how within-population variation in density determines between-individual interaction rates based on connectedness in spatial and social networks. We fit multiple linear and nonlinear relationships to identify the slope and shape of density effects within each system, and we use meta-analyses to investigate general rules determining their slope and shape across systems. In particular, we focus on comparing space sharing with social interactions and associations as a cross-system case study. Ultimately, we present a *de novo* cross-system analysis of individuals' social and spatial behaviour that traverses fields of behavioural,

population, and disease ecology, which could help to inform general rules governing the structure of social systems.





Figure 1: Schematic detailing the rationale underlying this study, outlining how population density drives the formation of spatial and social networks. This depiction uses the Wytham Wood great tits as an example. Ultimately, we aim to ask whether spatial or social connections generally show a stronger relationship with density, partly functioning as a proxy for indirect and direct contact events with the potential to transmit pathogens. This framework moves between concepts of network and contact formation traversing behavioural ecology, spatial and social network ecology, and disease ecology.

142



143

144 *Figure 2: The phylogenetic (A) and geographic (B) distribution of our 34 examined datasets of spatial*
 145 *and social behaviour, with (C) schematic depicting the methodology for deriving local density values,*
 146 *using the Isle of Rum red deer data as an example. The X and Y axes are bivariate spatial*
 147 *coordinates. The panels within (C) show raw observations of individuals in space that we then*
 148 *average at the individual level to make centroids; we use the centroids to generate annual density*
 149 *distributions, which are then assigned to individuals in the form of local density measures. Animal*
 150 *silhouettes are from phylopic.org; a list of attributions is in the supplement. NB the Potomac dolphins*
 151 *are now defined as Tursiops erebennus; they are currently incorporated in Panel A as T. truncatus,*
 152 *following the Open Tree of Life nomenclature.*

153 Methods

154 Data standardisation and behavioural pipeline

155 Data were manipulated and analysed using R version 4.2.3 [36], and all R code is available
 156 at <https://github.com/gfalbery/DensityMetaAnalysis>. Our 34 datasets each involved at least
 157 one continuous uninterrupted spatial distribution of observations in a single population; some
 158 datasets comprised multiple such populations. These datasets covered 28 different host
 159 species, including sharks, carnivores, cetaceans, ungulates, rodents, elephants, birds,

reptiles, and one insect (Figure 2). In one case (The Firth of Tay and Moray Dolphins) we used two distinct replicates despite being composed of overlapping groups of individuals, because of their distinct spatial distributions, which made it difficult to fit a coherent density distribution.

To standardise the timescale across studies, all systems were analysed as annual replicates – i.e., social and spatial networks were summarised within each year. Our analyses used 60 system-behaviour replicates, listed in Supplementary Table 1, and totalled 151,507 unique system-individual-year-behaviour data points.

All spatial coordinates were converted to the scale of kilometres or metres to allow comparison across systems. To provide an approximation of local density, following prior methodology [12,37], we took each individual's average location across the year (their centroid) and created a spatial density kernel using the `adehabitat` package [38], which provides a probabilistic distribution of population density across each study system based on the local frequencies of observed individuals. Each individual was assigned an annual estimate of local density based on their centroid's location within this spatial density distribution. We made these density distributions as comparable as possible between systems by incorporating the density raster using metre squares; however, there were large differences in density across populations that were difficult to resolve and put on the same scale (e.g. interactions per individual/km² unit of density). Consequently, we scaled and centred density to have a mean of zero and a standard deviation of one within each population, which allowed us to focus on differences in relative slope and shape across systems.

To provide a measure of asynchronous space sharing, we constructed home range overlap (HRO) networks based on proportional overlap of two individuals' minimum convex polygon (MCP; i.e., the bounding polygon around all observations of each individual in a given year). These HRO networks were restricted to only individuals with five or more observations in a given year to allow us to create convex polygons effectively; 10/34 (29%) systems did not have sufficient sampling for this analysis. We also repeated our analyses with a series of higher sampling requirements for observation numbers to ensure that our findings were robust to this assumption. The MCP approach is relatively low-resolution, and assumes uniform space use across an individual's home range; however, this approach is less data intensive – and less sensitive to assumptions – than density kernel-based approaches that would estimate variation in space use across the home range, allowing us to apply the models across more systems, more generalisably, and more conservatively.

To provide a measure of social connectedness, we built social networks using various approaches as defined by the original studies: direct observations of dyadic interactions (e.g. fighting or mating); gambit of the group (GoG; i.e. membership of the same group) [39]; co-trapping (i.e. trapped together or in adjacent traps within a given number of trapping sessions); or direct contact measured by proximity sensors (defined by a certain distance-based detection threshold). Notably some analyses use indirect interactions – i.e., spatial overlap – to *approximate* direct interactions, which requires spatiotemporal coincidence, which we caution against particularly when modelling pathogen transmission [30,40]. While the two do often correlate, here we are not using HRO to approximate direct interaction

rates, but rather as a measure of indirect interactions (e.g., indicative of transmission of environmental parasites).

For each social network, we scaled connection strength relative to the number of observations of each individual in a dyad (i.e., simple ratio index [41]). Our response variable therefore took the form of strength centrality, scaled to between 0-1 for each dyad, for each social and spatial network. We focus on comparing density effects on social interactions and associations with density's effects on space sharing.

Density-connectedness models: what forms do density effects take?

We developed a novel workflow to allow us to derive and compare density's effects on connectedness – and their drivers – in a standardised way across our wild animal systems. These took three forms: linear models fitted to the whole dataset, nonlinear models fitted to the whole dataset, and linear models fitted separately to low- and high-density subsets of each dataset.

Linear models: For each system-behaviour replicate, we first fitted a linear model using the ``lm`` function in R, fitting scaled density as an explanatory variable to estimate linear density effect slopes.

Generalised additive models (GAMs): We fitted GAMs in the ``mgcv`` package [42] to identify whether each density effect was better described by a linear or nonlinear relationship, and to identify the shape of these nonlinear relationships. For each model, we fitted a default thin plate spline with $k=4$ knots. This knot number was selected to reduce overfitting in our models, which formed several fits to the data that were difficult to reconcile with functional formats. To assess whether nonlinear models fit better than linear models, we used Akaike Information Criterion (AIC), with a contrast of $2\Delta AIC$ designated to distinguish between models. See Supplementary Figure 1 for a schematic depicting our analytical workflow.

Saturation models: To quantify whether density effects were generally saturating (i.e., that density had steeper relationships with individuals' connectedness at lower density values), we split the data into two portions: all values below the median density value, and all values above the median. We then re-ran linear models examining the relationship between density and strength in each portion. We attempted to investigate nonlinear patterns (especially saturating effects) across all our systems using a range of other methods (e.g., comparing specific functional relationships with nonlinear least squares), but found that they were generally incapable of fitting well to the data in a standardised way across the many datasets (i.e., non-convergence of nonlinear least squares using semi-automated starting estimates across systems). As such, this approach represented a tradeoff between tractable, generalisable model fitting, interpretability, and accurate representation of the relationship's shape. All else being equal, we posit that investigating the relative slopes of two otherwise-identical portions of the data is a conservative and informative method of identifying saturation, which was our main hypothesis for the expected shape of density effects.

Meta-analysis: what factors determine the slope of density-connectedness relationships?

To characterise the typical relative slope of density effects across systems and identify the factors influencing their variation, we fitted hierarchical meta-analytical models using the ``metafor`` package in R. The response variable was the standardised slope of the linear density effect; because both individual network strength and density were scaled to have mean of zero and standard deviation of one in the linear regression, this is equivalent to the correlation coefficient (r) [43]. We converted all correlation coefficients into Fisher's Z (Z_i) and computed associated sampling variance. We then used an initial model that nested observations within a system-level random effect to account for within- and between-system heterogeneity [44], as 24/34 systems had more than one density effect. We used a random effect for species to account for repeat observations per animal species.

We then added a separate random effect for animal phylogeny [45]. This used a phylogenetic correlation matrix of our 28 animal species derived from the Open Tree of Life via the ``rotl`` package [46], with the ``ape`` package used to resolve multichotomies and provide branch lengths [47]. We fitted the model using the ``rma.mv()`` function with restricted maximum likelihood (REML), weighted by inverse sampling variance, and used variance components to quantify R^2 , the contribution of true heterogeneity to the total variance in effect size.

We next fitted models with the same random effects structure that included explanatory variables. To detect whether some animals were more likely to experience density effects, we fitted **Animal group** as a factor with six categories, representing a combination of species' taxonomy and general ecology: aquatic (fish and dolphins), birds, large herbivores (elephants and ungulates), small mammals (rodents and hyraxes), carnivores, and ectotherms (insects and reptiles). We also fitted several explanatory variables indicative of greater statistical power that might increase the strength of density effects: **Geographic area** (km^2 , \log_{10} -transformed), **Number of years** of study, and **Number of individuals**, all of which we fitted as continuous covariates. Broadly, the animal group model was highly uninformative and competed with the other effects, and we expected that the phylogeny would be more informative, so we report the results of the model without the host group effect fitted.

We ran several different versions of these meta-analyses: first, we fitted meta-analytical models to the **overall linear models** of spatial and social interaction types separately, and then together, to investigate differences between the spatial and social networks in terms of their mean density slope. Next, we fitted duplicated versions of these models, but with the **saturation models**. These models were identical, but each system replicate had two linear estimates: one taken from the first 50% of the data (up to the median), and one to the latter 50%. By fitting a binary fixed effect of "data portion" to the meta-analytical models, this model would tell us whether the slopes were generally higher in the first portion of the data than the last (and therefore showed generally saturating shapes). We were unable to fit meta-analytical models to our GAMMs, as methods capable of meta-analysing nonlinear estimates are not yet well defined.




Figure 3: Meta-analysis revealed drivers of variation in linear density effects on individual network connectedness across systems. A) Our fitted linear model estimates of density effects on network strength. Each point represents the mean estimate from a given system; the error bars denote 95% confidence intervals. Opaque error bars were significant (i.e., do not overlap with 0); transparent ones were not. The estimates are in units of standard deviations for both density and network strength. The colour of the point denotes whether the network being examined was defined using spatial or social connections. B) Meta-analyses revealed that centrality in spatial networks (i.e., home range overlap; red points) had a significantly steeper relationship with density than social networks (blue points). C) When linear models were fitted separately to two portions of the data (values below and above the median), the slopes for the latter portion (pink points) were generally less positive than the former portion (purple points), implying a general saturation shape. In panels B) and C), each coloured point represents a study replicate fitted to the strength estimate; points are sized according to sample size, and jittered slightly on the x axis to reduce overplotting. The large black points represent the mean slope estimated from the meta-analysis, and the error bars represent 95% confidence intervals.




Figure 4: Relationships between density and network connectedness varied substantially across animal systems. Density in individuals per area is on the x axis; network connectedness (strength centrality) is on the y axis. Both values have been standardised to have a mean of zero and a standard deviation of 1 within each system; the axis ticks are in units of 1 standard deviation. Each point represents an individual-year-behaviour replicate; the lines portray the model fit from our GAMs. Red lengths of the smooth=significantly positive; grey=not significantly different from zero; blue=significantly negative. Points are semi-transparent to enhance visibility. Panels are arranged phylogenetically following the tree displayed in Figure 2A; GOG=gambit of the group; HRO=home range overlap. Supplementary Figure 2 shows a version of this plot with variable smoothing parameters to show variation in the possible shape of the curves. Animal silhouettes are from phylopic.org; a set of links and attributions are in the Supplement.

Results and Discussion

We compiled an unprecedented comparative meta-dataset of over ten million observations of individual animals' spatial and social behaviour, across an expansive range of ecological systems. We then ran a standardised pipeline to align their spatial and social observations, identifying strong and predictable relationships between local density and network connectedness at the individual level.

We observed strong positive relationships between individuals' local population density and their connectedness in spatial and social networks across a wide range of wild animals: of our 60 replicates, 48 (80%) were significantly positive when analysed using linear models (Figure 3A; Supplementary Table 1). Meta-analyses identified a highly significant positive mean correlation between density and connectedness, both for social networks (Estimate 0.22; 95% CI 0.16, 0.27) and spatial networks (0.45; 0.36, 0.53; Figure 3B). Our study therefore provides fundamental evidence that high local population density broadly drives greater connectedness within ecological systems, at the individual level. Slopes were highly variable across systems for both spatial and social networks (Figure 3A; Q-test of heterogeneity across systems: $Q_{35} = 5744.83$ and $Q_{23} = 1559.08$, both $P < 0.0001$), indicating that quantifying these slopes within and between multiple systems and comparing them is important for understanding animal socio-spatial structure. That is, relationships between density and individual connectedness differ substantially between populations, and the biological mechanisms underlying these divergent trends are likely important. As well as adding resolution and allowing comparisons of density effects across systems, our methodology facilitated fitting of nonlinear relationships. This approach has only rarely been applied before, and then at much coarser resolution (see [8,9,11]). As such, this study fills an important empirical gap by providing insights into the slope and shape of density-connectedness relationships for a variety of animal groups and their social and spatial behaviours (Figure 4). Nevertheless, we were able to identify several further general trends in our data.

Remarkably, density's effect more than doubled in size for spatial compared to social networks (Figure 3B; $r=0.45$ *versus* 0.22); there was a difference of 0.28 (CI 0.17, 0.38, $P < 0.0001$) for this effect when we meta-analysed the two contact types together. This finding indicates that as density increases, wild animals are more likely to share space with each other, but that social connections increase at a much slower rate. Similarly, we discovered that saturating shapes were extremely common: as density increased, its effect on connectedness decreased, such that 44/60 systems (73%) had a steeper slope at low density values than at high ones. This effect was strong for social networks (effect on $r = -0.11$; CI -0.20, -0.03; $P=0.01$) and similar in size but marginally insignificant for spatial networks (-0.14; -0.29, 0.01; $P=0.058$); due to the greater overall effect for space sharing, the latter half of the density-spatial connections effect was still higher than the first half of the density-social connections effect (Figure 3C). In fact, the second half of the social effect overlapped with zero (Figure 3C; lower CI=-0.0119): that is, considering only the upper half of density values would not have uncovered a significant effect of density on individuals' connectedness in social networks. Together, these observations suggest that density-dependent processes act to limit the increase in social connectedness with density, but without limiting spatial overlaps to the same extent. Consequently, higher-density areas are characterised disproportionately by individuals asynchronously sharing space rather than

socially associating, while in lower-density areas individuals are disproportionately more socially connected proportional to their shared space.

There are many possible social reasons for saturating nonlinearity in density-dependent network structuring: for example, individuals in higher density areas may begin to avoid each other, seeking to avoid competition or aggression [31] or exposure to infectious disease [48]. Eastern water dragons (*Intellagama lesueurii*) show greater avoidance at higher densities [13], supporting this mechanism. Alternatively, in species with high social cognition or stable bonds, saturation could reflect lower social effort or ability to keep track of social affiliates at higher densities [49]. In general, individuals likely have a preferred social interaction rate or group size – a preference that they may increasingly exert at higher densities [7]. It remains to be seen how this preference varies among individuals, and whether individuals vary in their preferred social network position given a certain density. Given that individuals vary in their movement and spatial phenotypes [50–52], and social phenotypes [52–54] in ways that should manifest for density-dependent behaviours specifically [16], it seems likely that these slopes could vary between individuals as they do between populations. Future analyses might fit variable density-connectedness slopes between individuals to identify socio-spatial syndromes across systems, as has been done previously in single systems including caribou (*Rangifer tarandus*) [55] and red squirrels (*Tamiasciurus hudsonicus*) [56].

We considered spatial behaviours might explain these trends: for example, density could create greater competition over resources and therefore reduce energy to roam (and contact others). Individuals may partition their niches [57], or reduce their territory or home range sizes [56,58,59], potentially driven by years of plentiful resources supporting higher densities alongside smaller home ranges which could drive lower association rates. However, our findings do not seem to support explanations related to small home ranges, because such explanations should produce an equivalent or stronger reduction in (relative) spatial connectedness. In contrast, we observed that density drove individuals to become spatially connected faster than socially, such that the underlying mechanisms likely involve behaviours and demographic processes that specifically affect social collocation in space and time. Testing the precise underlying mechanism might require finer-scale behavioural observations, as described below. Regardless of mechanism, these saturating density-connectedness relationships strongly support the idea that examining density effects at the individual level – rather than between populations – is highly informative. For many systems, “mean field” expectations of homogenous interactions under increasing density likely produce an inaccurate (i.e., inflated) picture of density’s effects.

The fact that spatial networks show stronger and more linear density dependence than social networks is likely to have important implications for the ecology of wild animal systems. For example, indirectly transmitted (i.e., environmentally latent) parasites may exhibit greater density dependence than directly transmitted ones, given that individuals likely experience disproportionately more indirect contact at higher densities. This observation contrasts with orthodoxy that contagious parasites are most likely to be density dependent [60], and supports the value of investigating nonlinear changes in socio-spatial behaviour and grouping patterns in response to density when considering density dependence. Saturating density-connectedness functions further have implications for disease modelling and control. Rather than assuming constant behavioural mixing at higher densities, epidemiological models could benefit from incorporating density-dependent shifts

in behaviours and demography that influence direct and indirect interaction frequencies, as previously suggested empirically and by epidemiological theory [15]. These relationships could influence our targets for culling or vaccination coverage [61]. Given that animals at high density seem likely to have a relatively shallow relationship between density and contact rates, reducing population density – for example via culling – might not be effective at reducing pathogen transmission initially, particularly when considering socially transmitted pathogens, where contact rates are particularly likely to have saturated (Figure 3C). Similar problems with culling have already been acknowledged in specific systems – e.g. in canine rabies [33,62,63] – but our study implies that shallow nonlinear density-contact trends could be more general than previously thought and could be driven by flexible density-dependent changes in behaviour and demographic processes.

Operationally, the common nature of saturating density effects will impact researchers' ability to detect density dependence: that is, density dependence could be harder to observe in higher-density areas given the shallower slopes we observed. Most of the systems in this study are relatively long-term studies of known individuals; these populations tend to be in carefully selected, high-density areas that make it convenient to study the focal animal with relatively low operational costs. For example, it has previously been noted that the badgers of Wytham Wood, the red deer of the Isle of Rum, and the Soay sheep of St Kilda are all at high densities for their respective species [64–66]. As such, we may be inherently investigating the upper end of density-connectedness relationships in the wild, and it could be difficult and costly to investigate the effects of low density so widely.

Beyond these general trends, our GAMs revealed that 46/60 density effects on network connectedness (77%) were significantly nonlinear ($\Delta AIC > 2$); these relationships took a wide variety of shapes, representing a range of nonlinear functions that are hard to generalise (Figure 4; Supplementary Table 2). Notably, while many GAM smooths were eventually significantly negative (Figure 4), the vast majority of linear models fitted to the second half of the data were positive (Figure 3C); this result is likely an artefact of restricted model fitting, rather than true downturns in connectedness with density. Nonlinearity did not cluster according to connection type definitions, or according to host group. These observations were largely corroborated by our meta-analytical models, which found no factors influencing the slope and shape of density effects ($P > 0.05$). This observation speaks to the complexity of these relationships within and across systems, while accentuating that simple functional relationships are often likely to be complicated by contravening ecological factors like habitat selection [67,68], parasite avoidance [69], and demographic structuring [70]. While we were unable to identify specific between-system predictors of nonlinearity of density-connectedness relationships, the finding that most such relationships are strongly nonlinear is an important consideration for future work.

We acknowledge several limitations of our study, which we nevertheless believe could be remedied in the future. First, many of our social networks were formed of general spatiotemporal associations, and relatively few from specific social interactions – particularly those involving direct physical contact. Our current dataset could therefore benefit from supplementation with a broader range of direct interactions, particularly involving antagonism or bonding. For example, datasets concerning aggression or dominance interactions (e.g. [71]) or grooming alongside spatial behaviour could inform how density dependence affects the transmission of certain parasites such as mycobacteria [72] or tattoo skin disease [73].

The meta-dataset was also unevenly distributed across animal taxa (Figure 1): there were no primates or bony fishes and only one invertebrate, while rodents and ungulates were over-represented. These biases likely emerge through differences in data collection approaches: for example, although primate social behaviour is often studied with observations of direct interactions that could augment our data as described above (e.g. [74]), the spatial data required to build density distributions are rarely collected in these systems. This is linked to their social structures: our workflow was best suited to studies of fission-fusion societies or relatively asocial animals, rather than those with wide-ranging fixed social groups that are more common in primate systems. Finally, given that our data were observational, we could not account for (or estimate) bidirectional causality between density and social relationships (point 4 in Figure 1): that is, as well as encountering more conspecifics in areas of high density, individuals may be drawn to conspecifics, *creating* areas of high density [16]. To do so might require creating in-depth, high-resolution models of animal movement and group formation (e.g. [75]), potentially making use of telemetry approaches and drawing from large-scale open movement repositories like Movebank [76].

Aside from incorporating more specific interaction types, there exist a range of potential extensions to our analysis. For example, density dependence often varies between age or sex classes (e.g. [77,78]), and age effects on infection are common and ecologically important [79,80], as are sex differences [81]. We chose not to analyse how individual animals' traits alter the shape or slope of density's effects for brevity and simplicity; however, given that many of the systems nevertheless include these data, future analyses could make use of this meta-dataset to investigate how density affects connectedness of different classes of hosts. Further, researchers could investigate other behavioural questions such as the role of observation biases; the factors influencing the correlation between spatial and social networks; and the role of environmental drivers and spatial autocorrelation in driving observed patterns of connectivity [12,16]. Finally, as our analysis approximated density-contact relationships and not host-parasite interactions specifically, important future work could investigate whether contact rates (as approximated by network connectedness) ultimately translate to greater infection risk or parasite burden. Although some previous investigations have linked density-related metrics to aspects of infection [82,83], density covaries with a range of other factors including nutrition, cooperation, and competition, all of which could complicate density-driven increases in exposure [2]. For example, in the case of ectoparasite transmission, although contact rates in general would likely increase with density, so too might grooming behaviours that remove parasites; in cases such as these, density's overall effect on ectoparasite infection may be neutral. In the future, verifying that within- and between-population variation in density-contact relationships translate to variation in infection – and whether these trends might be influenced by flexible avoidance behaviours [37] – will be a vital part of understanding and predicting density-dependent disease dynamics.

Density is a universal factor underlying the dynamics of animal populations, and its linear and nonlinear effects on spatial and social network structure are likely to impact myriad processes in behaviour, ecology, and evolution. Similar to other studies that have reported general scaling patterns in network analysis [84] and in food web ecology [85], the patterns we report strongly suggest that animal systems generally become better connected spatially than socially under increasing density. These might extrapolate to human networks, given that other scaling patterns in animal networks do [84]. As these patterns seemingly manifest

regardless of animal group and interaction type, they may reflect a generalisable rule governing the socio-spatial structure of ecological systems. Further refining and implementing these models could facilitate prediction of network structure in novel systems.

Acknowledgements

GFA and SB were supported by NSF grant number DEB-2211287. GFA acknowledges support from a College for Life Sciences Fellowship at the Wissenschaftskolleg zu Berlin and WAI (C-2023-00057). DJB acknowledges support from the Edward Mallinckrodt, Jr. Foundation and NSF BII 2213854. JAF acknowledges funding from BBSRC (BB/S009752/1), NERC (NE/S010335/1 and NE/V013483/1), and WAI (C-2023-00057). For a list of system-specific acknowledgements, please see the supplement.

Code availability

All R code is available at <https://github.com/gfalbery/DensityMetaAnalysis>.

Affiliations

1. Department of Biology, Georgetown University, Washington, DC, USA
2. Wissenschaftskolleg zu Berlin, Berlin, Germany
3. School of Biological Sciences, University of Oklahoma, Norman, OK, USA
4. Department of Biology, University of Oxford, Oxford, UK
5. School of Biology, University of Leeds, Leeds, UK
6. CEFÉ, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
7. Centre for Ecology and Conservation, University of Exeter, Penryn, UK
8. School of Biosciences, University of Sheffield, Sheffield, UK
9. Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, UK
10. Department of Biology, Memorial University of Newfoundland, 45 Arctic Avenue, St. John's NL
11. University of Guelph, Guelph, Canada
12. Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK
13. Institute of Zoology, Zoological Society of London, Regents Park, London, UK
14. University of Glasgow, Glasgow, UK
15. Department of Biological Sciences, Old Dominion University, Norfolk, VA, USA
16. Avian Ecology Program, Archbold Biological Station, Venus, FL, USA
17. University of Salford, Salford, UK
18. Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA
19. Canine Companions, 2965 Dutton Ave, Santa Rosa, CA, USA
20. Hopkins Marine Station, Stanford University, Stanford, CA, USA
21. University of Nottingham, Nottingham, UK
22. Pitzer and Scripps Colleges, Claremont, CA, USA
23. Wildlife Conservation Research Unit, The Recanati-Kaplan Centre, Department of Zoology, University of Oxford, Oxford, UK
24. The University of British Columbia, Irving K. Barber Faculty of Sciences, Okanagan Department of Biology, Kelowna, British Columbia, Canada

- 535 25. UK Centre for Ecology and Hydrology, Bush Estate, Midlothian, UK
- 536 26. University of Delaware, Newark, DE, USA
- 537 27. Marine Science Institute, UC Santa Barbara, Santa Barbara, CA, USA
- 538 28. ISI Foundation, Torino, Italy
- 539 29. Evolutionary Ecology Group, Station d'Ecologie Theorique et Experimentale du CNRS,
- 540 Moulis, France
- 541 30. Oregon State University, Corvallis, OR, USA
- 542 31. Lighthouse Field Station, School of Biological Sciences, University of Aberdeen, UK
- 543 32. Department of Zoology, University of Cambridge, Cambridge, UK
- 544 33. Institute of Zoology, Zoological Society of London, London, UK
- 545 34. National Wildlife Management Centre, Animal and Plant Health Agency, Sand Hutton,
- 546 York, UK
- 547 35. Department of Collective Behavior, Max Planck Institute for Animal Behavior, 78464
- 548 Konstanz, Germany
- 549 36. Department of Evolutionary Biology and Environmental Studies, University of Zurich,
- 550 8057 Zurich, Switzerland
- 551 37. Department of Ornithology, National Museum of Kenya, P.O. Box 40241-001000,
- 552 Nairobi, Kenya
- 553 38. Division of Ecology and Evolution, Research School of Biology, Australian National
- 554 University, 46 Sullivan Creek Road, Canberra, ACT 2600, Australia
- 555 39. Virginia Tech, Blacksburg, VA, USA
- 556 40. The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK
- 557 41. Environment and Sustainability Institute, University of Exeter, Penryn, UK
- 558 42. Department of Marine Biology, Texas AandM University at Galveston, USA
- 559 43. School of the Environment, University of Queensland, Brisbane, Queensland, Australia
- 560 44. College of Science and Engineering, Flinders University, Adelaide, Australia
- 561 45. School of Zoology, Faculty of Life Science, Tel Aviv University, Tel Aviv, Israel
- 562 46. Department of Zoology, University of Otago, Otago, New Zealand
- 563 47. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ,
- 564 USA
- 565 48. Sea Mammal Research Unit, South Oceans Institute, University of St Andrews, UK
- 566 49. Department of Visitor Management and National Park Monitoring, Bavarian Forest
- 567 National Park, Grafenau, Germany
- 568 50. Chair of Wildlife Ecology and Wildlife Management, Faculty of Environment and Natural
- 569 Resources, University of Freiburg, Freiburg, Germany
- 570 51. Department of Forestry and Wildlife Management, Faculty of Applied Ecology,
- 571 Agricultural Sciences and Biotechnology, Campus Evenstad, Inland Norway University of
- 572 Applied Sciences, Postboks 400 Vestad, 2418 Elverum, Norway
- 573 52. Faculty of Life Sciences, Bar Ilan University, Ramat Gan, Israel
- 574 53. Lancaster University, Lancaster, UK
- 575 54. Division of Marine Science and Conservation, Nicholas School of the Environment, Duke
- 576 University Marine Laboratory, Beaufort, NC, USA
- 577 55. Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood
- 578 Sciences, Czech University of Life Sciences, Prague, Czech Republic
- 579 56. Evolutionary Ecology Group, EVECO, Universiteit Antwerpen, Antwerp, Belgium
- 580 57. Department of Nature Conservation and Research, Hainich National Park, Bad
- 581 Langensalza, Germany
- 582 58. Mpala Research Center, P.O Box 555-10400, Nanyuki, 10400, Kenya

- 583 59. Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78464
584 Konstanz, Germany
- 585 60. School of Environment and Natural Science, Bangor University, Bangor, Wales
- 586 61. Department of Biology, Memorial University of Newfoundland, St John's, NL, Canada
- 587 62. Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada
- 588 63. University of Aberdeen, Aberdeen, UK
- 589 64. School of Animal and Veterinary Sciences, The University of Adelaide, Adelaide,
590 Australia
- 591 65. School of Biological Sciences, University of Nebraska-Lincoln, NE, USA
- 592 66. Department of Ecology, Behavior, and Evolution, University of California, San Diego, CA
- 593 67. Department of Evolutionary Biology and Environmental Studies, University of Zurich,
594 Zurich, Switzerland
- 595 68. Institute of Pest Management, Sokoine University of Agriculture, Morogoro, Tanzania
- 596 69. Max Planck Institute of Animal Behavior, Radolfzell, Germany
- 597 70. MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France
- 598 71. PSL Research University, Université de Perpignan, Perpignan, France
- 599 72. Laboratoire d'Excellence "CORAIL", Papetoai, Moorea, French Polynesia
- 600 73. University of Nevada, Reno, NV, USA
- 601 74. University of Eldoret, School of Natural Resource Management, Department of Wildlife,
602 1125-30100 Eldoret, Kenya
- 603 75. Department of Veterinary Sciences, University of Turin, Grugliasco (TO), Italy
- 604 76. Florida International University, FL, USA
- 605 77. University of Liverpool, Liverpool, UK
- 606 78. Department of Environmental Science and Policy, University of California Davis, Davis,
607 CA, USA
- 608 79. Department of Ecology and Evolutionary Biology, University of California, Los Angeles,
609 CA, USA
- 610 80. Stanford University, Stanford, CA, USA
- 611 81. School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA
- 612 82. Department of Wildlife, California State Polytechnic University, Humboldt, Arcata, CA,
613 USA
- 614 83. Texas Tech University, Lubbock, TX, USA
- 615 84. University of Western Australia, Crawley WA 6009, Australia
- 616 85. Department of Veterinary Population Medicine, University of Minnesota, Minneapolis, US
- 617 86. University of Surrey, Guildford, UK

618 References

- 619 1. Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latto J, Sanchez MS,
620 Smith AB, Swei A. 2005 Should we expect population thresholds for wildlife disease?
621 *Trends Ecol Evol* **20**, 511–519. (doi:10.1016/j.tree.2005.07.004)
- 622 2. Albery GF. 2022 Density dependence and disease dynamics: moving towards a
623 predictive framework. *EcoEvoRxiv Preprints* (doi:10.32942/OSF.IO/GAW49)
- 624 3. Hopkins SR, Fleming-Davies AE, Belden LK, Wojdak JM. 2020 Systematic review of
625 modeling assumptions and empirical evidence: does parasite transmission increase

- 626 nonlinearly with host density? *Methods Ecol Evol* **11**, 0–2. (doi:10.1111/2041-
627 210x.13361)
- 628 4. Begon M, Bennett M, Bowers RG, French NP, Hazel SM, Turner J. 2002 A
629 clarification of transmission terms in host-microparasite models: Numbers, densities
630 and areas. *Epidemiol Infect* **129**, 147–153. (doi:10.1017/S0950268802007148)
- 631 5. Ferrari MJ, Perkins SE, Pomeroy LW, Bjrnstad ON. 2011 Pathogens, social networks,
632 and the paradox of transmission scaling. *Interdiscip Perspect Infect Dis* **2011**.
633 (doi:10.1155/2011/267049)
- 634 6. Webber QMR, Vander Wal E. 2020 Heterogeneity in social network connections is
635 density-dependent: implications for disease dynamics in a gregarious ungulate. *Behav*
636 *Ecol Sociobiol* **74**, 1–13.
- 637 7. Beauchamp G. 2011 Functional relationship between group size and population
638 density in Northwest Atlantic seabirds. *Mar Ecol Prog Ser* **435**, 225–233.
639 (doi:10.3354/meps09239)
- 640 8. Borremans B, Reijniers J, Hens N, Leirs H. 2017 The shape of the contact–density
641 function matters when modelling parasite transmission in fluctuating populations. *R*
642 *Soc Open Sci* **4**. (doi:10.1098/rsos.171308)
- 643 9. Davis S, Abbasi B, Shah S, Telfer S, Begon M. 2015 Spatial analyses of wildlife
644 contact networks. *J R Soc Interface* **12**. (doi:10.1098/rsif.2014.1004)
- 645 10. Carson BD, Orians CM, Crone EE. 2023 When does spatial clustering in caterpillars
646 inuence the relationship between population density and contact rates?
647 (doi:10.21203/rs.3.rs-2842538/v1)
- 648 11. Mancy R *et al.* 2022 Rabies shows how scale of transmission can enable acute
649 infections to persist at low prevalence. *Science (1979)* **376**, 512–516.
- 650 12. Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, Firth
651 JA. 2021 Multiple spatial behaviours govern social network positions in a wild
652 ungulate. *Ecol Lett* **24**, 676–686. (doi:10.1101/2020.06.04.135467)
- 653 13. Strickland K, Patterson EM, Frère CH. 2018 Eastern water dragons use alternative
654 social tactics at different local densities. *Behav Ecol Sociobiol* **72**.
655 (doi:10.1007/s00265-018-2563-x)
- 656 14. Strickland K, Frère CH. 2019 Individual Variation in the Social Plasticity of Water
657 Dragons. *Am Nat* **194**, 194–206. (doi:10.5061/dryad.5j5gh63)
- 658 15. Hu H, Nigmatulina K, Eckhoff P. 2013 The scaling of contact rates with population
659 density for the infectious disease models. *Math Biosci* **244**, 125–134.
660 (doi:10.1016/j.mbs.2013.04.013)

- 661 16. Webber QMR, Albery GF, Farine DR, Pinter-wollman N, Sharma N, Vander Wal E,
662 Manlove K. 2023 Behavioural ecology at the spatial-social interface. *Biological*
663 *Reviews* , 1–40.
- 664 17. Albon SD, Staines HJ, Guinness FE, Clutton-Brock TH. 1992 Density-Dependent
665 Changes in the Spacing Behaviour of Female Kin in Red Deer. *Source: Journal of*
666 *Animal Ecology*. **61**.
- 667 18. Vas J, Andersen IL. 2015 Density-dependent spacing behaviour and activity budget in
668 pregnant, domestic goats (*Capra hircus*). *PLoS One* **10**.
669 (doi:10.1371/journal.pone.0144583)
- 670 19. Connor EF, Cargain MJ. 1994 Density-related foraging behaviour in *Closterocerus*
671 *tricinctus*, a parasitoid of the leaf-mining moth, *Cameraria hamadryadella*. *Ecol*
672 *Entomol* **19**, 327–334. (doi:10.1111/J.1365-2311.1994.TB00249.X)
- 673 20. Matthysen E. 2005 Density-dependent dispersal in birds and mammals. *Ecography*
674 **28**, 403–416. (doi:10.1111/j.0906-7590.2005.04073.x)
- 675 21. Avgar T, Betini GS, Fryxell JM. 2020 Habitat selection patterns are density dependent
676 under the ideal free distribution. *Journal of Animal Ecology* **89**, 2777–2787.
677 (doi:10.1111/1365-2656.13352)
- 678 22. Vander Wal E, Laforge MP, Mcloughlin PD, Wal E Vander, Laforge MP, Mcloughlin
679 PD, Vander Wal E, Laforge MP, Mcloughlin PD. 2014 Density dependence in social
680 behaviour: Home range overlap and density interacts to affect conspecific encounter
681 rates in a gregarious ungulate. *Behav Ecol Sociobiol* **68**, 383–390.
682 (doi:10.1007/s00265-013-1652-0)
- 683 23. Kokko H, Rankin DJ. 2006 Lonely hearts or sex in the city? Density-dependent effects
684 in mating systems. In *Philosophical Transactions of the Royal Society B: Biological*
685 *Sciences*, pp. 319–334. Royal Society. (doi:10.1098/rstb.2005.1784)
- 686 24. Modlmeier AP, Colman E, Hanks EM, Bringenberg R, Bansal S, Hughes DP. 2019
687 Ant colonies maintain social homeostasis in the face of decreased density. *Elife* **8**, 1–
688 17. (doi:10.7554/eLife.38473)
- 689 25. Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting animal
690 social network analysis. *Journal of Animal Ecology* **84**, 1144–1163.
691 (doi:10.1111/1365-2656.12418)
- 692 26. Farine DR. 2024 Modelling animal social networks: New solutions and future
693 directions. *Journal of Animal Ecology* (doi:10.1111/1365-2656.14049)
- 694 27. Webber QMR, Vander Wal E. 2019 Trends and perspectives on the use of animal
695 social network analysis in behavioural ecology: a bibliometric approach. *Anim Behav*
696 **149**, 77–87. (doi:10.1016/j.anbehav.2019.01.010)

- 697 28. Pinter-Wollman N *et al.* 2014 The dynamics of animal social networks: Analytical,
698 conceptual, and theoretical advances. *Behavioral Ecology* **25**, 242–255.
699 (doi:10.1093/beheco/art047)
- 700 29. Sah P, Mann J, Bansal S. 2018 Disease implications of animal social network
701 structure: A synthesis across social systems. *Journal of Animal Ecology* **87**, 546–558.
702 (doi:10.1111/1365-2656.12786)
- 703 30. Albery GF, Kirkpatrick L, Firth JA, Bansal S. 2021 Unifying spatial and social network
704 analysis in disease ecology. *Journal of Animal Ecology* **90**, 1–17. (doi:10.1111/1365-
705 2656.13356)
- 706 31. Cubaynes S, Macnulty DR, Stahler DR, Quimby KA, Smith DW, Coulson T. 2014
707 Density-dependent intraspecific aggression regulates survival in northern Yellowstone
708 wolves (*Canis lupus*). *Journal of Animal Ecology* **83**, 1344–1356. (doi:10.1111/1365-
709 2656.12238)
- 710 32. Lloyd-Smith JO, Getz WM, Westerhoff H V. 2004 Frequency-dependent incidence in
711 models of sexually transmitted diseases: Portrayal of pair-based transmission and
712 effects of illness on contact behaviour. *Proceedings of the Royal Society B: Biological*
713 *Sciences* **271**, 625–634. (doi:10.1098/rspb.2003.2632)
- 714 33. Morders MK, Restif O, Hampson K, Cleaveland S, Wood JLN, Conlan AJK. 2013
715 Evidence-based control of canine rabies: A critical review of population density
716 reduction. *Journal of Animal Ecology* **82**, 6–14. (doi:10.1111/j.1365-
717 2656.2012.02033.x)
- 718 34. Gordon DM, Paul RE, Thorpe K. 1993 What is the function of encounter patterns in
719 ant colonies? *Anim Behav* **45**, 1083–1100. (doi:10.1006/ANBE.1993.1134)
- 720 35. Davis S, Abbasi B, Shah S, Telfer S, Begon M, Davis S. 2015 Spatial analyses of
721 wildlife contact networks. *Journal of the Royal Society, Interface / the Royal Society*
722 **12**, 20141004. (doi:10.1098/rsif.2014.1004)
- 723 36. R Development Core Team R. 2011 R: A Language and Environment for Statistical
724 Computing. *R Foundation for Statistical Computing*. **1**, 409. (doi:10.1007/978-3-540-
725 74686-7)
- 726 37. Albery GF, Newman C, Bright Ross J, Macdonald DW, Bansal S, Buesching CD.
727 2020 Negative density-dependent parasitism in a group-living carnivore. *Proceedings*
728 *of the Royal Society B: Biological Sciences* **287**, 20202655.
729 (doi:10.1101/2020.06.15.153726)
- 730 38. Calenge C. 2011 Home range estimation in R: the adehabitatHR package. See
731 <https://cran.r-project.org/web/packages/adehabitatHR/index.html> (accessed on 10
732 March 2020).
- 733 39. Franks DW, Ruxton GD, James R. 2010 Sampling animal association networks with
734 the gambit of the group. *Behav Ecol Sociobiol* **64**, 493–503. (doi:10.1007/s00265-009-
735 0865-8)

- 736 40. Wanelik KM, Farine DR. 2022 A new method for characterising shared space use
737 networks using animal trapping data. *Behav Ecol Sociobiol* **76**. (doi:10.1007/s00265-
738 022-03222-5)
- 739 41. Cairns SJ, Schwager SJ. 1987 A comparison of association indices. *Anim Behav* **35**,
740 1454–1469. (doi:10.1016/S0003-3472(87)80018-0)
- 741 42. Wood SN. 2011 Fast stable restricted maximum likelihood and marginal likelihood
742 estimation of semiparametric generalized linear models. *J R Stat Soc Series B Stat*
743 *Methodol* **73**, 3–36. (doi:10.1111/j.1467-9868.2010.00749.x)
- 744 43. Koricheva J, Gurevitch J, Mengersen KL. 2013 Handbook of meta-analysis in ecology
745 and evolution. , 498.
- 746 44. Konstantopoulos S. 2011 Fixed effects and variance components estimation in three-
747 level meta-analysis. *Res Synth Methods* **2**, 61–76. (doi:10.1002/jrsm.35)
- 748 45. Cinar O, Nakagawa S, Viechtbauer W. 2022 Phylogenetic multilevel meta-analysis: A
749 simulation study on the importance of modelling the phylogeny. *Methods Ecol Evol*
750 **13**, 383–395. (doi:10.1111/2041-210X.13760)
- 751 46. Michonneau F, Brown JW, Winter DJ. 2016 rotl: an R package to interact with the
752 Open Tree of Life data. *Methods Ecol Evol* **7**, 1476–1481. (doi:10.1111/2041-
753 210X.12593)
- 754 47. Paradis E, Schliep K. 2019 ape 5.0: an environment for modern phylogenetics and
755 evolutionary analyses in R. *Bioinformatics* **35**, 526–528.
756 (doi:10.1093/BIOINFORMATICS/BTY633)
- 757 48. Gibson AK, Amoroso CR. 2022 Evolution and Ecology of Parasite Avoidance. , 1–21.
- 758 49. Colman E, Colizza V, Hanks EM, Hughes DP, Bansal S. 2021 Social fluidity mobilizes
759 contagion in human and animal populations. *Elife* **10**. (doi:10.7554/ELIFE.62177)
- 760 50. Michelangeli M, Payne E, Spiegel O, Sinn D, Leu ST, Gardner M, Sih A. 2021
761 *Personality, spatiotemporal ecological variation, and resident/explorer movement*
762 *syndromes in the sleepy lizard*. (doi:10.1111/1365-2656.13616)
- 763 51. Spiegel O, Leu ST, Bull CM, Sih A. 2017 What's your move? Movement as a link
764 between personality and spatial dynamics in animal populations. *Ecol Lett* **20**, 3–18.
765 (doi:10.1111/ele.12708)
- 766 52. Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014 Individual-level personality
767 influences social foraging and collective behaviour in wild birds. *Proceedings of the*
768 *Royal Society of London B* **281**, 20141016. (doi:10.1098/rspb.2014.1016)
- 769 53. Hunt ER, Mi B, Fernandez C, Wong BM, Pruitt JN, Pinter-Wollman N. 2018 Social
770 interactions shape individual and collective personality in social spiders. *Proceedings*
771 *of the Royal Society B: Biological Sciences* **285**, 20181366.
772 (doi:10.1098/rspb.2018.1366)

- 773 54. Firth JA, Cole EF, Ioannou CC, Quinn JL, Aplin LM, Culina A, McMahon K, Sheldon
774 BC. 2018 Personality shapes pair bonding in a wild bird social system. *Nat Ecol Evol*
775 **2**, 1696–1699. (doi:10.1038/s41559-018-0670-8)
- 776 55. Webber Q, Laforge M, Bonar M, Wal E Vander. 2022 The adaptive value of density-
777 dependent habitat specialization and social network centrality. *Authorea Preprints*
778 (doi:10.22541/AU.165210484.46672609/V1)
- 779 56. Webber QMR, Dantzer B, Lane JE, Boutin S, McAdam AG. 2023 Density-dependent
780 plasticity in territoriality revealed using social network analysis. *Journal of Animal*
781 *Ecology* **92**, 207–221. (doi:10.1111/1365-2656.13846)
- 782 57. Sheppard CE, Inger R, McDonald RA, Barker S, Jackson AL, Thompson FJ,
783 Vitikainen EIK, Cant MA, Marshall HH. 2018 Intragroup competition predicts individual
784 foraging specialisation in a group-living mammal. *Ecol Lett* **21**, 665–673.
785 (doi:10.1111/ELE.12933)
- 786 58. Sanchez JN, Hudgens BR. 2015 Interactions between density, home range
787 behaviors, and contact rates in the Channel Island fox (*Urocyon littoralis*). *Ecol Evol* **5**,
788 2466–2477. (doi:10.1002/ece3.1533)
- 789 59. Wilkin TA, Garant D, Gosler AG, Sheldon BC. 2006 Density effects on life-history
790 traits in a wild population of the great tit *Parus major*: Analyses of long-term data with
791 GIS techniques. *Journal of Animal Ecology* **75**, 604–615. (doi:10.1111/j.1365-
792 2656.2006.01078.x)
- 793 60. Cote IM, Poulin R. 1995 Parasitism and group size in social animals: a meta-analysis.
794 *Behavioral Ecology* **6**, 159–165. (doi:10.1093/beheco/6.2.159)
- 795 61. McCallum H, Barlow N, Hone J. 2001 How should pathogen transmission be
796 modelled? *Trends Ecol Evol* **16**, 295–300. (doi:10.1016/S0169-5347(01)02144-9)
- 797 62. Rajeev M, Metcalf CJE, Hampson K. 2020 Modeling canine rabies virus transmission
798 dynamics. In *Rabies*, pp. 655–670. Elsevier. (doi:10.1016/b978-0-12-818705-
799 0.00020-0)
- 800 63. Townsend SE *et al.* 2013 Designing Programs for Eliminating Canine Rabies from
801 Islands: Bali, Indonesia as a Case Study. *PLoS Negl Trop Dis* **7**.
802 (doi:10.1371/JOURNAL.PNTD.0002372)
- 803 64. Macdonald DW, Newman C. 2022 *The Badgers of Wytham Woods: a model for*
804 *behaviour, ecology, and evolution*. Oxford University Press.
- 805 65. Clutton-Brock TH, Guinness FE, Albon SD. 1982 *Red Deer: Behavior and Ecology of*
806 *Two Sexes*. Chicago, IL: University of Chicago Press. See
807 https://books.google.co.uk/books/about/Red_Deer.html?id=x4SGuA3t-NoC&pgis=1.
- 808 66. Clutton-Brock TH, Pemberton JM. 2004 *Soay sheep: population dynamics and*
809 *selection on St. Kilda*. Cambridge, UK: Cambridge University Press.

- 810 67. Tardy O, Massé A, Pelletier F, Mainguy J, Fortin D. 2014 Density-dependent
811 functional responses in habitat selection by two hosts of the raccoon rabies virus
812 variant. *Ecosphere* **5**, 1–16. (doi:10.1890/ES14-00197.1)
- 813 68. Wilber MQ, Yang A, Boughton R, Manlove KR, Miller RS, Pepin KM, Wittemyer G.
814 2022 A model for leveraging animal movement to understand spatio-temporal disease
815 dynamics. *Ecol Lett* , 1–15. (doi:10.1111/ele.13986)
- 816 69. Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary Consequences
817 of Parasite Avoidance. *Trends Ecol Evol* **33**, 619–632.
818 (doi:10.1016/j.tree.2018.05.001)
- 819 70. Lion S, Boots M. 2010 Are parasites “prudent” in space? *Ecol Lett* **13**, 1245–1255.
820 (doi:10.1111/j.1461-0248.2010.01516.x)
- 821 71. Strauss ED, Decasien AR, Galindo G, Hobson EA, Shizuka D, Curley JP. 2022
822 DomArchive: a century of published dominance data. *Philosophical Transactions of*
823 *the Royal Society B* **377**. (doi:10.1098/RSTB.2020.0436)
- 824 72. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in
825 wild meerkats. *Proceedings. Biological sciences / The Royal Society* **277**, 633–42.
826 (doi:10.1098/rspb.2009.1775)
- 827 73. Powell SN, Wallen MM, Miketa ML, Krzyszczyk E, Foroughirad V, Bansal S, Mann J.
828 2021 Sociality and tattoo skin disease among bottlenose dolphins in Shark Bay,
829 Australia. *Behavioral Ecology* **31**, 459–466. (doi:10.1093/BEHECO/ARZ207)
- 830 74. Testard C *et al.* 2020 Rhesus Macaques Build New Social Connections after a Natural
831 Disaster. *Current Biology* , 1–11. (doi:10.2139/ssrn.3741230)
- 832 75. Spiegel O, Leu ST, Sih A, Bull CM. 2016 Socially interacting or indifferent
833 neighbours? Randomization of movement paths to tease apart social preference
834 and spatial constraints. *Methods Ecol Evol* , 971–979. (doi:10.1111/2041-
835 210X.12553)
- 836 76. Kays R *et al.* 2021 The Movebank system for studying global animal movement and
837 demography. *Methods Ecol Evol* **2021**, 1–13. (doi:10.1111/2041-210X.13767)
- 838 77. Batsuren E, Zhang X, Song M, Wan X, Li G, Liu J, Huang S, Zhang Z. 2022 Density-
839 dependent changes of mating system and family structure in Brandt's voles
840 (*Lasiopodomys brandtii*). *Ecol Evol* , 1–16. (doi:10.1002/ece3.9199)
- 841 78. Vander Wal E, van Beest FM, Brook RK. 2013 Density-Dependent Effects on Group
842 Size Are Sex-Specific in a Gregarious Ungulate. *PLoS One* **8**, e53777.
843 (doi:10.1371/JOURNAL.PONE.0053777)
- 844 79. Ashby B, Bruns E. 2018 The evolution of juvenile susceptibility to infectious disease.
845 *Proceedings of the Royal Society B: Biological Sciences* **285**, 20180844.
846 (doi:10.1098/rspb.2018.0844)

847 80. Albery GF, Sweeny AR, Webber Q. In press. How behavioural ageing affects
848 infectious disease.

849 81. Zuk M. 2009 The sicker sex. *PLoS Pathog* **5**, 1–3. (doi:10.1371/journal.ppat.1000267)

850 82. Arneberg P, Skorping A, Grenfell B, Read AF. 1998 Host densities as determinants of
851 abundance in parasite communities. *Proceedings of the Royal Society B: Biological*
852 *Sciences* **265**, 1283–1289. (doi:10.1098/rspb.1998.0431)

853 83. Arneberg P. 2001 An ecological law and its macroecological consequences as
854 revealed by studies of relationships between host densities and parasite prevalence.
855 *Ecography* **24**, 352–358. (doi:10.1111/j.1600-0587.2001.tb00208.x)

856 84. Rocha LEC, Ryckebusch J, Schoors K, Smith M. 2021 Scaling of social interactions
857 across animal species. *Sci Rep* **11**, 12584.

858 85. Lafferty KD, DeLeo G, Briggs CJ, Dobson AP, Gross T, Kuris AM. 2015 A general
859 consumer-resource population model. *Science (1979)* **349**, 854–857.
860 (doi:10.1126/SCIENCE.AAA6224/SUPPL_FILE/LAFFERTY.SM.PDF)

861