

AFTER VISIT SUMMARY

Melody D. Dashora MRN: 2000000307
CEID: THR-2SVP-22F2-QQQR
EMERGENCY DEPARTMENT

8/30/2026 Texas Health Frisco Emergency
Department 469-495-2070

Instructions



Read the attached information

1. General Headache Without Cause (English)
2. Motor Vehicle Collision Injury Adult (English)
3. Cervical Strain and Sprain Rehab (English)



Follow up with Sarah Uddin in 3 days (around 9/2/2026)

Contact: 19 Yawpo Avenue
Oakland NJ 07436
201-337-9600



Follow up with Texas Health Frisco Emergency Department

Why: As needed, If symptoms worsen
Specialty: Emergency Room
Contact: 12400 N. Dallas Parkway
Frisco Texas 75033
469-495-2070

Today's Visit

You were seen by Shannon Kelsey Della Monica

Reason for Visit

Motor Vehicle Crash

Diagnoses

- Motor vehicle collision, subsequent encounter
- Strain of neck muscle, subsequent encounter
- Acute nonintractable headache, unspecified headache type

Lab Tests Completed

CBC WITH DIFFERENTIAL
COMPREHENSIVE METABOLIC PANEL
MAGNESIUM
PREGNANCY TEST, SERUM

Imaging Tests

CT Head, WO IV Contrast (CT HEAD WO CON)
CTA Head & Neck Angio Combo (CTA HEAD NECK ANGIO COMBO)

Today's Visit (continued)

Medications Given

fentaNYL PF (Sublimaze) Last given at 8:09 PM
iohexol (Omnipaque-350) Last given at 8:15 PM
ketorolac (Toradol) Last given at 9:38 PM
NORMAL SALINE 1,000 mL Stopped at 9:38 PM
Normal Saline
ondansetron (Zofran) Last given at 8:09 PM



Blood Pressure
109/68



Temperature
(Temporal Artery)
97.2 °F



Pulse
70



Respiration
16



Oxygen Saturation
96%

What's Next

You currently have no upcoming appointments scheduled.

Changes to Your Medication List

You have not been prescribed any medications.



Texas Health
Resources®

MyChart

Access your Health Information with MyChart

Your MyChart account is already active. Log in to view your health records, test results, and more.

Forgot your login information? Visit <https://mychart.texashealth.org> click Sign in, then choose *Forgot Username* or *Forgot Password* to recover your account details.

Want to give someone access to your MyChart account? Visit the "*Need proxy access to a family or friend's health record?*" section and click the link titled [Here's how](#) to get started.

General Headache Without Cause

A headache is pain or discomfort felt around the head or neck area. There are many causes and types of headaches. A few common types include:

- Tension headaches.
- Migraine headaches.
- Cluster headaches.
- Chronic daily headaches.

Sometimes, the specific cause of a headache may not be found.

Follow these instructions at home:

Watch your condition for any changes. Let your health care provider know about them. Take these steps to help with your condition:

Managing pain

- Take over-the-counter and prescription medicines only as told by your health care provider. Treatment may include medicines for pain that are taken by mouth or applied to the skin.
- Lie down in a dark, quiet room when you have a headache.
- Keep lights dim if bright lights bother you or make your headaches worse.
- If directed, put ice on your head and neck area:
 - Put ice in a plastic bag.
 - Place a towel between your skin and the bag.
 - Leave the ice on for 20 minutes, 2–3 times per day.
 - Remove the ice if your skin turns bright red. This is very important. If you cannot feel pain, heat, or cold, you have a greater risk of damage to the area.
- If directed, apply heat to the affected area. Use the heat source that your health care provider recommends, such as a moist heat pack or a heating pad.
 - Place a towel between your skin and the heat source.
 - Leave the heat on for 20–30 minutes.
 - Remove the heat if your skin turns bright red. This is especially important if you are unable to feel pain, heat, or cold. You have a greater risk of getting burned.

Eating and drinking

- Eat meals on a regular schedule.
- If you drink alcohol:
 - Limit how much you have to:
 - 0–1 drink a day for women who are not pregnant.
 - 0–2 drinks a day for men.
 - Know how much alcohol is in a drink. In the U.S., one drink equals one 12 oz bottle of beer (355 mL), one 5 oz glass of wine (148 mL), or one 1½ oz glass of hard liquor (44 mL).
- Stop drinking caffeine, or decrease the amount of caffeine you drink.
- Drink enough fluid to keep your urine pale yellow.

General instructions

- Keep a headache journal to help find out what may trigger your headaches. For example, write down:
 - What you eat and drink.
 - How much sleep you get.
 - Any change to your diet or medicines.
- Try massage or other relaxation techniques.
- Limit stress.
- Sit up straight, and **do not** tense your muscles.
- **Do not** use any products that contain nicotine or tobacco. These products include cigarettes, chewing tobacco, and vaping devices, such as e-cigarettes. If you need help quitting, ask your health care provider.
- Exercise regularly as told by your health care provider.
- Sleep on a regular schedule. Get 7–9 hours of sleep each night, or the amount recommended by your health care provider.
- Keep all follow-up visits. This is important.

Contact a health care provider if:

- Medicine does not help your symptoms.
- You have a headache that is different from your usual headache.
- You have nausea or you vomit.
- You have a fever.

Get help right away if:

- Your headache:
 - Becomes severe quickly.
 - Gets worse after moderate to intense physical activity.
- You have any of these symptoms:
 - Repeated vomiting.
 - Pain or stiffness in your neck.
 - Changes to your vision.
 - Pain in an eye or ear.
 - Problems with speech.
 - Muscular weakness or loss of muscle control.
 - Loss of balance or coordination.
- You feel faint or pass out.
- You have confusion.
- You have a seizure.

These symptoms may represent a serious problem that is an emergency. Do not wait to see if the symptoms will go away. Get medical help right away. Call your local emergency services (911 in the U.S.). Do not drive yourself to the hospital.

Summary

- A headache is pain or discomfort felt around the head or neck area.
- There are many causes and types of headaches. In some cases, the cause may not be found.
- Keep a headache journal to help find out what may trigger your headaches. Watch your condition for any changes. Let your health care provider know about them.
- Contact a health care provider if you have a headache that is different from the usual headache, or if your symptoms are not helped by medicine.
- Get help right away if your headache becomes severe, you vomit, you have a loss of vision, you lose your balance, or you have a seizure.

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 05/18/2022 Document Reviewed: 05/18/2022
Elsevier Patient Education © 2025 Elsevier Inc.

Motor Vehicle Collision Injury, Adult

After a motor vehicle collision, it is common to have injuries to the head, face, arms, and body. These injuries may include cuts, burns, and bruises. The collision can also cause sore muscles, muscle strains, headaches, and broken bones.

You may have stiffness and soreness for the first several hours. You may feel worse after waking up the first morning after the collision. These injuries tend to feel worse for the first 24–48 hours. Your injuries should then begin to improve with each day. How quickly you improve often depends on:

- The severity of the collision.
- The number of injuries you have.
- The location and nature of the injuries.
- Whether you were wearing a seat belt and whether your airbag deployed.

A head injury may result in a concussion, which is a brain injury that can have serious effects. If you have a concussion, you should rest as told by your health care provider. You must be very careful to avoid having a second concussion.

Follow these instructions at home:

Medicines

- Take over-the-counter and prescription medicines only as told by your health care provider.
- If you were prescribed antibiotics, take or apply it as told by your health care provider. **Do not** stop using the antibiotic even if you start to feel better.

Wound or burn care

Follow instructions from your health care provider about how to take care of your wound or burn. Make sure you:

- Clean your wound or burn. To do this:
 - Wash it with mild soap and water.
 - Rinse it with water to remove all soap.
 - Pat it dry with a clean towel. **Do not** rub it.

- Put an ointment or cream on the wound, if you were told to do so.
- Know when and how to change or remove your bandage (*dressing*). Always wash your hands with soap and water for at least 20 seconds before and after you change your dressing. If soap and water are not available, use hand sanitizer.
- Leave any stitches (*sutures*), skin glue, or adhesive strips in place. These skin closures may need to stay in place for 2 weeks or longer. If adhesive strip edges start to loosen and curl up, you may trim the loose edges. **Do not** remove adhesive strips completely unless your health care provider tells you to do that.
- Avoid exposing your burn or wound to the sun.
- Keep the surface of the wound or burn intact.
 - **Do not** scratch or pick at the wound or burn.
 - **Do not** break any blisters you may have.
 - **Do not** peel any skin.
- Check your wound or burn every day for signs of infection. Check for:
 - Redness, swelling, or pain.
 - Fluid or blood.
 - Warmth.
 - Pus or a bad smell.

Managing pain, stiffness, and swelling

- If directed, put ice on the injured areas. This can help with pain and swelling. To do this:
 - Put ice in a plastic bag.
 - Place a towel between your skin and the bag.
 - Leave the ice on for 20 minutes, 2–3 times a day.
 - If your skin turns bright red, remove the ice right away to prevent skin damage. The risk of skin damage is higher if you cannot feel pain, heat, or cold.
 - Raise (*elevate*) the wound or burn above the level of your heart while you are sitting or lying down. This will help reduce pain, pressure, and swelling.
 - If you have a wound or burn on your face, you may want to sleep with your head elevated. You may do this by putting an extra pillow under your head.

Activity

- Rest. Rest helps your body to heal. Make sure you:
 - Get plenty of sleep at night. Avoid staying up late.
 - Keep the same bedtime hours on weekends and weekdays.
- You may have to avoid lifting. Ask your health care provider how much you can safely lift. Lifting can make neck or back pain worse.
- Ask your health care provider when you can drive, ride a bicycle, or use machinery. Your ability to react may be slower if you injured your head. **Do not** do these activities if you are dizzy.

General instructions

- If you have a splint, brace, or sling, follow your health care provider's instructions on how to use your device.
- Drink enough fluid to keep your urine pale yellow.
- **Do not** drink alcohol.
- Eat a healthy diet. Ask your health care provider what foods you should eat.

Contact a health care provider if:

- You have any new or worsening symptoms, such as:
 - A worsening headache
 - Pain or swelling in an arm or leg.
 - Numbness, tingling, or weakness in your arms or legs.
 - Trouble moving an arm or leg.
 - New neck or back pain.
 - Nausea or vomiting
- You have signs of infection in a wound or burn.
- You have a fever.
- You have a head injury and any of the following symptoms for more than 2 weeks after your motor vehicle collision:
 - Headaches that do not go away.
 - Dizziness or balance problems.
 - Nausea or vomiting.
 - Increased sensitivity to noise or light.
 - Depression, anxiety, or irritability and mood swings.

- Memory problems or trouble concentrating.
- Sleep problems or feeling more tired than usual.
- You have changes in bowel or bladder control.
- You have blood in your urine, stool, or you vomit.

Get help right away if:

- You have increasing pain in the chest, neck, back, or abdomen.
- You have shortness of breath.

These symptoms may be an emergency. Get help right away. Call 911.

- **Do not wait to see if the symptoms will go away.**
- **Do not drive yourself to the hospital.**

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 06/12/2023 Document Reviewed: 06/12/2023
Elsevier Patient Education © 2025 Elsevier Inc.

Cervical Strain and Sprain Rehab

Ask your health care provider which exercises are safe for you. Do exercises exactly as told by your health care provider and adjust them as directed. It is normal to feel mild stretching, pulling, tightness, or discomfort as you do these exercises. Stop right away if you feel sudden pain or your pain gets worse. **Do not** begin these exercises until told by your health care provider.

Stretching and range-of-motion exercises

Cervical side bending

1. Using good posture, sit on a stable chair or stand up.
2. Without moving your shoulders, slowly tilt your left / right ear to your shoulder until you feel a stretch in the neck muscles on the opposite side. You should be looking straight ahead.
3. Hold for _____ seconds.
4. Repeat with the other side of your neck.

Repeat _____ times. Complete this exercise _____ times a day.

Cervical rotation

1. Using good posture, sit on a stable chair or stand up.
2. Slowly turn your head to the side as if you are looking over your left / right shoulder.
 - Keep your eyes level with the ground.
 - Stop when you feel a stretch along the side and the back of your neck.
3. Hold for _____ seconds.
4. Repeat this by turning to your other side.

Repeat _____ times. Complete this exercise _____ times a day.

Thoracic extension and pectoral stretch

1. Roll a towel or a small blanket so it is about 4 inches (10 cm) in diameter.
2. Lie down on your back on a firm surface.

3. Put the towel in the middle of your back across your spine. It should not be under your shoulder blades.
4. Put your hands behind your head and let your elbows fall out to your sides.
5. Hold for _____ seconds.

Repeat _____ times. Complete this exercise _____ times a day.

Strengthening exercises

Upper cervical flexion

1. Lie on your back with a thin pillow behind your head or a small, rolled-up towel under your neck.
2. Gently tuck your chin toward your chest and nod your head down to look toward your feet. **Do not** lift your head off the pillow.
3. Hold for _____ seconds.
4. Release the tension slowly. Relax your neck muscles completely before you repeat this exercise.

Repeat _____ times. Complete this exercise _____ times a day.

Cervical extension

1. Stand about 6 inches (15 cm) away from a wall, with your back facing the wall.
2. Place a soft object, about 6–8 inches (15–20 cm) in diameter, between the back of your head and the wall. A soft object could be a small pillow, a ball, or a folded towel.
3. Gently tilt your head back and press into the soft object. Keep your jaw and forehead relaxed.
4. Hold for _____ seconds.
5. Release the tension slowly. Relax your neck muscles completely before you repeat this exercise.

Repeat _____ times. Complete this exercise _____ times a day.

Posture and body mechanics

Body mechanics refer to the movements and positions of your body while you do your daily activities. Posture is part of body mechanics. Good posture and healthy body mechanics can help to relieve stress in your body's tissues and joints. Good posture

means that your spine is in its natural S-curve position (your spine is *neutral*), your shoulders are pulled back slightly, and your head is not tipped forward.

The following are general guidelines for using improved posture and body mechanics in your everyday activities.

Sitting

- When sitting, keep your spine neutral and keep your feet flat on the floor. Use a footrest, if needed, and keep your thighs parallel to the floor. Avoid rounding your shoulders. Avoid tilting your head forward.
- When working at a desk or a computer, keep your desk at a height where your hands are slightly lower than your elbows. Slide your chair under your desk so you are close enough to maintain good posture.
- When working at a computer, place your monitor at a height where you are looking straight ahead and you do not have to tilt your head forward or downward to look at the screen.

Standing

- When standing, keep your spine neutral and keep your feet about hip-width apart. Keep a slight bend in your knees. Your ears, shoulders, and hips should line up.
- When you do a task in which you stand in one place for a long time, place one foot up on a stable object that is 2–4 inches (5–10 cm) high, such as a footstool. This helps keep your spine neutral.

Resting

When lying down and resting, avoid positions that are most painful for you. Try to support your neck in a neutral position. You can use a contour pillow or a small rolled-up towel. Your pillow should support your neck but not push on it.

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 04/22/2024 Document Reviewed: 07/10/2023
Elsevier Patient Education © 2025 Elsevier Inc.

Pain Medication After Care Information

Pain/Sedating Medicine

After Care Instructions

You have been given pain/sedating medicine during your Emergency Department visit today. This medicine might change the way you think and you might not be able to do some things safely.

FOR THE NEXT 8 HOURS:

- **DO NOT** drive.
- **DO NOT** use machines.
- **DO NOT** use power tools.
- **DO NOT** sign legal papers.
- **DO NOT** take care of children by yourself.
- **DO NOT** climb or go to high places.
- **DO NOT** swim without an adult nearby to help you.
- **DO NOT** drink alcohol.
- **DO NOT** take sleeping pills.
- **DO NOT** take other medicines unless your doctor says it is OK.

GET HELP RIGHT AWAY IF:

- You feel dizzy or pass out (*faint*).
- You feel there are other problems that might be caused by your medicine.
- Your medicine is not helping the pain go away.
- You throw up (*vomit*) or have watery poop (*diarrhea*).
- You have pain in an area that did not hurt before.

MAKE SURE YOU:

- Understand these instructions.
- Will watch your condition.
- Will get help right away if you are not doing well or get worse.

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Released: 06/05/2009 Document Revised: 03/11/2013 Document Reviewed: 12/02/2011
ExitCare® Patient Information ©2016 ExitCare, LLC.

PLEASE NOTE: A Summary of Care document will be provided to your Primary Care Physician to facilitate your transition of care.

Physician Referral Service

THR Well Call Number:

1-877-THR-WELL (1-877-847-9355)

Allergies as of 8/30/2026

Acetaminophen

Egg

Fish

Shellfish

Medications:

Medication changes done in the Emergency Department are not necessarily long term changes. Follow up with your physician for ongoing medications.

Note: You should keep an updated list of your home medications, including vitamins and herbal products, and take this list with you to every visit to your physician, to the Emergency Department and to the hospital.

Radiology Procedures:

If you had an X-Ray, Cat Scan, Sonogram, Ultrasound, or an MRI during your stay, you can review these results with your primary care physician. Other findings, that were not necessarily related to your treatment, may be present.

If you had a test that included IV dye (contrast containing iodine), you will need to drink plenty of water to help your body flush out the dye. Some medications should be stopped for 48 hours after your test. These medications are listed below:

For instructions on restarting any of these medications after your test, call or visit your doctor:

- | | |
|-----------------|----------------|
| • Glucophage XR | • Riomet |
| • Glucophage | • Actoplus Met |
| • Glucovance | • Avandamet |
| • Glumetza | • Fortamet |
| • Metaglip | • Or _____ |
| • Metformin | |

QUITTING SMOKING IS ONE OF THE MOST IMPORTANT THINGS YOU WILL EVER DO:

- Quitting will lower your chance of having a heart attack, stroke, or cancer.
- If you are pregnant, quitting smoking will improve your chances of having a healthy baby.
- By quitting, you protect your children and adolescents from illnesses caused by second-hand smoke.

Resources:

You may want to contact these organizations for further information on smoking and how to quit.

- American Heart Association - 800-AHAUSA1 (242-8721) - www.heart.org
- American Lung Association - 800-LUNGUSA (586-4872) - www.lung.org
- American Cancer Society - 800-227-2345 - www.cancer.org
- National Cancer Institute - 800-4-CANCER (422-6237) - www.cancer.gov

GET HELP FOR ABUSE, NEGLECT, EXPLOITATION, DOMESTIC/FAMILY VIOLENCE

- National Domestic Violence Hotline (Nationwide) - 800-799-SAFE (7233) - www.thehotline.org
- National Human Trafficking Hotline (24-hour hotline) - 888-373-7888 - Humantraffickinghotline.org

- Texas Youth Helpline (24-hour hotline) - 800-989-6884 - www.dfps.texas.gov/youth-helpline/
- Texas Dept of Family & Protective Services (To report child, adult, disabled person abuse/neglect) - 800-252-5400

If you or someone you know is thinking about suicide or is having a mental health crisis, please call or text the Suicide & Crisis Lifeline at 988.

If you or someone you care about is struggling with substance use, support is available.

Addiction Recovery Center by Texas Health – 682-812-6150 – TexasHealthRecovery.com

Substance Abuse and Mental Health Services Administration (SAMHSA) National Helpline - 1-800-662-HELP (4357)

SUTURE REMOVAL REMINDER:

If you had sutures or staples placed today, you have several options for suture or staple removal:

- You can go to your primary care physician or other healthcare provider to have the sutures or staples removed.
- You can contact local clinics to see if they provide suture or staple removal services
- You can return to the Emergency Room to have the sutures or staples removed. **Please be aware you may incur an additional ED visit with additional costs.**

General guidelines to return for suture removal are:

5 days for the face.

7 days for the scalp.

10 days for the extremities and the trunk.

We value Your opinion. Please complete the satisfaction survey you may receive in the mail. Thank You.

HANDOUTS:



What are Emmi Patient Education Videos?

Learning about your health is an important part of your care. Your THR health care team would like you to watch video(s). The video(s) help explain your health condition in an easy-to-understand way.

How Do I Get My Video(s)?

- Log into your MyChart account
- Select Message Center tab
- Select Inbox
- Click on the Emmi link to watch your video(s)

Call 1-866-294-3664 or email support@emmisolutions.com if you have questions about your Emmi patient education video(s).

Remember, Emmi is NOT for urgent medical needs. Call 911 for medical emergencies.

[Excuse Letter](#) (720h ago, onward)

None