

AFTER VISIT SUMMARY

Melody Dashora MRN: 15608293
CEID: SCW-D3VH-7T9W-8ND8

8/28/2026 BAYLOR SCOTT & WHITE MEDICAL
CENTER - GRAPEVINE 817-481-1588

Instructions



Your medications have changed



START taking:

ketorolac (TORADOL)

methocarbamol (ROBAXIN)

traMADol (Ultram)

Review your updated medication list below.



Read the attached information

1. Bruise on a Bone (Periosteal Hematoma): What to Know (English)
2. Stretched or Torn Ligament in the Neck (Cervical Sprain): What to Know (English)
3. Head Injury in Adults: What to Know Easy-to-Read (English)
4. Motor Vehicle Crash Injury in Adults: What It Means Easy-to-Read (English)



Pick up these medications at CVS/pharmacy #10523 - Frisco, TX - 270 Stonebrook Pkwy

ketorolac • methocarbamol • traMADol

Address: 270 Stonebrook Pkwy, Frisco TX 75036
Phone: 214-469-1143



Call PCP in 1 day (around 8/29/2026)



Follow up with BAYLOR SCOTT & WHITE MEDICAL CENTER - GRAPEVINE

Why: As needed

Specialty: Emergency Medicine

Contact: 1650 W College Street
Grapevine Texas 76051
817-481-1588



Follow up with Baylor Scott & White Health - Baylor Scott & White Douglass Community Clinic
Plano

Contact: 801 East Plano Parkway Ste 160
Plano Texas 75074
844-279-3627

Today's Visit

Reason for Visit

Motor Vehicle Crash

Today's Visit (continued)

Diagnoses

- MVC (motor vehicle collision), initial encounter
- Closed head injury, initial encounter
- Neck sprain, initial encounter
- Blunt trauma

Lab Tests Completed

(P) CBC WITH DIFFERENTIAL

Comprehensive Metabolic Panel

Imaging Tests

CT Abdomen Pelvis w Contrast

CT Cervical Spine wo Contrast

CT Chest w Contrast

CT Head wo Contrast

CT Lumbar Spine Reconstructed Images

CT Thoracic Spine Reconstructed Images

XR Knee 1 or 2 Views Left

XR Knee 1 or 2 Views Right

Medications Given

fentaNYL (SUBLIMAZE) Last given at 7:29 AM

iopromide (ULTRAVIST) Last given at 6:25 AM

morphine (PF) Last given at 6:08 AM

ondansetron (ZOFTRAN) Last given at 6:08 AM

sodium chloride Last given at 6:25 AM

sodium chloride Last given at 6:25 AM



Blood Pressure

122/89



Weight

229 lb



Height

5' 6"



Temperature

97.6 °F



Pulse

101



Respiration

18



Oxygen Saturation

96%

What's Next

You currently have no upcoming appointments scheduled.

Thank you for choosing **BAYLOR SCOTT & WHITE MEDICAL CENTER - GRAPEVINE** for your care today. The examination and treatment you have received in the Emergency Department today have been rendered on an emergency basis only and are not intended to be a substitute for an effort to provide complete medical care. You should contact your follow-up physician as it is important that you let him or her check you and report any new or remaining problems since it is impossible to recognize and treat all elements of an injury or illness in a single emergency care center visit.

Need Help?

For questions regarding your discharge instructions, please call your provider.

Your provider's office can help you with the following:

- Help with referrals to a specialist
- Letters for going back to work
- Insurance pre-approved
- Refills of medications (including pain medications and other controlled medications)
- Lab or test results

If you are unable to reach your provider, please call the **Baylor Scott & White Health Patient Advisory Nurse line at 1-800-724-7037**.

You were seen by

1. Travis E Odom, DO
2. Jonathan Donoho Holt, MD

Procedures and tests performed during your visit

(P) CBC WITH DIFFERENTIAL
CT Abdomen Pelvis w Contrast
CT Cervical Spine wo Contrast
CT Chest w Contrast
CT Head wo Contrast
CT Lumbar Spine Reconstructed Images
CT Thoracic Spine Reconstructed Images
Comprehensive Metabolic Panel
XR Knee 1 or 2 Views Left
XR Knee 1 or 2 Views Right

You are allergic to the following

Allergen

Tylenol (Acetaminophen)
Eggshell Membrane

Patient Survey Information

We'd love to hear all about your visit. Press Ganey may send you a survey asking about your experience. Your feedback will help us recognize where your care stood out and pinpoint areas where we can do better for you in the future. Thanks in advance and be well!

Your Medication List



ketorolac 10 mg tablet
Commonly known as: TORADOL

Take 1 tablet (10 mg total) by mouth every 6 (six) hours.



methocarbamol 500 MG tablet
Commonly known as: ROBAXIN

Take 2 tablets (1,000 mg total) by mouth 4 (four) times daily.



traMADol 50 mg tablet
Commonly known as: Ultram

Take 1 tablet (50 mg total) by mouth every 6 (six) hours. Indications: acute pain



Our records indicate that you have an active MyBSWHealth account.

You can view your After Visit Summary by accessing your MyBSWHealth account through the MyBSWHealth mobile app or at my.bswhealth.com and logging in with your username and password. If you don't have a MyBSWHealth username and password but a parent or guardian has access to your record, the parent or guardian should login with their own MyBSWHealth username and password and access your record to view the After Visit Summary.

If you have questions, you can call 855-691-0180 to talk to our MyBSWHealth technical support staff. Remember, MyBSWHealth is NOT to be used for urgent needs. For medical emergencies, dial 911.

MyBSWHealth Mobile App

Manage your health from anywhere with the MyBSWHealth App. There are four options to find the MyBSWHealth App in your phone's app store:

- Text "BETTER" to 88408.
- Scan this code with your smartphone.
- Enter <https://my.bswhealth.com/download> into your phone's internet browser.
- Search for "MyBSWHealth" in your phone's app store.



Discharge Instructions

Bruise on a Bone (Periosteal Hematoma): What to Know

Periosteal hematoma is a bone bruise that can cause tenderness and swelling close to the bone. It can result from a small, hidden break (*fracture*) in the bone, other types of injuries near the bone, or surgery. It usually happens in bones that are close to the surface of the skin. This includes bones of the wrist, shin, knee, or heel. It may take several weeks to heal completely.

What are the causes?

This condition is usually caused by:

- A bone injury. This may result from:
 - A hard, direct hit.
 - A fracture.
 - A sports injury.
- Landing improperly after a jump.
- Running too hard or too much without stretching.
- Twisting injuries.

What are the signs or symptoms?

Symptoms of this condition include:

- Severe pain around the injury. This usually lasts longer than a normal bruise.
- Trouble using the bruised area.
- A tender, swollen area close to the bone.
- A change in color at the injury.

How is this diagnosed?

Periosteal hematoma may be diagnosed based on a physical exam and imaging tests, such as an MRI or X-rays.

How is this treated?

Treatment depends on how severe the injury is. Your health care provider may recommend that you:

- Rest and put ice on the injured area.
- Take over-the-counter or prescription medicine.
- Raise (*elevate*) the injured area to reduce swelling.
- Use crutches to help you walk, if needed.
- Wear a splint, brace, or cast around the injured area.
- Do exercises to improve movement and strength in the area (*physical therapy*).

Follow these instructions at home:

If you have a cast:

- **Do not** put pressure on any part of the cast or bear weight on it until it is fully hardened. This may take several hours. For the first 24 hours, rest the cast on a soft surface, such as a pillow.
- **Do not** stick anything inside the cast to scratch your skin. Doing that increases your risk of infection.
- Check the skin around the cast every day. Tell your provider about any concerns.
- You may put lotion on dry skin around the edges of the cast. **Do not** put lotion on the skin under the cast.
- Keep the cast clean.

If you have a removable splint or brace:



- Wear and remove it only as told.
- Check the skin around it every day. Let your provider know if you see any changes.
- Loosen it if your fingers or toes tingle, become numb, or turn cold and blue.
- If told, you may remove your splint or brace before bathing, before applying ice, and at night.
- Keep the splint or brace clean.

Bathing

If the cast, splint, or brace is not waterproof:

- **Do not** get it wet.
- Cover it with a watertight covering to take a bath or shower.

Managing pain, stiffness, and swelling



- If told, put ice on the injured area.
 - If you have a removable splint or brace, remove it as told by your provider.
 - Put ice in a plastic bag.
 - Place a towel between your skin and the bag or between your cast and the bag.
 - Leave the ice on for 20 minutes, 2–3 times a day.

- If your skin turns bright red, remove the ice right away to prevent skin damage. The risk of damage is higher if you cannot feel pain, heat, or cold.
- Move your fingers or toes often to reduce stiffness and swelling.
- Raise the injured area above the level of your heart while you are sitting or lying down.

Medicines

- Take over-the-counter and prescription medicines only as told.
- Ask your provider if your medicine:
 - Requires you to avoid driving or using machinery.
 - Can cause constipation. You may need to take these actions to prevent or treat constipation:
 - Drink enough fluid to keep your pee (*urine*) pale yellow.
 - Take over-the-counter or prescription medicines.
 - Eat foods that are high in fiber, such as beans, whole grains, and fresh fruits and vegetables.

Activity

- Rest as told by your provider.
- **Do not** use the injured limb to support your body weight until your provider says that you can. Use crutches as told.
- Return to your normal activities as told by your provider. Ask what activities are safe for you. Your provider may recommend low-impact activities, such as swimming or cycling, instead of your normal activities.
- **Do not** take part in any activities that involve the injured area until your provider approves or you are fully healed.
- Do exercises as told.

General instructions

- Ask when it is safe to drive if you have a cast, splint, or brace on your arm or leg.
- Keep all follow-up visits. Your provider will check to see how you are healing.

Contact a health care provider if:

- You have any of these symptoms that get worse:
 - Bruising.
 - Swelling.
 - Tenderness.
 - Warmth in the injured area.
 - Trouble bearing weight on the injured body part.
- Your toes or fingers feel unusually cold, especially if this does not improve after removing a splint or brace.
- You have pain that gets worse or does not improve with medicine.
- You have problems with your cast, such as:
 - Itching under your cast that will not go away.
 - Red or irritated skin around the cast.
 - Cracks or breaks in the cast.
 - A feeling that the cast is too tight.

Get help right away if:

- You have severe pain near the injured area or severe pain when moving.
- You have increased swelling or a loss of feeling in the injured area.

- You have pale, cool skin below the injury in an arm or leg, and this does not go away after removing a splint or brace.
- You cannot move your toes or fingers, even after taking off the splint or brace.

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 10/29/2025 Document Reviewed: 10/31/2023

Elsevier Patient Education © 2026 Elsevier Inc.

Stretched or Torn Ligament in the Neck (Cervical Sprain): What to Know

A cervical sprain is a stretch or tear in one or more of the ligaments in the neck. Ligaments are the tissues that connect bones to each other. Cervical sprains can range from mild to severe. Severe cervical sprains can cause the spinal bones (*vertebrae*) in the neck to be unstable. This can result in spinal cord damage and serious nervous system problems.

Healing time for a cervical sprain depends on the cause and extent of the injury. Most cervical sprains heal in 4–6 weeks.

What are the causes?

Cervical sprains may be caused by trauma, such as an injury from a motor vehicle accident, a fall, or a sudden forward and backward whipping movement of the head and neck (*whiplash injury*). Mild cervical sprains may be caused by wear and tear over time.

What increases the risk?

You are more likely to get a cervical sprain if:

- You take part in activities that have a high risk of trauma to the neck. These include contact sports, gymnastics, and diving.
- You have:
 - Osteoarthritis of the spine.
 - Poor strength and flexibility of the neck.
 - Poor posture.
- You have had a neck injury in the past.
- You spend long periods in positions that put stress on the neck, such as sitting at a computer.

What are the signs or symptoms?

Symptoms of this condition include:

- Any of these problems in the neck, shoulders, or upper back:
 - Pain or tenderness.
 - Stiffness.
 - Swelling.
 - A burning feeling.
- Sudden tightening of neck muscles (*spasms*).
- Limited ability to move the neck.
- Headache.
- Dizziness.
- Nausea or vomiting.
- Weakness, numbness, or tingling in a hand or an arm.

Symptoms may develop right away after injury or may develop over a few days. In some cases, symptoms may go away with treatment and return (*recur*) over time.

How is this diagnosed?

This condition may be diagnosed based on:

- Your symptoms, medical history, and a physical exam.
- Any recent injuries or known neck problems that you have, such as arthritis in the neck.
- Imaging tests, such as X-rays, an MRI, or a CT scan.

How is this treated?



This condition is treated by resting and icing the injured area and doing physical therapy exercises to improve movement and strength. Heat therapy may be used 2–3 days after the injury if there is no swelling. Depending on the severity of your condition, treatment may also include:

- Keeping your neck in place (*immobilized*) for periods of time. This may be done using:
 - A cervical collar. This supports your chin and the back of your head.
 - A cervical traction device. This is a sling that holds up your head. It removes weight and pressure from your neck.
- Medicines for pain or other symptoms.
- Surgery. This is rare.

Follow these instructions at home:

Medicines

- Take over-the-counter and prescription medicines only as told by your health care provider.
- Ask your provider if the medicine prescribed to you:
 - Requires you to avoid driving or using machinery.
 - Can cause constipation. You may need to take these actions to prevent or treat constipation:
 - Drink enough fluid to keep your pee pale yellow.
 - Take over-the-counter or prescription medicines.
 - Eat foods that are high in fiber, such as beans, whole grains, and fresh fruits and vegetables.
 - Limit foods that are high in fat and processed sugars, such as fried or sweet foods.

If you have a cervical collar:

- Wear the collar as told by your provider. **Do not** remove it unless told.
- Ask before making any adjustments to your collar.
- If you have long hair, keep it outside of the collar.

- If you are allowed to remove the collar for cleaning and bathing:
 - Follow instructions about how to remove it safely.
 - Clean it by hand with mild soap and water and air-dry it completely.
 - If your collar has removable pads, remove them every 1–2 days and wash them by hand with soap and water. Let them air-dry completely before putting them back in the collar.
- Tell your provider if your skin under the collar has irritation or sores.

Managing pain, stiffness, and swelling



- Use a cervical traction device as told.
- If told, put ice on the affected area.
 - Put ice in a plastic bag.
 - Place a towel between your skin and the bag.
 - Leave the ice on for 20 minutes, 2–3 times a day.
- If told, apply heat to the affected area before you exercise or as often as told by your provider. Use the heat source that your provider recommends, such as a moist heat pack or a heating pad.
 - Place a towel between your skin and the heat source.
 - Leave the heat on for 20–30 minutes.
- If your skin turns bright red, remove the ice or heat right away to prevent skin damage. The risk of damage is higher if you cannot feel pain, heat, or cold.

Activity

- **Do not** drive while wearing a cervical collar. If you do not have a cervical collar, ask if it is safe to drive while your neck heals.
- **Do not** lift anything that is heavier than 10 lb (4.5 kg) until your provider says that it is safe.
- Rest as told by your provider.
- Avoid positions and activities that make your symptoms worse.
- Do physical therapy exercises as told by your provider or physical therapist.
- Return to your normal activities as told by your provider. Ask your provider what activities are safe for you.

General instructions

- **Do not** use any products that contain nicotine or tobacco. These products include cigarettes, chewing tobacco, and vaping devices, such as e-cigarettes. These can delay healing. If you need help quitting, ask your provider.
- Keep all follow-up visits. Your provider will monitor your injury and activity level.

How is this prevented?

To prevent a cervical sprain from happening again:

- Use and maintain good posture. Make any needed adjustments to your workstation to help you do this.
- Exercise regularly as told by your provider or physical therapist.
- Avoid risky activities that may cause a cervical sprain.

Contact a health care provider if:

- You have symptoms that get worse or do not get better after 2 weeks of treatment.
- You have new symptoms.
- Your pain gets worse or does not get better with medicine.
- You have sores or irritated skin on your neck from wearing your cervical collar.

Get help right away if:


- You have severe pain.
- You develop numbness, tingling, or weakness in any part of your body.
- You cannot move a part of your body (you have *paralysis*).
- You have neck pain along with severe dizziness or headache.

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 10/24/2025 Document Reviewed: 07/21/2023

Elsevier Patient Education © 2026 Elsevier Inc.

Head Injury in Adults: What to Know



There are many types of head injuries. They can be as minor as a small bump. Some head injuries can be worse. Worse injuries include:

- A strong hit to the head that shakes the brain back and forth, causing damage (*concussion*).
- A bruise (*contusion*) of the brain. This means there is bleeding in the brain that can cause swelling.
- A cracked skull (*skull fracture*).
- Bleeding in the brain that gathers, gets thick (makes a *clot*), and forms a bump (*hematoma*).

Most problems from a head injury come in the first 24 hours. However, you may still have side effects up to 7–10 days after your injury. It is important to watch your condition for any changes. You may need to be watched in the emergency department or urgent care, or you may need to stay in the hospital.

What are the causes?

There are many possible causes of a head injury. A serious head injury may be caused by:

- A car accident.
- Bicycle or motorcycle accidents.
- Sports injuries.
- Falls.
- Being hit by an object.

What are the signs or symptoms?

Symptoms of a head injury include a bruise, bump, or bleeding where the injury happened. Other physical symptoms may include:

- Headache.
- Feeling like you may vomit (*nauseous*) or vomiting.
- Dizziness.
- Blurred or double vision.
- Being uncomfortable around bright lights or loud noises.
- Feeling tired.
- Trouble waking up.

- Severe symptoms such as:
 - Feeling weak or numb on one side of the body.
 - Slurred speech.
 - Swallowing problems.
 - Fainting.
 - Shaking movements that you cannot control (*seizures*).

Mental or emotional symptoms may include:

- Feeling grumpy or cranky.
- Confusion and memory problems.
- Having trouble paying attention or concentrating.
- Changes in eating or sleeping habits.
- Feeling worried or nervous (*anxious*).
- Feeling sad (*depressed*).

How is this treated?

Treatment for this condition depends on how bad the injury is and the type of injury you have. The main goal is to prevent problems and to allow the brain time to heal.

Mild head injury

If you have a mild head injury, you may be sent home, and treatment may include:

- Being watched. A responsible adult should stay with you for 24 hours after your injury and check on you often.
- Physical rest.
- Brain rest.
- Pain medicines.

Very bad head injury

If you have a very bad head injury, treatment may include:

- Being watched closely. This includes staying in the hospital.
- Medicines to:
 - Help with pain.
 - Prevent seizures.
 - Help with brain swelling.
- Protecting your airway and using a machine that helps you breathe (*ventilator*).
- Watching for and manage swelling inside the brain.
- Brain surgery. This may be needed to:
 - Remove a collection of blood or blood clots.
 - Stop the bleeding.
 - Remove a part of the skull. This allows room for the brain to swell.

Follow these instructions at home:

Activity

- Rest.

- Avoid activities that are hard or tiring.
- Make sure you get enough sleep.
- Let your brain rest. Do fewer activities that need a lot of thought or attention, such as:
 - Watching TV.
 - Playing memory games and doing puzzles.
 - Job-related work or homework.
 - Working on the computer, social media, and texting.
- Avoid activities that could cause another head injury until your doctor says it is okay. This includes playing sports.
- Ask your doctor when it is safe for you to go back to your normal activities, such as work or school.
- Ask your doctor when you can drive, ride a bicycle, or use machines. **Do not** do these activities if you are dizzy.

Lifestyle



- **Do not** drink alcohol until your doctor says it is okay.
- **Do not** use drugs.
- If it is hard to remember things, write them down.
- If you are easily distracted, try to do one thing at a time.
- Talk with family members or close friends when making important decisions.
- Tell your friends, family, a trusted co-worker, and work manager about your injury, symptoms, and limits (*restrictions*). Have them watch for any problems that are new or getting worse.

General instructions

- Take over-the-counter and prescription medicines only as told by your doctor.
- Have a responsible adult stay with you for 24 hours after your head injury. This person should watch you for any changes in your symptoms and be ready to get help.
- Keep all follow-up visits to catch any new problems early.

How is this prevented?

Having another head injury can be dangerous. Another injury can lead to brain damage, brain swelling, or death. You can avoid this by:

- Working on your balance and strength. This can help you avoid falls.
- Wearing a seat belt when you are in a moving vehicle.
- Wearing a helmet when you:
 - Ride a bicycle.
 - Ski.
 - Do any other sport or activity that has a risk of injury.

- Making your home safer by:
 - Getting rid of clutter from the floors and stairs. This includes things that can make you trip.
 - Using grab bars in bathrooms and handrails by stairs.
 - Placing non-slip mats on floors and in bathtubs.
 - Putting more light in dim areas.

Where to find more information

- Brain Injury Association: biausa.org

Contact a doctor if:

- These symptoms do not go away:
 - Headaches.
 - Dizziness.
 - Double vision or vision changes.
 - Trouble sleeping.
 - Changes in mood.
- You have new symptoms.

Get help right away if:

- You have sudden:
 - Headache that is very bad.
 - Vomiting that does not stop.
 - Changes in the size of one of your pupils. Pupils are the black centers of your eyes.
 - Changes in how you see (*vision*).
 - More confusion or more grumpy moods.
- You have a seizure.
- Your symptoms get worse.
- You have a clear or bloody fluid coming from your nose or ears.

These symptoms may be an emergency. Get help right away. Call 911.

- **Do not wait to see if the symptoms will go away.**
- **Do not drive yourself to the hospital.**

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 10/27/2025 Document Reviewed: 10/05/2023

Elsevier Patient Education © 2026 Elsevier Inc.

Motor Vehicle Crash Injury in Adults: What It Means

After a car accident (*motor vehicle collision*), it is common to have injuries to your head, face, arms, and body. These injuries may include cuts, burns, and bruises. The injuries may also include sore muscles, muscles strains, headaches, and broken bones.

You may feel stiff and sore for the first several hours. You may feel worse after waking up the first morning after the accident. These injuries often feel worse for the first 24–48 hours. After that, you will usually begin to get better with each day. How quickly you get better often depends on:

- How bad the accident was.
- How many injuries you have.
- Where your injuries are.
- What types of injuries you have.
- If you were wearing a seat belt.
- If your airbag was used.

A head injury may result in a concussion. This is a type of brain injury that can have serious effects. If you have a concussion, you should rest as told by your doctor. You must be very careful to avoid having a second concussion.

Follow these instructions at home:

Medicines

- Take over-the-counter and prescription medicines only as told by your doctor.
- If you were prescribed antibiotics, take or apply them as told by your doctor. **Do not** stop using them even if you start to feel better.

Wound care



Follow instructions from your doctor about how to take care of your wound. Make sure you:

- Clean your wound. To do this:
 - Wash it with mild soap and water.
 - Rinse it with water to get all the soap off.
 - Pat it dry with a clean towel. **Do not** rub it.
 - Put an ointment or cream on the wound, if you were told to do so.
- Know when and how to change or remove your bandage (*dressing*).

- Always wash your hands with soap and water for at least 20 seconds before and after you change your bandage. If you cannot use soap and water, use hand sanitizer.
- Leave stitches or skin glue in place for at least 2 weeks.
- Leave tape strips alone unless you are told to take them off. You may trim the edges of the tape strips if they curl up.
- Avoid getting sun on your wound.
- **Do not** disturb the wound. This means:
 - **Do not** scratch or pick at the wound.
 - **Do not** break any blisters you may have.
 - **Do not** peel any skin.

Check your wound every day for signs of infection. Check for:

- More redness, swelling, or pain.
- More fluid or blood.
- Warmth.
- Pus or a bad smell.

Managing pain, stiffness, and swelling



- If told, put ice on the injured areas.
 - Put ice in a plastic bag.
 - Place a towel between your skin and the bag.
 - Leave the ice on for 20 minutes, 2–3 times a day.
 - If your skin turns bright red, take off the ice right away to prevent skin damage. The risk of skin damage is higher if you cannot feel pain, heat, or cold.
- Raise (*elevate*) the wound above the level of your heart while you are sitting or lying down.
 - Sleep with your head raised if the wound is on your face. You may do this by putting an extra pillow under your head.

Activity

- Rest. Rest helps your body to heal. Make sure you:
 - Get plenty of sleep at night. Avoid staying up late.
 - Go to bed at the same time on weekends and weekdays.
- You may have to avoid lifting. Ask your doctor how much you can safely lift.
- Ask your doctor when you can drive, ride a bicycle, or use machinery. **Do not** do these activities if you are dizzy.
- If you are told to wear a brace on an injured arm, leg, or other part of your body, follow instructions from your doctor about activities. Your doctor may give you instructions about driving, bathing, exercising, or working.

General instructions

- If you have a splint, brace, or sling, follow your doctor's instructions on how to use the device.
- Drink enough fluid to keep your pee (*urine*) pale yellow.
- **Do not** drink alcohol.
- Eat healthy foods.

Contact a doctor if:

- You have very bad neck pain, especially pain in the middle of the back of your neck.
- You have loss of feeling (*numbness*), tingling, or weakness in your arms or legs.
- You have a change in your ability to control your pee or poop (*stool*).
- You have swelling in any area of your body, especially your legs.
- You have signs of infection in a wound.
- You have a fever.
- You have blood in your pee, poop, or vomit.
- You have any of the following symptoms for more than 2 weeks after your car accident:
 - Long-term (*chronic*) headaches.
 - Dizziness or balance problems.
 - Feeling like you may vomit.
 - Problems with how you see (*vision*).
 - More sensitivity to noise or light.
 - Sleep problems.
 - Feeling tired all the time.
 - Mental health changes such as:
 - Depression or mood swings.
 - Feeling worried or nervous (*anxiety*).
 - Getting upset or bothered easily.
 - Memory problems.
 - Trouble concentrating or paying attention.

Get help right away if:

- You have shortness of breath.
- You have light-headedness or you faint.
- You have chest pain.
- You have these eye or vision changes:
 - Sudden vision loss or double vision.
 - Your eye suddenly turns red.
 - The black center of your eye (*pupil*) is an odd shape or size.

These symptoms may be an emergency. Get help right away. Call 911.

- **Do not wait to see if the symptoms will go away.**
- **Do not drive yourself to the hospital.**

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Adult Mobility/Fall Prevention Education

How to Prevent Falls in the Home for Adults

Falls can cause injuries. Falls can happen to people of all ages. There're many things you can do to make your home safe and to help prevent falls.

If you need it, ask for help making these changes.

What can I do to prevent falls?

Things you can do in the bathroom



- Keep the floor dry. Clean up any water that's on the floor right away.
- Remove soap buildup. Clean the bathtub or shower to prevent slippery spots.
- Use non-skid mats or decals on the floor of the bathtub or shower.
- Secure bath mats. Use double-sided, non-slip rug tape to keep bath mats in place.
- Use a non-slip stool if you need to sit while showering.
- Install grab bars by the toilet and in the bathtub and shower. **Do not** use towel bars as grab bars.

Adult Mobility/Fall Prevention Education (continued)

Things you can do in the bedroom

- Have a light by your bed. Make sure it is easy to reach.
- **Do not** use any sheets or blankets on your bed that hang to the floor.
- Use a firm chair. Have a sturdy chair with arms that you can use for support when you get dressed.
- Keep a phone by your bed. This way, you can call for help if needed.

Things you can do in the kitchen

- Clean up any spills right away.
- If you need to reach something above you, use a step stool that has a grab bar.
- Keep electrical cords out of the way.
- **Do not** use floor polish or wax that makes floors slippery.

Things you can do with your stairs

- Keep stairs clear. **Do not** leave anything on the stairs.
- Make sure that you have a light switch at the top and the bottom of the stairs. Have them installed if you don't have them.
- Check handrails. Ensure there're handrails on both sides of the stairs. Fix any broken or loose handrails.
- Install non-slip stair treads on all stairs in your home if they don't have carpet.
- **Do not** place throw rugs at the top or bottom of stairs. You can also secure the rugs with carpet tape to prevent them from moving.
- Choose safe carpet. Use carpet that doesn't hide the edge of the steps and is firmly attached.

Things you can do on the outside of your home

- Use bright outdoor lighting.
- Repair walkways and driveways. Fix cracks and remove anything that can trip you, like tools or rocks.
- Mark steps. Use color or contrast paint to mark raised steps or doorways.
- Trim any bushes or trees on paths to your home.
- Check handrails. Make sure handrails are not loose or broken.
- Install guardrails along the edges of any raised decks or porches.
- Remove leaves, snow, and ice regularly. Use sand, salt, or ice melt on walkways during winter months.
- Clean up garage spills. Clean up any grease or oil spills right away.

Other things you can do

- Review your medicines. Talk to your health care provider about your medicines. Some can make you dizzy or confused.
- Wear supportive shoes. Choose closed-toe shoes that fit well and have rubber soles.
- Use a cane, walker, scooter, or crutches to help you move around, if needed.
- Talk to your provider. Ask about exercises that can improve your strength and balance. A physical therapist can help you with this.

General tips

- Use good lighting. Make sure all rooms are well-lit.
 - Replace any light bulbs that burn out.

Adult Mobility/Fall Prevention Education (continued)

- Turn on lights if it's dark.
- Use night lights.
- Keep items that you use often in easy-to-reach places. Lower the shelves around your home if needed.
- Move furniture to create clear paths.
- **Do not** keep throw rugs or other things on the floor. They can make you trip.
- Fix uneven floors.
- Add color or contrast paint or tape to clearly mark and help you see:
 - Grab bars or handrails.
 - First and last steps of staircases.
 - Where the edge of each step is.
- Use ladders safely.
 - Make sure that it's fully opened and locked. **Do not** climb a closed ladder.
 - Have someone hold the ladder.
- Know where your pets are as you move through your home.

Where to find more information

- Centers for Disease Control and Prevention, STEADI: [cdc.gov](https://www.cdc.gov)
- National Council on Aging: [ncoa.org](https://www.ncoa.org)
- National Institute on Aging: [nia.nih.gov](https://www.nia.nih.gov)

Contact a health care provider if:

- You're afraid of falling.
- You feel weak, drowsy, or dizzy.
- You fall.

Get help right away if:

- You faint or have trouble moving after a fall.
- You fall and you hurt your head.

These symptoms may be an emergency. Call 911 right away.

- **Do not wait to see if the symptoms will go away.**
- **Do not drive yourself to the hospital.**

This information is not intended to replace advice given to you by your health care provider. Make sure you discuss any questions you have with your health care provider.

Document Revised: 05/12/2026 Document Reviewed: 01/30/2025

Elsevier Patient Education © 2026 Elsevier Inc.