

# Notes – Linear Algebra

Zhijie Chen

March 8, 2025

## Contents

<i>Linear Maps</i>	3
<i>Products and Quotients of Vector Spaces</i>	3
<i>Duality</i>	3
<i>Polynomials</i>	4
<i>Eigenvalues and Eigenvectors</i>	5
<i>Invariant Subspaces</i>	5
<i>The Minimal Polynomial</i>	5
<i>Commuting Operators</i>	6
<i>Inner Product Spaces</i>	7
<i>Inner Products and Norms</i>	7
<i>Orthonormal Bases</i>	7
<i>Orthogonal Complements and Minimization Problems</i>	7
<i>Operators on Inner Product Spaces</i>	8
<i>Self-Adjoint and Normal Operators</i>	8
<i>Spectral Theorem</i>	9
<i>Positive Operators</i>	9
<i>Singular Value Decomposition</i>	9
<i>Complexification</i>	10

Don't just read it; fight it!  
Ask your own questions.  
Look for your own examples.  
Discover your own proofs.  
Is the hypothesis necessary?  
Is the converse true?  
What happens in the classical special case?  
What about the degenerate cases?  
Where does the proof use the hypothesis?

— Paul Holmos

## Linear Maps

### Products and Quotients of Vector Spaces

**Proposition 1.** *A nonempty subset  $A$  of  $V$  is a translate of some subspace of  $V$  if and only if  $\lambda v + (1 - \lambda)w \in A$  for all  $v, w \in A$  and all  $\lambda \in \mathbf{F}$ .*

**Exercise 2.** Suppose  $U$  is a subspace of  $V$  such that  $V/U$  is finite-dimensional.

- (a) Prove that if  $W$  is a finite-dimensional subspace of  $V$  and  $V = U + W$ , then  $\dim W \geq \dim V/U$ .
- (b) Prove that there exists a finite-dimensional subspace  $W$  of  $V$  such that  $V = U \oplus W$  and  $\dim W = \dim V/U$ .

*Proof.* Let  $\bar{w}_1 + U, \dots, \bar{w}_m + U$  be a basis of  $V/U$ . It can be shown that  $V = \text{span}(\bar{w}_1, \dots, \bar{w}_m) \oplus U$ . Let  $W_0 = \text{span}(\bar{w}_1, \dots, \bar{w}_m)$ . Then  $V = U \oplus W_0$ , as desired.

Now we prove that for each subspace  $W$  of  $V$  such that  $V = U + W$ , we have  $\dim W \geq m = \dim V/U$ . For each  $\bar{w}_k$  above, by definition we have  $\bar{w}_k = u_k + w_k$  for some  $u_k \in U$  and  $w_k \in W$ . It can be shown from the linear independence of  $\bar{w}_1 + U, \dots, \bar{w}_m + U$  that  $\bar{w}_1 - u_1, \dots, \bar{w}_m - u_m$  are independent vectors in  $W$ . Hence  $\dim W \geq m$ .  $\square$

### Duality

**Proposition 3.** *Suppose  $V$  is finite-dimensional and  $U$  is a subspace of  $V$ . Then<sup>1</sup>*

$$U = \left\{ v \in V : \varphi(v) = 0 \text{ for every } \varphi \in U^0 \right\}.$$

<sup>1</sup> Compare this result with  $(U^\perp)^\perp = U$  where  $U$  is a finite-dimensional subspace of inner product space  $V$ .

**Exercise 4.** Suppose  $V$  is finite-dimensional and  $\varphi_1, \dots, \varphi_n$  is a basis of  $V'$ . Prove that there exists a basis of  $V$  whose dual basis is precisely  $\varphi_1, \dots, \varphi_n$ .

*Proof.* We start from an arbitrary basis  $u_1, \dots, u_n$  of  $V$ . Let  $\psi_1, \dots, \psi_n$  be its dual basis. In this proof, we take the standard basis  $e_1, \dots, e_n$  as the basis of  $\mathbf{F}^n$ .

Define  $S, T \in \mathcal{L}(V, \mathbf{F}^n)$  by

$$T(v) = (\varphi_1(v), \dots, \varphi_n(v)), \quad S(v) = (\psi_1(v), \dots, \psi_n(v)).$$

Then  $\mathcal{M}(S, (u_1, \dots, u_n)) = I$ .

Let  $A$  be the change of basis matrix from  $\psi$ 's to  $\varphi$ 's, i.e.,

$$\begin{bmatrix} \varphi_1 & \cdots & \varphi_n \end{bmatrix} = \begin{bmatrix} \psi_1 & \cdots & \psi_n \end{bmatrix} A.$$

Then by the definition of change of basis matrix, we have

$$\mathcal{M}(T, (u_1, \dots, u_n)) = \begin{bmatrix} \mathcal{M}(\varphi_1, (u_1, \dots, u_n)) \\ \vdots \\ \mathcal{M}(\varphi_n, (u_1, \dots, u_n)) \end{bmatrix} = A^t \begin{bmatrix} \mathcal{M}(\psi_1, (u_1, \dots, u_n)) \\ \vdots \\ \mathcal{M}(\psi_n, (u_1, \dots, u_n)) \end{bmatrix}$$

$$= A^t \cdot \mathcal{M}(S, (u_1, \dots, u_n)) = A^t.$$

Consider basis  $v_1, \dots, v_n$  of  $V$  such that the change of basis matrix from  $u$ 's to  $v$ 's is  $(A^t)^{-1}$ .<sup>2</sup> Thus

$$\mathcal{M}(T, (v_1, \dots, v_n)) = \mathcal{M}(T, (u_1, \dots, u_n)) \cdot \mathcal{M}(I, (v_1, \dots, v_n), (u_1, \dots, u_n)) = I.$$

Then the dual basis of  $v_1, \dots, v_n$  is precisely  $\varphi_1, \dots, \varphi_n$ , as desired.  $\square$

**Exercise 5** (A natural isomorphism from primal space onto double dual space). Define  $\Lambda : V \rightarrow V''$  by

$$(\Lambda v)(\varphi) = \varphi(v)$$

for each  $v \in V$  and  $\varphi \in V'$ .

- (a) Prove that if  $T \in \mathcal{L}(V)$ , then  $T'' \circ \Lambda = \Lambda \circ T$ .
- (b) Prove that if  $V$  is finite-dimensional, then  $\Lambda$  is an isomorphism from  $V$  onto  $V''$ .

**Remarks 6.** (a) Suppose  $V$  is finite-dimensional. Then  $V$  and  $V'$  are isomorphic, but finding an isomorphism from  $V$  onto  $V'$  generally requires choosing a basis of  $V$ . In contrast, the isomorphism  $\Lambda$  from  $V$  onto  $V''$  does not require a choice of basis and thus is more natural.

- (b) Another natural isomorphism is  $\pi' \in \mathcal{L}((V/U)', V')$  where  $\pi$  is the normal quotient map.

## Polynomials

**Theorem 7.** Suppose  $p \in \mathcal{P}(\mathbf{F})$  is a nonconstant polynomial of degree  $m$ . Then  $p$  has at most  $m$  zeros in  $\mathbf{F}$ .<sup>3</sup>

**Remark 8.** Theorem 7 implies that the coefficients of a polynomial are uniquely determined. In particular, the degree of a polynomial is well-defined.

**Theorem 9** (Division algorithm for polynomials). Suppose that  $p, s \in \mathcal{P}(\mathbf{F})$ , with  $s \neq 0$ . Then there exist unique polynomials  $q, r \in \mathcal{P}(\mathbf{F})$  such that  $p = sq + r$ .

*Proof.* Let  $n = \deg p$  and  $m = \deg s$ . The case where  $n < m$  is trivial. Thus we now assume that  $n \geq m$ .

The list

$$1, z, \dots, z^{m-1}, s, zs, \dots, z^{n-m}s$$

is linearly independent in  $\mathcal{P}_n(\mathbf{F})$ . And it also has length  $n + 1$ . Hence the list is a basis of  $\mathcal{P}_n(\mathbf{F})$ .

Because  $p \in \mathcal{P}_n(\mathbf{F})$ , there exist unique constants  $a_0, \dots, a_{m-1}, b_0, \dots, b_{n-m} \in \mathbf{F}$  such that

$$\begin{aligned} p &= a_0 + a_1z + \dots + a_{m-1}z^{m-1} + b_0s + b_1zs + \dots + b_{n-m}z^{n-m}s \\ &= (a_0 + a_1z + \dots + a_{m-1}z^{m-1}) + s(b_0 + b_1z + \dots + b_{n-m}z^{n-m}). \end{aligned} \quad \square$$

**Exercise 10.** Suppose  $p, q \in \mathcal{P}(\mathbf{C})$  are nonconstant polynomials with no zeros in common. Let  $m = \deg p$  and  $n = \deg q$ . Prove that there exist  $r \in \mathcal{P}_{n-1}(\mathbf{C})$  and  $s \in \mathcal{P}_{m-1}(\mathbf{C})$  such that  $rp + sq = 1$ .<sup>4</sup>

<sup>2</sup> *Proof Idea.* The change of basis for  $V' \rightarrow V'$  corresponds to the transpose of  $V \leftarrow V$ , where transpose and inverse both come from duality.

<sup>3</sup> A useful corollary of this theorem: when a polynomial  $p$  has too many zeros,  $p = 0$ .

<sup>4</sup> *Proof Idea.* Define  $T : \mathcal{P}_{n-1}(\mathbf{C}) \times \mathcal{P}_{m-1}(\mathbf{C}) \rightarrow \mathcal{P}_{m+n-1}(\mathbf{C})$  by  $T(r, s) = rp + sq$  and prove the surjectivity of  $T$ .

## Eigenvalues and Eigenvectors

### Invariant Subspaces

**Exercise 11.** Suppose that  $\lambda_1, \dots, \lambda_n \in \mathbf{R}$  are pairwise distinct. Prove that the list  $e^{\lambda_1 x}, \dots, e^{\lambda_n x}$  is linearly independent in  $\mathbf{R}^{\mathbf{R}}$ .

*Proof.* Let  $V = \text{span}(e^{\lambda_1 x}, \dots, e^{\lambda_n x})$ .<sup>5</sup> Define  $D \in \mathcal{L}(V)$  by  $Df = f'$ . Then  $e^{\lambda x}$  is an eigenvector of  $D$  corresponding to  $\lambda$ . A list of eigenvectors corresponding to distinct eigenvalues is linearly independent.  $\square$

<sup>5</sup> Alternatively we can let  $V = D(\mathbf{R})$ .

**Exercise 12.** Suppose  $V$  is finite-dimensional,  $T \in \mathcal{L}(V)$ , and  $U$  is a subspace of  $V$  invariant under  $T$ . Prove that each eigenvalue of the quotient operator  $T/U$  is an eigenvalue of  $T$ .

*Proof.* It suffices to show that  $T/U - \lambda I = (T - \lambda I)/U$  is not injective  $\implies T - \lambda I$  is not injective. We prove that  $T - \lambda I$  is invertible  $\implies (T - \lambda I)/U$  is injective.

Suppose  $T - \lambda I$  is invertible.  $U$  being invariant under  $T$  implies that  $U$  is invariant under  $T - \lambda I$ . Thus  $(T - \lambda I)v \in U \iff v \in U$ . Suppose  $((T - \lambda I)/U)(v + U) = 0$ . Then  $(T - \lambda I)v \in U$ , which implies that  $v \in U$ , i.e.,  $v + U = 0$ . That proves the injectivity of  $(T - \lambda I)/U$ .  $\square$

**Exercise 13.** Suppose  $V$  is finite-dimensional and  $T \in \mathcal{L}(V)$ . Prove that  $T$  has an eigenvalue if and only if there exists a subspace of  $V$  of dimension  $\dim V - 1$  that is invariant under  $T$ .<sup>6</sup>

<sup>6</sup> *Proof Idea.* Consider the zero entries in  $\mathcal{M}(T)$  and its transpose matrix  $\mathcal{M}(T')$ .

*Proof.* We first suppose that  $T$  has an eigenvalue  $\lambda$ . Then  $\lambda$  is an eigenvalue of  $T'$ . There exists  $\varphi \in V'$  such that  $\varphi \circ T = T'\varphi = \lambda\varphi$ . Extend  $\varphi$  to a basis  $\varphi, \varphi_2, \dots, \varphi_n$  of  $V'$  and let  $v, v_2, \dots, v_n$  be the basis of  $V$  whose dual basis is  $\varphi, \varphi_2, \dots, \varphi_n$ . Then  $(\varphi \circ T)v_k = \lambda\varphi(v_k) = 0$  for every  $k$ . Because  $\varphi(Tv_k) = 0$  for every  $k$ , we have  $Tv_k \in \text{span}(v_2, \dots, v_n)$ . That proves that  $\text{span}(v_2, \dots, v_n)$  is invariant under  $T$ .

Reversing the steps above leads to an eigenvector of  $T'$ , completing the proof.  $\square$

### The Minimal Polynomial

**Exercise 14** (Companion matrix of a polynomial). Suppose  $a_0, \dots, a_{n-1} \in \mathbf{F}$ . Let  $T \in \mathcal{L}(\mathbf{F}^n)$  be such that  $\mathcal{M}(T)$  (with respect to the standard basis) is

$$\begin{bmatrix} 0 & & & & -a_0 \\ 1 & 0 & & & -a_1 \\ & 1 & 0 & & -a_2 \\ & & \ddots & & \vdots \\ & & & 0 & -a_{n-2} \\ & & & 1 & -a_{n-1} \end{bmatrix}$$

Prove that the minimal polynomial of  $T$  is the polynomial

$$a_0 + a_1 z + \dots + a_{n-1} z^{n-1} + z^n.$$

**Remark 15.** This exercise implies that every monic polynomial is the minimal polynomial of some operator. Hence an algorithm that could produce exact eigenvalues for each operators on each  $\mathbf{F}^n$  does not exist.

**Exercise 16.** Prove that every operator on a finite-dimensional vector space of dimension at least 2 has an invariant subspace of dimension 2.

*Proof.* Let  $T \in \mathcal{L}(V)$  and  $n = \dim V$ . We use induction on  $n$ . The base case  $n = 2$  is trivial. Now suppose  $n > 2$  and the desired result holds for all smaller positive integers. Let  $p$  be the minimal polynomial of  $T$ .

If  $T$  has an eigenvalue  $\lambda$ , then  $p(z) = q(z)(z - \lambda)$  for some monic polynomial  $q$  with  $\deg q = \deg p - 1$ . Because  $q(T)|_{\text{im}(T - \lambda I)} = 0$ , the desired result holds by induction hypothesis if  $\dim \text{im}(T - \lambda I) \geq 2$ . If  $T - \lambda I = 0$  the desired result trivially holds. If  $\dim \text{im}(T - \lambda I) = 1$ , then  $(T - \lambda I)v$  is a scalar multiple of some fixed  $u \in V$  for all  $v \in V$ . Take  $w \in V \setminus \text{span}(u)$  and  $\text{span}(u, w)$  will satisfy the desired property.

If  $T$  has no eigenvalues, then  $\mathbf{F} = \mathbf{R}$  and  $p(z) = q(z)(z^2 + bz + c)$  for some  $b, c \in \mathbf{R}$  with  $b^2 < 4c$  and monic polynomial  $q$  with  $\deg q = \deg p - 2$ . If  $\dim \text{im}(T^2 + bT + cI) \geq 2$  the desired result holds by induction hypothesis. If  $T^2 + bT + cI = 0$ , then let  $w \in V$  be such that  $w \neq 0$ . It can be verified that  $\text{span}(w, Tw)$  is invariant under  $T$ . If  $\dim \text{im}(T^2 + bT + cI) = 1$ , because  $\dim \ker(T^2 + bT + cI)$  is even,  $n$  is odd, which implies that  $T$  has an eigenvalue. That completes the proof.  $\square$

## Commuting Operators

**Exercise 17.** Suppose  $\mathcal{E} \subseteq \mathcal{L}(V)$  and every element of  $\mathcal{E}$  is diagonalizable. Prove that there exists a basis of  $V$  with respect to which every element of  $\mathcal{E}$  has a diagonal matrix if and only if every pair of elements of  $\mathcal{E}$  commutes.<sup>7</sup>

*Proof.* Suppose every pair of elements of  $\mathcal{E}$  commutes. We use induction on  $n = \dim V$ . The base case  $n = 1$  is trivial. Now suppose  $n > 1$  and the desired result holds for all smaller integers. Without loss of generality, suppose  $\mathcal{E} \cap \{\lambda I : \lambda \in \mathbf{F}\} = \emptyset$ , or else consider  $\mathcal{E} \setminus \{\lambda I : \lambda \in \mathbf{F}\}$ .

Let  $\lambda_1, \dots, \lambda_m$  be the distinct eigenvalues of  $T \in \mathcal{E}$ . Then  $V = E(\lambda_1, T) \oplus \dots \oplus E(\lambda_m, T)$ . Because  $E(\lambda_k, T)$  is invariant under every  $S \in \mathcal{E}$ , it suffices to show that the desired result holds on  $E(\lambda_1, T)$ . Because  $E(\lambda_1, T) \subsetneq V$ , it holds by induction hypothesis. The other direction is trivial.  $\square$

**Exercise 18.** Suppose  $V$  is a finite-dimensional nonzero complex vector space. Suppose that  $\mathcal{E} \subseteq \mathcal{L}(V)$  is such that  $S$  and  $T$  commute for all  $S, T \in \mathcal{E}$ .

- Prove that there is a vector in  $V$  that is an eigenvector for every element of  $\mathcal{E}$ .
- Prove that there exists a basis of  $V$  with respect to which every element of  $\mathcal{E}$  has an upper-triangular matrix.<sup>8</sup>

<sup>7</sup> This exercise extends simultaneous diagonalizability to more than 2 (possibly infinitely many) operators.

<sup>8</sup> This exercise extends simultaneous upper triangularizability to more than 2 (possibly infinitely many) operators.

## Inner Product Spaces

### Inner Products and Norms

**Theorems 19** (Polarization identities). (a) Suppose  $V$  is a real inner product space. Then

$$\langle u, v \rangle = \frac{\|u + v\|^2 - \|u - v\|^2}{4}.$$

(b) Suppose  $V$  is a complex inner product space. Then

$$\langle u, v \rangle = \frac{\|u + v\|^2 - \|u - v\|^2 + \|u + iv\|^2 i - \|u - iv\|^2 i}{4}.$$

**Exercise 20.** Prove that if  $\|\cdot\|$  is a norm on  $U$  satisfying the parallelogram equality, then there is an inner product  $\langle \cdot, \cdot \rangle$  on  $U$  such that  $\|u\| = \langle u, u \rangle^{1/2}$  for all  $u \in U$ .<sup>9</sup>

<sup>9</sup> Recall Theorems 19.

### Orthonormal Bases

**Exercise 21.** Suppose  $v_1, \dots, v_m$  is a linearly independent list in  $V$ . Prove that the orthonormal list produced by the formulas of the Gram-Schmidt procedure is the only orthonormal list  $e_1, \dots, e_m$  in  $V$  such that  $\langle v_k, e_k \rangle > 0$  and  $\text{span}(v_1, \dots, v_k) = \text{span}(e_1, \dots, e_k)$  for each  $k$ .

**Exercise 22.** Suppose  $v_1, \dots, v_n$  is a basis of  $V$ . Prove that there exists a basis  $u_1, \dots, u_n$  of  $V$  such that<sup>10</sup>

$$\langle u_j, v_k \rangle = \begin{cases} 0 & \text{if } j \neq k, \\ 1 & \text{if } j = k. \end{cases}$$

<sup>10</sup> *Proof Idea.* Linearity in inner product and patterns of 0, 1 inspire the use of linear functionals.

*Proof.* Define  $\varphi_k \in V'$  for each  $k$  by  $\varphi_k(u) = \langle u, v_k \rangle$ . By the Riesz representation theorem,  $\varphi_1, \dots, \varphi_n$  is a spanning list in  $V'$ . Thus it is a basis of  $V'$ . Let  $u_1, \dots, u_n$  be the basis of  $V$  whose dual basis is  $\varphi_1, \dots, \varphi_n$ . Then  $u_1, \dots, u_n$  satisfies the desired property.  $\square$

### Orthogonal Complements and Minimization Problems

**Theorem 23** (Riesz representation theorem). Suppose  $V$  is finite-dimensional. For each  $v \in V$ , define  $\varphi_v \in V'$  by

$$\varphi_v(u) = \langle u, v \rangle$$

for each  $u \in V$ . Then  $v \mapsto \varphi_v$  is a one-to-one map from  $V$  onto  $V'$ .

*Proof Idea.* If  $\varphi(u) = \langle u, v \rangle$  holds for all  $u \in V$ , then  $v \in (\ker \varphi)^\perp$ . However,  $(\ker \varphi)^\perp$  has dimension 1 (except when  $\varphi = 0$ ). Hence we can obtain the right  $v$  by choosing an arbitrary nonzero  $w \in (\ker \varphi)^\perp$  and then multiplying by an appropriate scalar.  $\square$

**Exercise 24.** Suppose  $V$  is finite-dimensional and  $P \in \mathcal{L}(V)$  is such that  $P^2 = P$ .

- (a) Suppose every vector in  $\ker P$  is orthogonal to every vector in  $\operatorname{im} P$ . Prove that there exists a subspace  $U$  of  $V$  such that  $P = P_U$ .<sup>11</sup>
- (b) Suppose  $\|Pv\| \leq \|v\|$  for every  $v \in V$ . Prove that there exists a subspace  $U$  of  $V$  such that  $P = P_U$ .<sup>12</sup>

<sup>11</sup> *Proof Idea.* Observe that  $\ker P = (\operatorname{im} P)^\perp$ . Thus we prove that  $P$  and  $P_{\operatorname{im} P}$  agree on  $\ker P$  and  $\operatorname{im} P$ .

<sup>12</sup> *Proof Idea.* Observe that among all  $v$ 's with the same  $Pv$ , the one in  $(\ker P)^\perp$  is the shortest. Applying this  $v$  makes the best use of the inequality. Thus we prove that  $P$  and  $P_{(\ker P)^\perp}$  agree on  $\ker P$  and  $(\ker P)^\perp$ .

**Propositions 25** (Algebraic properties of the pseudoinverse). *Suppose  $V$  is finite-dimensional and  $T \in \mathcal{L}(V, W)$ .*

- (a) *The pseudoinverse of an orthogonal projection is the operator itself.*
- (b)  $\ker T^\dagger = (\operatorname{im} T)^\perp$  and  $\operatorname{im} T^\dagger = (\ker T)^\perp$ .
- (c)  $TT^\dagger T = T$  and  $T^\dagger TT^\dagger = T^\dagger$ .
- (d)  $(T^\dagger)^\dagger = T$ .

## Operators on Inner Product Spaces

From now on, we suppose that  $V$  and  $W$  are nonzero finite-dimensional inner product spaces over  $\mathbb{F}$ .

### Self-Adjoint and Normal Operators

**Exercise 26.** Suppose  $T \in \mathcal{L}(V)$  and  $\lambda \in \mathbb{F}$ . Prove that

$$\lambda \text{ is an eigenvalue of } T \iff \overline{\lambda} \text{ is an eigenvalue of } T^*.$$

**Exercise 27.** Suppose  $T \in \mathcal{L}(V)$  and  $U$  is a subspace of  $V$ . Prove that

$$U \text{ is invariant under } T \iff U^\perp \text{ is invariant under } T^*.$$

**Exercise 28.** Suppose  $T \in \mathcal{L}(V)$  is normal. Prove that

$$\ker T^k = \ker T \quad \text{and} \quad \operatorname{im} T^k = \operatorname{im} T.$$

**Exercise 29.** Suppose  $T \in \mathcal{L}(V, W)$ . Prove that under the standard identification of  $V$  with  $V'$  (by Riesz representation theorem) and  $W$  with  $W'$ , the adjoint map  $T^*$  corresponds to the dual map  $T'$ . More precisely, prove that

$$T'(\varphi_w) = \varphi_{T^*w}$$

for all  $w \in W$ .

*Remark 30.* Furthermore, under this identification of  $V$  with  $V'$ , the orthogonal complement corresponds to the annihilator; the formulas for  $\ker T^*$  and  $\operatorname{im} T^*$  become identical to the formulas for  $\ker T'$  and  $\operatorname{im} T'$ . Note that orthogonal complements and adjoints are easier to deal with than annihilators and dual maps.



## Spectral Theorem

**Theorems 31.** (a) Suppose  $T \in \mathcal{L}(V)$ . Suppose  $\mathbf{F} = \mathbf{R}$ . Then  $T$  is self-adjoint if and only if all pairs of eigenvectors corresponding to distinct eigenvalues of  $T$  are orthogonal and  $V = E(\lambda_1, T) \oplus \cdots \oplus E(\lambda_m, T)$ , where  $\lambda_1, \dots, \lambda_m$  denote the distinct eigenvalues of  $T$ .

(b) Suppose  $T \in \mathcal{L}(V)$ . Suppose  $\mathbf{F} = \mathbf{C}$ . Then  $T$  is normal if and only if all pairs of eigenvectors corresponding to distinct eigenvalues of  $T$  are orthogonal and  $V = E(\lambda_1, T) \oplus \cdots \oplus E(\lambda_m, T)$ , where  $\lambda_1, \dots, \lambda_m$  denote the distinct eigenvalues of  $T$ .

**Exercise 32.** (a) Suppose  $\mathbf{F} = \mathbf{C}$  and  $\mathcal{E} \subseteq \mathcal{L}(V)$ . Prove that there is an orthonormal basis of  $V$  with respect to which every element of  $\mathcal{E}$  has a diagonal matrix if and only if  $S$  and  $T$  are commuting normal operators for all  $S, T \in \mathcal{E}$ .

(b) Suppose  $\mathbf{F} = \mathbf{R}$  and  $\mathcal{E} \subseteq \mathcal{L}(V)$ . Prove that there is an orthonormal basis of  $V$  with respect to which every element of  $\mathcal{E}$  has a diagonal matrix if and only if  $S$  and  $T$  are commuting self-adjoint operators for all  $S, T \in \mathcal{E}$ .<sup>13</sup>

<sup>13</sup> This exercise extends the spectral theorems to the context of a collection of commuting operators.

## Positive Operators

**Exercise 33.** For  $T \in \mathcal{L}(V)$  and  $u, v \in V$ , define  $\langle u, v \rangle_T$  by  $\langle u, v \rangle_T = \langle Tu, v \rangle$ .

- (a) Suppose  $T \in \mathcal{L}(V)$ . Prove that  $\langle \cdot, \cdot \rangle_T$  is an inner product on  $V$  if and only if  $T$  is an invertible positive operator (with respect to the original inner product).
- (b) Prove that every inner product on  $V$  is of the form  $\langle \cdot, \cdot \rangle_T$  for some invertible positive operator  $T \in \mathcal{L}(V)$ .

*Proof Idea.* (b) Let  $\langle \cdot, \cdot \rangle_1$  be an inner product on  $V$ . We need to find  $T$  such that  $\langle u, v \rangle_1 = \langle Tu, v \rangle$ . Fixing  $v$  is not sufficient to find the unique  $Tu$ . Thus we fix  $u$  and consider  $\langle v, u \rangle_1 = \langle v, Tu \rangle$ . The map  $v \mapsto \langle v, u \rangle_1$  is in  $V'$ . Hence we use the Riesz representation theorem to define  $Tu$ .  $\square$

## Singular Value Decomposition

**Exercise 34.** Suppose  $T \in \mathcal{L}(V, W)$  and  $s > 0$ . Prove that  $s$  is a singular value of  $T$  if and only if there exist nonzero vectors  $v \in V$  and  $w \in W$  such that<sup>14</sup>

$$Tv = sw \quad \text{and} \quad T^*w = sv.$$

<sup>14</sup> The vectors  $v, w$  satisfying both equations above are called a *Schmidt pair*.

**Exercise 35.** Suppose  $T \in \mathcal{L}(V, W)$ . Suppose  $s_1 \geq s_2 \geq \cdots \geq s_m > 0$  and  $e_1, \dots, e_m$  is an orthonormal list in  $V$  and  $f_1, \dots, f_m$  is an orthonormal list in  $W$  such that

$$Tv = s_1 \langle v, e_1 \rangle f_1 + \cdots + s_m \langle v, e_m \rangle f_m$$

for every  $v \in V$ .

- (a) Prove that  $s_1, \dots, s_m$  are the positive singular values of  $T$ .
- (b) Prove that if  $k \in \{1, \dots, m\}$ , then  $e_k$  is an eigenvector of  $T^*T$  with corresponding eigenvalue  $s_k^2$ .

## Complexification

**Propositions 36** (Properties of complexification). (a)  $\lambda \in \mathbf{R}$  is an eigenvalue of  $T$  if and only if  $\lambda$  is an eigenvalue of  $T_{\mathbf{C}}$ .

(b)  $\lambda \in \mathbf{C}$  is an eigenvalue of  $T_{\mathbf{C}}$  if and only if  $\bar{\lambda}$  is an eigenvalue of  $T_{\mathbf{C}}$ .

(c) The minimal polynomial of  $T_{\mathbf{C}}$  equals the minimal polynomial of  $T$ .

(d) Suppose  $V$  is a real inner product space. For  $u, v, w, x \in V$ , define

$$\langle u + iv, w + ix \rangle_{\mathbf{C}} = \langle u, w \rangle + \langle v, x \rangle + (\langle v, w \rangle - \langle u, x \rangle)i.$$

Then  $\langle \cdot, \cdot \rangle_{\mathbf{C}}$  makes  $V_{\mathbf{C}}$  into a complex inner product space. If  $u, v \in V$ , then

$$\langle u, v \rangle_{\mathbf{C}} = \langle u, v \rangle \quad \text{and} \quad \|u + iv\|_{\mathbf{C}}^2 = \|u\|^2 + \|v\|^2.$$

**Proposition 37.** Every operator on a finite-dimensional nonzero vector space has an invariant subspace of dimension 1 or 2.

*Proof.* The case where  $\mathbf{F} = \mathbf{C}$  is trivial. Now assume  $V$  is a real vector space and  $T \in \mathcal{L}(V)$ . Then  $T_{\mathbf{C}}$ <sup>15</sup> has an eigenvalue  $a + bi$  with  $a, b \in \mathbf{R}$ . Thus there exist  $u, v \in V$ , not both 0, such that

$$Tu + iTv = (au - bv) + (av + bu)i.$$

Hence  $\text{span}(u, v)$  is invariant under  $T$ , as desired.  $\square$

<sup>15</sup> *Proof Idea.* Field extension.