

# Math 121 Notes

Mealoud Mokhtarzad

This set of notes is very informal and tries its best to simplify often hard to digest ideas. Hopefully it's useful in your learning of the material.

**FINISH: put figures in right places**

# Contents

<b>1</b>	<b>Metric space topology</b>	<b>1</b>
1.1	Introduction to metric spaces . . . . .	1
<b>2</b>	<b>Basics of topology</b>	<b>1</b>
2.1	Super basics of topological spaces . . . . .	1
2.2	Subspaces . . . . .	3
2.3	Continuous functions . . . . .	4

# 1 Metric space topology

## 1.1 Introduction to metric spaces

test

# 2 Basics of topology

## 2.1 Super basics of topological spaces

We start with the titular definition.

**Definition 2.1** (Topology). A **topology** on a set  $X$  is a collection  $\mathcal{T}$  of subsets of  $X$  such that

- (1)  $\emptyset, X \in \mathcal{T}$ ;
- (2) any union of sets in  $\mathcal{T}$  is in  $\mathcal{T}$ ; and
- (3) any finite intersection of sets in  $\mathcal{T}$  is in  $\mathcal{T}$ .

The elements of  $\mathcal{T}$  are called **open sets**. A topological space is a pair  $(X, \mathcal{T})$ .

**Remark.** A topology is a prescription for which sets are open.

Let's relate this back to something more familiar and intuitive: open sets in metric spaces. Recall the following properties of open sets in metric spaces: the empty set and the whole space are open; any union of open sets is open; and any finite intersection of open sets is open. I think of our definition of topology as a generalization of that.

Now let's get to some examples of topologies. (Some are gross, which seems to be a common theme in topology. Things can get pretty pathological at times.)

### Example 2.2.

- (1) If  $X$  is a metric space, the collection of open sets is a topology called the **metric topology**.
- (2) If  $X$  is any set, the family  $\mathcal{T} := \{\emptyset, X\}$  forms a topology called the **indiscrete topology**.
- (3) If  $X$  is any set, the family  $\mathcal{T} := \mathcal{P}(X)$  (the collection of all subsets) forms the **discrete topology**.
- (4) Let  $X := \{a, b\}$ . Then  $\mathcal{T} := \{\emptyset, \{a\}, X\}$  is a topology.

Let's start translating some of our basic metric-space-type ideas to more general topological spaces. Next we have the neighborhood (another name for an open ball in metric spaces).

**Definition 2.3** (Neighborhood). Let  $X$  be a topological space. An open **neighborhood** of  $x \in X$  is an open set  $U$  with  $x \in U$ .

Recall that we would define open sets in metric spaces using interior points. We actually have a very similar idea in general topological spaces using open neighborhoods instead of the open ball.

**Definition 2.4.** Let  $X$  be a topological space and suppose  $S$  is a subset of  $X$ . A point  $x \in X$  is an **interior point** of  $S$  if and only if there exists an open neighborhood  $U$  of  $x$  that is contained in  $S$ . The **interior** of  $S$  is the set of interior points of  $S$  and is denoted  $S^\circ$ .

**Proposition 2.5.** *Let  $X$  be a topological space and let  $S$  be a subset of  $X$ . Then*

- (1)  $S^\circ$  is open and
- (2)  $S$  is open if and only if  $S = S^\circ$ .

**Proof.** (1) Suppose  $s \in S^\circ$ . Then there exists an open neighborhood  $U_s$  of  $s$  contained in  $S$ . In fact,  $U_s$  is an open neighborhood of all elements of  $S$  in  $U_s$ . Hence, all points of  $U_s$  are interior points of  $S$  and we have that  $U_s \subseteq S^\circ$ . Now, notice that  $\bigcup_{s \in S^\circ} U_s$  is open by properties of open sets. Moreover, notice that

$$S^\circ \subseteq \bigcup_{s \in S^\circ} U_s \subseteq S^\circ \implies S^\circ = \bigcup_{s \in S^\circ} U_s,$$

so  $S^\circ$  is open.

(2)  $(\implies)$  Suppose  $S$  is open. Then  $S$  is an open neighborhood of any of its points, so  $S \subseteq S^\circ \subseteq S$ . Thus  $S^\circ = S$ .  $(\impliedby)$  If  $S = S^\circ$ , then  $S$  is open since  $S^\circ$  is open. ■

Now let's get to closed sets. (They're what you think they are, if you're familiar with metric topology.)

**Definition 2.6** (Closed set). Let  $X$  be a topological space. A subset  $C$  of  $X$  is **closed** if and only if  $X \setminus C$  is open.

It turns out that our adherent points thing from metric spaces still works but this is the more topology-core definition of closed. (We'll see this more and more in topology as we continue to study it, but things are usually defined in terms of open sets.) Let's actually get into adherent points now.

**Definition 2.7** (Adherent point). Let  $X$  be a topological space. A point  $x \in X$  is **adherent** to a subset  $S$  of  $X$  if and only if every open neighborhood of  $x$  has nonempty intersection with  $S$ .

Now that we have that, we can get to closures.

**Definition 2.8** (Closure of a set). Let  $X$  be a topological space and let  $S$  be a subset of  $X$ . The **closure** of  $S$  is defined to be  $X \setminus (X \setminus S)^\circ$  and is denoted  $\overline{S}$ . Moreover,  $\overline{S}$  consists precisely of the points of  $X$  adherent to  $S$ .

**Proposition 2.9.** *The closure of a set is closed.*

**Proof.** This is clear considering that the closure is in fact the complement of an interior (which is always open). ■

**Proposition 2.10.** *Let  $X$  be a topological space and let  $S$  be a subset of  $X$ . Then,  $S$  is closed if and only if  $S = \overline{S}$ .*

**Proof.**  $(\impliedby)$  Suppose  $S = \overline{S}$ . Then  $S$  is closed (by Proposition 2.9).

$(\implies)$  Suppose  $S$  is closed. Then  $X \setminus S$  is open (by definition), so  $(X \setminus S)^\circ = X \setminus S$  by Proposition 2.5. Thus we have that

$$\overline{S} = X \setminus (X \setminus S)^\circ = X \setminus (X \setminus S) = S.$$

■

Now let's move on to sequences.

**Definition 2.11.** A sequence of points  $(x_n)$  in a topological space  $X$  **converges** to  $x \in X$  if and only if for every open neighborhood  $U$  of  $x$ , there exists an integer  $N$  such that  $x_n \in U$  whenever  $n \geq N$ .

Here's a neat little theorem.

**Theorem 2.12.** *If  $S$  is a subset of a topological space  $X$  and  $(x_n)$  is a sequence in  $S$  converging to  $x \in X$ , then  $x \in \bar{S}$ .*

**Proof.** Suppose towards a contradiction that  $x \notin \bar{S}$ . Then  $x \in X \setminus \bar{S}$ , which is open. Then there exists an open neighborhood of  $x$  entirely contained in  $X \setminus \bar{S}$  and since  $(x_n)$  converges to  $x$ , such a neighborhood contains infinitely many terms of  $(x_n)$ . But  $x_n \in S$  for all  $n$ . ■

Now let's talk a bit more about adherent points. It turns out that we can use adherent points to define what the boundary of a subset of a topological space is.

**Definition 2.13.** Let  $X$  be a topological space and let  $S$  be a subset of  $X$ . A point  $x \in X$  is a **boundary point** of  $S$  if and only if  $x$  is adherent to both  $S$  and  $X \setminus S$  (i.e.,  $x \in \bar{S} \cap \overline{(X \setminus S)}$ ).

The set of boundary points, or **boundary** of  $S$ , is  $\bar{S} \cap \overline{(X \setminus S)}$  and is denoted by  $\partial S$ .

**Proposition 2.14.** *Let  $X$  be a topological space and let  $S$  be a subset of  $X$ . Then,*

- (1)  $S^\circ \cap \partial S = \emptyset$  and
- (2)  $\bar{S} = S^\circ \cup \partial S$

**Proof.** (1) Suppose towards a contradiction that  $S^\circ \cap \partial S$  is nonempty. Then there exists an  $x$  in the intersection. Then there exists an open neighborhood of  $x$  that is completely contained inside  $S$ . However, since  $x \in \partial S$ , every open neighborhood intersects  $X \setminus S$  nontrivially. Thus we have reached a contradiction and  $S^\circ \cap \partial S$  is empty.

(2) ( $\supseteq$ ) Since  $S^\circ \subseteq S \subseteq \bar{S}$  and  $\partial S \subseteq \bar{S}$ , we have that  $S^\circ \cup \partial S \subseteq \bar{S}$ . ( $\subseteq$ ) Suppose  $x \in \bar{S}$ . Then, either there exists an open neighborhood  $U$  of  $x$  with  $U \subseteq S$  or not. In the former case,  $x \in S^\circ$ . In the latter case, any open neighborhood of  $x$  intersects  $X \setminus S$ , so  $x \in \overline{(X \setminus S)}$ , so  $x \in \bar{S} \cap \overline{(X \setminus S)}$ , so  $x \in \partial S$ . ■

Now let's move on to I think our shortest section: subspaces.

## 2.2 Subspaces

**Definition 2.15.** Let  $S$  be a subset of a topological space  $(X, \mathcal{T})$ . The family

$$\mathcal{S} := \{U \cap S : U \in \mathcal{T}\}$$

is a topology for  $S$  called the **subspace topology**. (This is also called the **relative topology**.) The elements of  $\mathcal{S}$  are called **relatively open** subsets of  $S$ .

Here's an example of this:

**Example 2.16.** If  $X$  is a metric space and  $Y$  is a subspace of  $X$ , the topology on  $Y$  obtained by restricting the metric to  $Y$  coincides with the relative topology.

## 2.3 Continuous functions

Because general topological spaces do not have metrics, we can't actually define metrics using our usual  $\epsilon$ - $\delta$  definition, so we use another idea (that's more topology-core tbh): open sets. In particular, we define a function to be continuous if and only if preimages preserve openness. Let's put this in one of our pretty boxes.

**Definition 2.17** (Continuity). Let  $X$  and  $Y$  be topological spaces and let  $f: X \rightarrow Y$  be a function.

The function  $f$  is **continuous** if and only if  $f^{-1}(V)$  is open whenever  $V$  is an open subset of  $Y$  (where  $f^{-1}(V)$  is the preimage of  $V$ ).

The function  $f$  is **continuous at a point**  $x \in X$  if and only if for every open set  $V \subseteq Y$  such that  $f(x) \in V$ , there exists an open set  $U \subseteq X$  such that  $x \in U$  and  $f(U) \subseteq V$ .

It turns out that being continuous at every point in your domain does indeed mean you're continuous. Here's it stated formally and its proof.

**Theorem 2.18.** *Let  $X$  and  $Y$  be topological spaces. A map  $f: X \rightarrow Y$  is continuous if and only if  $f$  is continuous at every point of  $X$ .*

**Proof.** ( $\Rightarrow$ ) Suppose  $f$  is continuous. Taken  $x \in X$  and let  $V$  be an open subset of  $Y$  containing  $f(x)$ . Then,  $U = f^{-1}(V)$  is open in  $X$  (by definition of being continuous), contains  $x$ , and satisfies  $f(U) \subseteq V$ .

( $\Leftarrow$ ) Suppose  $f$  is continuous at every point of  $X$ . Let  $V$  be an open subset of  $Y$  and take  $x \in f^{-1}(V)$ . Then there exists an open set  $U_x \subseteq X$  containing  $x$  with  $U_x \subseteq f^{-1}(V)$ , which means that  $f(U_x) \subseteq V$ .

**FINISH** ■

Honestly this last theorem kind of sucks tbh. I don't see this fact being very useful in my (brief) experience of topology. But now we have something definitely more useful: the composition of continuous functions is continuous.

**Theorem 2.19** (Composition of continuous functions is continuous). *Let  $X$ ,  $Y$ , and  $Z$  be topological spaces,  $f: X \rightarrow Y$ , and  $g: Y \rightarrow Z$  be continuous maps. Then  $g \circ f: X \rightarrow Z$  is continuous.*

**Proof.** Let  $V$  be an open subset of  $Z$ . Then  $g^{-1}(V) \subseteq Y$  is open and  $f^{-1}(g^{-1}(V)) \subseteq X$  is open. Since  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ , we have shown that  $g \circ f$  is continuous. ■

Here's a definition very related to continuity: openness (of a function).

**Definition 2.20** (Open function). Let  $X$  and  $Y$  be topological spaces and let  $f: X \rightarrow Y$  be a function. The function  $f$  is **open** if and only if for all open subsets  $U$  of  $X$ , the image  $f(U)$  is an open subset of  $Y$ .

Now we have a very special type of continuous function: homeomorphisms.

**Definition 2.21** (Homeomorphism). Let  $X$  and  $Y$  be topological spaces. A function  $f: X \rightarrow Y$  is a **homeomorphism** if and only if  $f$  is continuous, bijective, and open.

**Remark.** For a bijection  $f: X \rightarrow Y$  and topological spaces  $X$  and  $Y$ ,  $f$  is open if and only if  $f^{-1}$  is continuous.

This coming idea is what we call it when we have a homeomorphism, but it's not immediately relevant I'd say. This will come up more often as we continue to learn about things called **topological properties**, but that's a story for another day (or rather very soon lmao).

**Definition 2.22** (Homeomorphic spaces). Two topological spaces  $X$  and  $Y$  are **homeomorphic** if and only if there exists a homeomorphism  $f: X \rightarrow Y$ .

**Remark.** If  $f: X \rightarrow Y$  is a homeomorphism, then so is  $f^{-1}: Y \rightarrow X$ . Compositions of homeomorphisms are also homeomorphisms and the identity map is a homeomorphism. Thus, “being homeomorphic” is an equivalence relation on topological spaces.

**Example 2.23.** In  $\mathbb{R}$ , let  $X = [0, r)$  and  $Y = [0, \infty)$ . Then

$$f(t) := \frac{t}{r-t}$$

is a homeomorphism from  $X$  to  $Y$  (exercise lmao).

This can be used to show that in  $\mathbb{R}^n$ , if  $X = B(\mathbf{0}; r)$  and  $Y = \mathbb{R}^n$ , then  $X$  and  $Y$  are homeomorphic via  $g: B(\mathbf{0}; r) \rightarrow \mathbb{R}^n$

$$g(\mathbf{x}) := f(\|\mathbf{x}\|)\mathbf{x}.$$

More generally, any ball  $B(\mathbf{x}_0; r)$  around any  $\mathbf{x}_0 \in \mathbb{R}^n$  is homeomorphic to  $\mathbb{R}^n$  itself via

$$h(\mathbf{x}) := f(\|\mathbf{x} - \mathbf{x}_0\|)(\mathbf{x} - \mathbf{x}_0).$$

**Remark.** A property of a topological space is a **topological invariant** or **topological property** if and only if it is preserved under homeomorphisms. We’ll get some examples of these soon.