

Mathematics

Senior 3 Part II

MELVIN CHIA

Started on 12 June 2023

Finished on XX XX 2023

Actual time spent: XX days

Preface

Why this book?

Disclaimer

Acknowledgements

Contents

26 Applications of Differentiation	4
26.1 Tangent and Normal Lines	4
26.1.1 Practice 1	4
26.1.2 Exercise 26.1	4
26.2 Increasing and Decreasing Functions	6
26.2.1 Practice 2	7
26.2.2 Exercise 26.2	7
26.3 Relative Maximum and Minimum Values of Functions	8
26.3.1 Practice 3	9
26.3.2 Exercise 26.3	9
26.4 Absolute Maximum and Minimum Values of Functions	10
26.4.1 Practice 4	10
26.4.2 Exercise 26.4	11
26.5 The Convexity and the Point of Inflection of Functions	12
26.5.1 Practice 5	13
26.5.2 Practice 26.5	13
26.6 Curve Sketching	14
26.6.1 Practice 6	14
26.6.2 Exercise 26.6	14
26.7 Rate of Change and Related Rate of Change	14
26.7.1 Practice 7	15
26.7.2 Exercise 26.7	15
26.8 Approximate Calculation	16
26.8.1 Practice 8	16
26.8.2 Exercise 26.8	16
26.9 Revision Exercise 26	18
27 Indefinite Integrals	27
27.1 Indefinite Integrals as the Inverse of Differentiation	27
27.2 Arithmetic Properties of Indefinite Integrals	28

27.2.1	Practice 1	28
27.2.2	Exercise 27.2a	29
27.2.3	Practice 2	30
27.2.4	Practice 3	31
27.2.5	Exercise 27.2b	31
27.3	Integration by Substitution	34
27.3.1	Practice 4	35
27.3.2	Exercise 27.3a	35
27.3.3	Practice 5	38
27.3.4	Exercise 27.3b	39
27.4	Integration by Partial Fractions	41
27.4.1	Practice 6	41
27.4.2	Exercise 27.4	42
27.5	Applications of Indefinite Integrals	44
27.5.1	Practice 7	44
27.5.2	Exercise 27.5	44
27.6	Revision Exercise 27	46
27	Definite Integrals	27
27.1	Concept of Definite Integrals and their Relationship with Indefinite Integrals	27
27.1.1	Practice 2	30
27.1.2	Exercise 28.1	30
27.2	Properties and Calculations of Definite Integrals	32
27.2.1	Practice 3	32
27.2.2	Practice 4	34
27.2.3	Exercise 28.2	35
27.3	Area	41
27.3.1	Practice 5	42
27.3.2	Practice 6	42
27.3.3	Exercise 28.3	43
27.3.4	Practice 7	46
27.3.5	Practice 8	46
27.3.6	Exercise 28.4	46
27.4	Revision Exercise 28	47


Chapter 27

Definite Integrals


27.1 Concept of Definite Integrals and their Relationship with Indefinite Integrals

Concept of Definite Integrals

A lot of practical problems, for example finding area and volumes, can be reduced to finding the limit of a certain type of sum. Let's take finding area as an example to explain the method of solving this kind of problem, and hence introduce the concept of definite integrals.




Shown in the diagram above (shaded area) is the area bounded by the line $x = a$, $x = b$, $y = 0$, and the curve $y = f(x)$ where $f(x) \geq 0$. This kind of graph is called the curved trapezoid.



As shown in the diagram above, in order to find the area of this curved trapezoid, we can split it into multiple small curved trapezoid, each of them being substituted by their respective rectangular shape. As such, an approximate value of the area of the curved trapezoid can be acquired by summing up of the area of each rectangle. As the curved trapezoid is being split into smaller and smaller pieces, the approximate value we get will get closer and closer to its actual area.

With this concept in mind, we can split the interval $[a, b]$ into n smaller interval $[x_0, x_1]$, $[x_1, x_2]$, \dots , $[x_{n-1}, x_n]$ where $x_0 = a$, $x_n = b$. From drawing lines that are perpendicular to the x -axis through the points x_1, x_2, \dots, x_{n-1} , we can split the curved trapezoid into n smaller curved trapezoid.



Choose any point ξ_i in the i -th interval $[x_{i-1}, x_i]$, then the area ΔA of the i -th curved trapezoid can be approximated by the area of the rectangle with width $\Delta x_i = x_i - x_{i-1}$ and height $f(\xi_i)$, as shown in the diagram above, i.e.

$$\Delta A_i \approx f(\xi_i) \Delta x$$

And the approximated value of the area of the original curved trapezoid is the sum of the area of all the smaller rectangle, i.e.

$$A = \sum_{i=1}^n \Delta A_i \approx \sum_{i=1}^n f(\xi_i) \Delta x$$

As the number of smaller interval n increases and the width of each interval decreases, the approximated value of the area of the original curved trapezoid gets closer and closer to its actual area. To find the value of A , we split the interval $[a, b]$ into indefinitely many smaller interval such that $\Delta x \rightarrow 0$ (i.e. $n \rightarrow \infty$), hence the area of the original curved trapezoid can be defined as the limit of the sum of the area of all the smaller rectangle, i.e.

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(\xi_i) \Delta x$$

This limit is called the definite integral of $f(x)$ from a to b , and is denoted by the symbol $\int_a^b f(x) dx$, i.e.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(\xi_i) \Delta x$$

where $f(x)$ is called the integrand, $[a, b]$ is called the interval of integration, a and b are called the lower and upper limits of integration respectively.

If $f(x) \geq 0$ in the interval $[a, b]$, we know from the above discussion that the value of the definite integral $\int_a^b f(x) dx$ is the area of the curved trapezoid bounded by the curve $y = f(x)$, the x -axis, and the lines $x = a$ and $x = b$, i.e. $A = \int_a^b f(x) dx$.

If $f(x) \leq 0$ in the interval $[a, b]$, as shown in the diagram above, $f(\xi_i) \Delta x$ is the negative value of the area of the i -th smaller rectangle. Hence, the definite integral $\int_a^b f(x) dx$ is negative, and its absolute value is the area of the curved trapezoid bounded by the curve $y = f(x)$, the x -axis, and the lines $x = a$ and $x = b$, i.e. $A = - \int_a^b f(x) dx$.

The Relationship between Definite Integrals and Indefinite Integrals

The definite integrals and the indefinite integrals has inseparable relationship between them. Consider the case of finding the area of curved trapezoid. Let $x_0 > a$ and $f(x) \geq 0$. The area of the curved trapezoid bounded by the curve $y = f(x)$, the x -axis, and the lines $x = a$, $x = x_0$ and $y = 0$ is $A(x_0) = \int_a^{x_0} f(x)dx$.

When x_0 changes, the area $A(x_0)$ also changes. From the diagram above, we know that

$$m\Delta x \leq A(x_0 + \Delta x) - A(x_0) \leq M\Delta x,$$

where m and M are the minimum and maximum values of $f(x)$ in the interval $[x_0, x_0 + \Delta x]$. Hence,

$$m \leq \frac{A(x_0 + \Delta x) - A(x_0)}{\Delta x} \leq M.$$

Apparently, as Δx approaches 0, both m and M approach $f(x_0)$. Besides, from the definition of derivative, $\lim_{\Delta x \rightarrow 0} \frac{A(x_0 + \Delta x) - A(x_0)}{\Delta x} = A'(x_0)$. Hence, we get $A'(x_0) = f(x_0)$. This relational expression is true for any $x_0 > a$. In other words, $A'(x) = f(x)$, i.e. the derivative of the area function $A(x)$ is the integrand $f(x)$.

Let $\int f(x)dx = F(x) + C$, i.e. $F(x)$ is the primitive of $f(x)$. Then, from $A'(x) = f(x) = F'(x)$, we get $A(x) = F(x) + C$, where C is a constant. When $A(a) = 0$, $x = a$, we get $C = -F(a)$. Hence, $C = -F(a)$, i.e. for any $x_0 > a$, we get $A(x_0) = F(x_0) - F(a)$.

Let $x_0 = b$, we get $A(b) = \int_a^b f(x)dx = F(b) - F(a)$. This relational expression is true for any continuous function $f(x)$. In order to make the relational expression true for any a and b , when $a > b$, the following definition is made.

$$\int_a^b f(x)dx = - \int_b^a f(x)dx.$$

Above all are the relationship between definite integrals and indefinite integrals, i.e.

If $\int f(x)dx = F(x) + C$, then $\int_a^b f(x)dx = F(b) - F(a)$.

This relationship is called the fundamental theorem of calculus. Generally, we express the expression as follows:

$$\int_a^b f(x)dx = [F(x)]_a^b = F(b) - F(a).$$

where $F(x)$ is any primitive of $f(x)$.

When finding the definite integral $\int_a^b f(x)dx$, we only have to find any primitive $F(x)$ of $f(x)$. The other primitive of $f(x)$ can be expressed as $F(x) + C$, where C is a constant. Hence, the definite integral $\int_a^b f(x)dx$ is independent of the choice of primitive of $f(x)$. But,

$$\begin{aligned} & [F(x) + C]_a^b \\ &= [F(b) + C] - [F(a) + C] \\ &= F(b) - F(a) \end{aligned}$$

Hence, the value of the definite integral has nothing to do with the constant C . The value of the definite integral remains the same no matter which primitive of $f(x)$ is chosen.

Note that $[F(x)]_a^b$ can also be written as $F(x) \Big|_a^b$.

27.1.1 Practice 2

1. $\int_2^8 x dx$

Sol.

$$\begin{aligned} I &= \left[\frac{1}{2} x^2 \right]_2^8 \\ &= \frac{1}{2} (64 - 4) \\ &= 30 \end{aligned}$$

2. $\int_{-2}^4 x^3 dx$

Sol.

$$\begin{aligned} I &= \left[\frac{1}{4} x^4 \right]_{-2}^4 \\ &= \frac{1}{4} (256 - 16) \\ &= 60 \end{aligned}$$

3. $\int_{-\pi}^{\pi} \cos x dx$

Sol.

$$\begin{aligned} I &= \left[\sin x \right]_{-\pi}^{\pi} \\ &= \sin \pi - \sin(-\pi) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

4. $\int_0^{\frac{\pi}{4}} \sec^2 x dx$

Sol.

$$\begin{aligned} I &= \left[\tan x \right]_0^{\frac{\pi}{4}} \\ &= \tan \frac{\pi}{4} - \tan 0 \\ &= 1 - 0 \\ &= 1 \end{aligned}$$

27.1.2 Exercise 28.1

1. $\int_{-2}^4 x^2 dx$

Sol.

$$\begin{aligned} I &= \left[\frac{1}{3} x^3 \right]_{-2}^4 \\ &= \frac{1}{3} (64 + 8) \\ &= 24 \end{aligned}$$

2. $\int_1^4 \frac{1}{x^2} dx$

Sol.

$$\begin{aligned} I &= \left[-\frac{1}{x} \right]_1^4 \\ &= -\frac{1}{4} + 1 \\ &= \frac{3}{4} \end{aligned}$$

3. $\int_4^9 \sqrt{x} dx$

Sol.

$$\begin{aligned} I &= \left[\frac{2}{3} x^{\frac{3}{2}} \right]_4^9 \\ &= \frac{2}{3} (27 - 8) \\ &= \frac{38}{3} \end{aligned}$$

4. $\int_1^{27} \frac{1}{\sqrt[3]{x^5}} dx$

Sol.

$$\begin{aligned} I &= \left[-\frac{3}{2} x^{-\frac{2}{3}} \right]_1^{27} \\ &= -\frac{3}{2} \left(\frac{1}{9} - 1 \right) \\ &= \frac{4}{3} \end{aligned}$$

$$5. \int_0^{\frac{\pi}{3}} \cos x dx$$

Sol.

$$\begin{aligned} I &= \left[\sin x \right]_0^{\frac{\pi}{3}} \\ &= \sin \frac{\pi}{3} - \sin 0 \\ &= \frac{\sqrt{3}}{2} - 0 \\ &= \frac{\sqrt{3}}{2} \end{aligned}$$

$$7. \int_0^2 e^x dx$$

Sol.

$$\begin{aligned} I &= e^2 - e^0 \\ &= e^2 - 1 \end{aligned}$$

$$9. \int_1^2 \frac{1}{x} dx$$

Sol.

$$\begin{aligned} I &= \left[\ln |x| \right]_1^2 \\ &= \ln 2 - \ln 1 \\ &= \ln 2 \end{aligned}$$

$$6. \int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} \sin x dx$$

Sol.

$$\begin{aligned} I &= \left[-\cos x \right]_{-\frac{\pi}{4}}^{\frac{\pi}{2}} \\ &= -\cos \frac{\pi}{2} - (-\cos(-\frac{\pi}{4})) \\ &= 0 + \frac{\sqrt{2}}{2} \\ &= \frac{\sqrt{2}}{2} \end{aligned}$$

$$8. \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \operatorname{cosec}^2 x dx$$

Sol.

$$\begin{aligned} I &= \left[-\cot x \right]_{\frac{\pi}{4}}^{\frac{\pi}{2}} \\ &= -\cot \frac{\pi}{2} + \cot \frac{\pi}{4} \\ &= 0 + 1 \\ &= 1 \end{aligned}$$

$$10. \int_0^2 \frac{1}{x+1} dx$$

Sol.

$$\begin{aligned} I &= \left[\ln |x+1| \right]_0^2 \\ &= \ln 3 - \ln 1 \\ &= \ln 3 \end{aligned}$$

27.2 Properties and Calculations of Definite Integrals

Properties of Definite Integrals

The definite integrals have the following basic properties:

$$\text{Property 1} \quad \int_a^b k f(x) dx = k \int_a^b f(x) dx$$

$$\text{Property 2} \quad \int_a^b [f(x) \pm g(x)] dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

$$\text{Property 3} \quad \int_c^a f(x) dx + \int_b^c f(x) dx = \int_b^a f(x) dx$$

27.2.1 Practice 3

1. Find the following definite integrals (Question 1 to 4):

$$(a) \int_1^4 \frac{2x^2 + 3x + 2}{x} dx$$

Sol.

$$\begin{aligned} I &= \int_1^4 \left(2x + 3 + \frac{2}{x} \right) dx \\ &= \left[x^2 + 3x + 2 \ln |x| \right]_1^4 \\ &= 16 + 12 + 2 \ln 4 - 1 - 3 - 2 \ln 1 \\ &= 24 + 2 \ln 4 \end{aligned}$$

$$(c) \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} (2 \cos x - 3 \sec^2 x) dx$$

Sol.

$$\begin{aligned} I &= \left[2 \sin x - 3 \tan x \right]_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \\ &= 2 \sin \frac{\pi}{3} - 3 \tan \frac{\pi}{3} - 2 \sin \left(-\frac{\pi}{3} \right) + 3 \tan \left(-\frac{\pi}{3} \right) \\ &= \sqrt{3} - 3\sqrt{3} + \sqrt{3} - 3\sqrt{3} \\ &= -4\sqrt{3} \end{aligned}$$

$$(b) \int_{-1}^1 (3e^{2x} - 5x) dx$$

Sol.

$$\begin{aligned} I &= \int_{-1}^1 (3e^{2x} - 5x) dx \\ &= \left[\frac{3}{2} e^{2x} - \frac{5}{2} x^2 \right]_{-1}^1 \\ &= \frac{3}{2} e^2 - \frac{5}{2} - \frac{3}{2} e^{-2} + \frac{5}{2} \\ &= \frac{3}{2} (e^2 - e^{-2}) \end{aligned}$$

$$(d) \int_{-2}^4 f(x) dx, f(x) = \begin{cases} x^2 - 2, & -2 \leq x < 2 \\ 4 - x, & 2 \leq x \leq 4 \end{cases}$$

Sol.

$$\begin{aligned} I &= \int_{-2}^2 (x^2 - 2) dx + \int_2^4 (4 - x) dx \\ &= \left[\frac{1}{3} x^3 - 2x \right]_{-2}^2 + \left[4x - \frac{1}{2} x^2 \right]_2^4 \\ &= \frac{8}{3} - 4 + \frac{8}{3} - 4 + 16 - 8 - 8 + 2 \\ &= -\frac{2}{3} \end{aligned}$$

2. Given that $\int_{-2}^5 f(x)dx = 2$, $\int_{-2}^3 f(x)dx = -1$, $\int_3^4 g(x)dx = 3$ and $\int_4^5 g(x)dx = 2$, find:

(a) $\int_3^5 f(x)dx$;

Sol.

$$\begin{aligned}\int_3^5 f(x)dx &= \int_{-2}^5 f(x)dx - \int_{-2}^3 f(x)dx \\ &= 2 - (-1) \\ &= 3\end{aligned}$$

(b) $\int_3^5 \left(\frac{1}{3}g(x) + \frac{1}{2}f(x) \right) dx$.

Sol.

$$\begin{aligned}\int_3^5 \left(\frac{1}{3}g(x) + \frac{1}{2}f(x) \right) dx &= \frac{1}{3} \int_3^5 g(x)dx + \frac{1}{2} \int_3^5 f(x)dx \\ &= \frac{1}{3} \left(\int_3^4 g(x)dx + \int_4^5 g(x)dx \right) + \frac{1}{2} \int_3^5 f(x)dx \\ &= \frac{1}{3}(3 + 2) + \frac{1}{2}(3) \\ &= \frac{19}{6}\end{aligned}$$

3. Given that $f(x) = \sqrt{x^2 + 1}$, find $f'(x)$. Hence, find $\int_0^1 \frac{x}{\sqrt{x^2 + 1}} dx$.

Sol.

$$\begin{aligned}f'(x) &= \frac{1}{2}(x^2 + 1)^{-\frac{1}{2}}(2x) \\ &= \frac{x}{\sqrt{x^2 + 1}}\end{aligned}$$

$$\begin{aligned}\int_0^1 \frac{x}{\sqrt{x^2 + 1}} dx &= \left[\sqrt{x^2 + 1} \right]_0^1 \\ &= \sqrt{2} - 1\end{aligned}$$

Integration by Substitution

In the last chapter, we have learned how to solve indefinite integrals using the method of integration by substitution: $\int f(g(x))g'(x)dx = \int f(u)du$ where $u = g(x)$. From the basic theorem of calculus, the method of integration by substitution of definite integrals can be derived:

$$\int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du$$

The method of integration by substitution of definite integrals is similar to that of indefinite integrals. The only difference is that the limits of integration are changed accordingly.

27.2.2 Practice 4

1. $\int_0^3 4e^{2x} dx$

Sol.

Let $u = 2x$, $du = 2dx$.

When $x = 0$, $u = 0$.

When $x = 3$, $u = 6$.

$$\begin{aligned} I &= \int_0^6 2e^u du \\ &= \left[2e^u \right]_0^6 \\ &= 2e^6 - 2 \\ &= 2(e^6 - 1) \end{aligned}$$

3. $\int_0^4 \frac{x}{25 - x^2} dx$

Sol.

Let $u = 25 - x^2$, $du = -2x dx$.

When $x = 0$, $u = 25$.

When $x = 4$, $u = 9$.

$$\begin{aligned} I &= -\frac{1}{2} \int_{25}^9 \frac{1}{u} du \\ &= -\frac{1}{2} \left[\ln |u| \right]_{25}^9 \\ &= -\frac{1}{2} \ln 9 + \frac{1}{2} \ln 25 \\ &= \frac{1}{2} \ln \frac{25}{9} \\ &= \ln \frac{5}{3} \end{aligned}$$

2. $\int_1^3 \frac{x}{3x^2 + 5} dx$

Sol.

Let $u = 3x^2 + 5$, $du = 6x dx$.

When $x = 1$, $u = 8$.

When $x = 3$, $u = 32$.

$$\begin{aligned} I &= \frac{1}{6} \int_8^{32} \frac{1}{u} du \\ &= \frac{1}{6} \left[\ln |u| \right]_8^{32} \\ &= \frac{1}{6} \ln 32 - \frac{1}{6} \ln 8 \\ &= \frac{1}{6} \ln 4 \\ &= \frac{1}{3} \ln 2 \end{aligned}$$

4. $\int_0^\pi 2 \sin x \cos^2 x dx$

Sol.

Let $u = \cos x$, $du = -\sin x dx$.

When $x = 0$, $u = 1$.

When $x = \pi$, $u = -1$.

$$\begin{aligned} I &= 2 \int_1^{-1} u^2 du \\ &= 2 \left[\frac{1}{3} u^3 \right]_1^{-1} \\ &= \frac{4}{3} \end{aligned}$$

27.2.3 Exercise 28.2

1. $\int_0^4 (x^2 - 2x) dx$

Sol.

$$\begin{aligned} I &= \left[\frac{1}{3}x^3 - x^2 \right]_0^4 \\ &= \frac{64}{3} - 16 \\ &= \frac{16}{3} \end{aligned}$$

3. $\int_{-3}^3 (x+3)^2 dx$

Sol.

Let $u = x + 3$, $du = dx$.

When $x = -3$, $u = 0$.

When $x = 3$, $u = 6$.

$$\begin{aligned} I &= \int_0^6 u^2 du \\ &= \left[\frac{1}{3}u^3 \right]_0^6 \\ &= 72 \end{aligned}$$

5. $\int_0^{\frac{\pi}{2}} (2 \sin 3\theta - 3 \cos 2\theta) d\theta$

Sol.

$$\begin{aligned} I &= \left[-\frac{2}{3} \cos 3\theta - \frac{3}{2} \sin 2\theta \right]_0^{\frac{\pi}{2}} \\ &= \frac{2}{3} + \frac{3}{2} + \frac{2}{3} \\ &= \frac{13}{6} \end{aligned}$$

2. $\int_1^4 \frac{2x^2 - 3\sqrt{x} + 1}{x} dx$

Sol.

$$\begin{aligned} I &= \int_1^4 (2x - 3x^{-\frac{1}{2}} + x^{-1}) dx \\ &= \left[x^2 - 6\sqrt{x} + \ln |x| \right]_1^4 \\ &= 16 - 12 + \ln 4 - 1 + 6 - \ln 1 \\ &= 9 + \ln 4 \end{aligned}$$

4. $\int_{-1}^1 (2+x)(2-x^2) dx$

Sol.

$$\begin{aligned} I &= \int_{-1}^1 (4 - 2x^2 + 2x - x^3) dx \\ &= \left[4x - \frac{2}{3}x^3 + x^2 - \frac{1}{4}x^4 \right]_{-1}^1 \\ &= 4 - \frac{2}{3} + 1 - \frac{1}{4} + 4 - \frac{2}{3} - 1 + \frac{1}{4} \\ &= \frac{20}{3} \end{aligned}$$

6. $\int_0^{\frac{\pi}{3}} \tan \theta d\theta$

Sol.

$$I = \int_0^{\frac{\pi}{3}} \frac{\sin \theta}{\cos \theta} d\theta$$

Let $u = \cos \theta$, $du = -\sin \theta d\theta$.

When $\theta = 0$, $u = 1$.

When $\theta = \frac{\pi}{3}$, $u = \frac{1}{2}$.

$$\begin{aligned} I &= - \int_1^{\frac{1}{2}} \frac{1}{u} du \\ &= - \left[\ln |u| \right]_1^{\frac{1}{2}} \\ &= -\ln \frac{1}{2} + \ln 1 \\ &= \ln 2 \end{aligned}$$

7. $\int_0^1 (e^x - 1)^2 dx$

Sol.

$$\begin{aligned} I &= \int_0^1 (e^{2x} - 2e^x + 1) dx \\ &= \left[\frac{1}{2}e^{2x} - 2e^x + x \right]_0^1 \\ &= \frac{1}{2}e^2 - 2e + 1 - \frac{1}{2} + 2 \\ &= \frac{1}{2}e^2 - 2e + \frac{5}{2} \\ &= \frac{e^2 - 4e + 5}{2} \end{aligned}$$

9. $\int_{-1}^3 \sqrt{2x+3} dx$

Sol.

Let $u = 2x + 3$, $du = 2dx$.

When $x = -1$, $u = 1$.

When $x = 3$, $u = 9$.

$$\begin{aligned} I &= \frac{1}{2} \int_1^9 \sqrt{u} du \\ &= \frac{1}{3} \left[u^{\frac{3}{2}} \right]_1^9 \\ &= \frac{1}{3} (27 - 1) \\ &= \frac{26}{3} \end{aligned}$$

11. $\int_{-1}^1 x^2 (x^3 - 1)^4 dx$

Sol.

Let $u = x^3 - 1$, $du = 3x^2 dx$.

When $x = -1$, $u = -2$.

When $x = 1$, $u = 0$.

$$\begin{aligned} I &= \frac{1}{3} \int_{-2}^0 u^4 du \\ &= \frac{1}{15} \left[u^5 \right]_{-2}^0 \\ &= \frac{32}{15} \end{aligned}$$

8. $\int_1^4 \frac{2}{4x-1} dx$

Sol.

Let $u = 4x - 1$, $du = 4dx$.

When $x = 1$, $u = 3$.

When $x = 4$, $u = 15$.

$$\begin{aligned} I &= \frac{1}{2} \int_3^{15} \frac{1}{u} du \\ &= \frac{1}{2} \left[\ln |u| \right]_3^{15} \\ &= \frac{1}{2} (\ln 15 - \ln 3) \\ &= \frac{1}{2} \ln 5 \end{aligned}$$

10. $\int_1^2 \frac{1}{(2x-1)^3} dx$

Sol.

Let $u = 2x - 1$, $du = 2dx$.

When $x = 1$, $u = 1$.

When $x = 2$, $u = 3$.

$$\begin{aligned} I &= \frac{1}{2} \int_1^3 u^{-3} du \\ &= -\frac{1}{4} \left[u^{-2} \right]_1^3 \\ &= -\frac{1}{4} \left(\frac{1}{9} - 1 \right) \\ &= -\frac{1}{4} \cdot \left(-\frac{8}{9} \right) \\ &= \frac{2}{9} \end{aligned}$$

12. $\int_1^6 x \sqrt{3x-2} dx$

Sol.

Let $u = 3x - 2$, $du = 3dx$, $x = \frac{u+2}{3}$.

When $x = 1$, $u = 1$.

When $x = 6$, $u = 16$.

$$\begin{aligned} I &= \frac{1}{3} \int_1^{16} \frac{u+2}{3} \sqrt{u} du \\ &= \frac{1}{9} \int_1^{16} \left(u^{\frac{3}{2}} + 2u^{\frac{1}{2}} \right) du \\ &= \frac{1}{9} \left[\frac{2}{5} u^{\frac{5}{2}} + \frac{4}{3} u^{\frac{3}{2}} \right]_1^{16} \\ &= \frac{1}{9} \left(\frac{2048}{5} + \frac{256}{3} - \frac{2}{5} - \frac{4}{3} \right) \\ &= \frac{274}{5} \end{aligned}$$

13. $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^2 \theta d\theta$

Sol.

$$\begin{aligned} I &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (1 - \cos 2\theta) d\theta \\ &= \frac{1}{2} \left[\theta - \frac{1}{2} \sin 2\theta \right]_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \\ &= \frac{1}{2} \left(\frac{\pi}{4} - \frac{1}{2} + \frac{\pi}{4} - \frac{1}{2} \right) \\ &= \frac{\pi}{4} - \frac{1}{2} \\ &= \frac{\pi - 2}{4} \end{aligned}$$

14. $\int_3^5 \frac{1}{x^2 - x - 2} dx$

Sol.

$$I = \int_3^5 \frac{1}{(x-2)(x+1)} dx$$

Let $\frac{1}{(x-2)(x+1)} = \frac{A}{x-2} + \frac{B}{x+1}$.

$$Ax + A + Bx - 2B = 1$$

$$(A+B)x + (A-2B) = 1$$

$$\begin{cases} A+B=0 \\ A-2B=1 \end{cases} \Rightarrow \begin{cases} A=\frac{1}{3} \\ B=-\frac{1}{3} \end{cases}$$

$$\begin{aligned} I &= \int_3^5 \left(\frac{1}{3(x-2)} - \frac{1}{3(x+1)} \right) dx \\ &= \frac{1}{3} \int_3^5 \left(\frac{1}{x-2} - \frac{1}{x+1} \right) dx \\ &= \frac{1}{3} \left[\ln|x-2| - \ln|x+1| \right]_3^5 \\ &= \frac{1}{3} (\ln 3 - \ln 6 - \ln 1 + \ln 4) \\ &= \frac{1}{3} \ln 2 \end{aligned}$$

$$15. \int_2^5 \frac{x}{x^3 - x^2 - x + 1} dx$$

Sol.

$$\begin{aligned} I &= \int_2^5 \frac{x}{x^2(x-1) - (x-1)} dx \\ &= \int_2^5 \frac{x}{(x^2-1)(x-1)} dx \\ &= \int_2^5 \frac{x}{(x+1)(x-1)^2} dx \end{aligned}$$

$$\text{Let } \frac{x}{(x+1)(x-1)^2} = \frac{A}{x+1} + \frac{B}{x-1} + \frac{C}{(x-1)^2}.$$

$$\begin{aligned} Ax^2 - 2Ax + A + Bx^2 - B + Cx + C &= x \\ (A+B)x^2 + (-2A+C)x + (A-B+C) &= x \end{aligned}$$

$$\begin{cases} A+B=0 \\ -2A+C=1 \\ A-B+C=0 \end{cases} \Rightarrow \begin{cases} A=-\frac{1}{4} \\ B=\frac{1}{4} \\ C=\frac{1}{2} \end{cases}$$

$$\begin{aligned} I &= \int_2^5 \left(-\frac{1}{4(x+1)} + \frac{1}{4(x-1)} + \frac{1}{2(x-1)^2} \right) dx \\ &= \left[-\frac{1}{4} \ln|x+1| + \frac{1}{4} \ln|x-1| - \frac{1}{2(x-1)} \right]_2^5 \\ &= -\frac{1}{4} \ln 6 + \frac{1}{4} \ln 4 - \frac{1}{8} + \frac{1}{4} \ln 3 - \frac{1}{4} \ln 1 + \frac{1}{2} \\ &= -\frac{1}{4} \ln 6 + \frac{1}{4} \ln 4 - \frac{1}{8} + \frac{1}{4} \ln 3 + \frac{1}{2} \\ &= \frac{1}{4} \ln 2 + \frac{3}{8} \end{aligned}$$

$$17. \int_0^3 \frac{x}{\sqrt{25-x^2}} dx$$

Sol.

$$\text{Let } u = 25 - x^2, du = -2x dx.$$

$$\text{When } x = 0, u = 25.$$

$$\text{When } x = 3, u = 16.$$

$$\begin{aligned} I &= \frac{1}{2} \int_{16}^{25} \frac{1}{\sqrt{u}} du \\ &= \left[\sqrt{u} \right]_{16}^{25} \\ &= 5 - 4 \\ &= 1 \end{aligned}$$

$$16. \int_0^1 \frac{x^2}{x^2 + 2x + 1} dx$$

Sol.

$$I = \int_0^1 \frac{x^2}{(x+1)^2} dx$$

$$\text{Let } u = x + 1, du = dx, x = u - 1.$$

$$\text{When } x = 0, u = 1.$$

$$\text{When } x = 1, u = 2.$$

$$\begin{aligned} I &= \int_1^2 \frac{(u-1)^2}{u^2} du \\ &= \int_1^2 \frac{u^2 - 2u + 1}{u^2} du \\ &= \int_1^2 \left(1 - \frac{2}{u} + \frac{1}{u^2} \right) du \\ &= \left[u - 2 \ln|u| - \frac{1}{u} \right]_1^2 \\ &= 2 - 2 \ln 2 - \frac{1}{2} - 1 + 2 \ln 1 + 1 \\ &= \frac{3}{2} - 2 \ln 2 \end{aligned}$$

$$18. \int_0^{\frac{\pi}{6}} \sin^2 \theta \cos \theta d\theta$$

Sol.

$$\text{Let } u = \sin \theta, du = \cos \theta d\theta.$$

$$\text{When } \theta = 0, u = 0.$$

$$\text{When } \theta = \frac{\pi}{6}, u = \frac{1}{2}.$$

$$\begin{aligned} I &= \int_0^{\frac{1}{2}} u^2 du \\ &= \left[\frac{1}{3} u^3 \right]_0^{\frac{1}{2}} \\ &= \frac{1}{24} \end{aligned}$$

19. Given that $f(x) = \begin{cases} 2x^2 - 1, & -2 \leq x \leq 2 \\ 3x + 1, & 2 < x \leq 4 \end{cases}$, find $\int_{-2}^4 f(x)dx$.

Sol.

$$\begin{aligned} I &= \int_{-2}^2 (2x^2 - 1)dx + \int_2^4 (3x + 1)dx \\ &= \left[\frac{2}{3}x^3 - x \right]_{-2}^2 + \left[\frac{3}{2}x^2 + x \right]_2^4 \\ &= \frac{16}{3} - 2 + \frac{16}{3} - 2 + 24 + 4 - 6 - 2 \\ &= \frac{80}{3} \end{aligned}$$

20. Given that $\int_3^5 f(x)dx = 6$, $\int_5^9 f(x)dx = 18$, $\int_1^4 g(x)dx = 4$ and $\int_3^4 g(x)dx = -4$. Find:

(a) $\int_1^3 g(x)dx$;

Sol.

$$\begin{aligned} \int_1^3 g(x)dx &= \int_1^4 g(x)dx - \int_3^4 g(x)dx \\ &= 4 - (-4) \\ &= 8 \end{aligned}$$

(b) $\int_1^3 f(3x)dx$;

Sol.

Let $u = 3x$, $du = 3dx$.

When $x = 1$, $u = 3$.

When $x = 3$, $u = 9$.

$$\begin{aligned} \int_1^3 f(3x)dx &= \frac{1}{3} \int_3^9 f(u)du \\ &= \frac{1}{3} \int_3^5 f(u)du + \frac{1}{3} \int_5^9 f(u)du \\ &= \frac{1}{3}(6 + 18) \\ &= 8 \end{aligned}$$

(c) $\int_1^3 [f(3x) - 3g(x)]dx$.

Sol.

$$\begin{aligned} \int_1^3 [f(3x) - 3g(x)]dx &= \int_1^3 f(3x)dx - 3 \int_1^3 g(x)dx \\ &= 8 - 3 \cdot 8 \\ &= -16 \end{aligned}$$

21. Given the function $y = x\sqrt{x+1}$,
find $\frac{dy}{dx}$. Hence, find $\int_3^8 \frac{3x+2}{\sqrt{x+1}} dx$.

Sol.

$$\begin{aligned}\frac{dy}{dx} &= \sqrt{x+1} + \frac{x}{2\sqrt{x+1}} \\ &= \frac{2x + x + 2}{2\sqrt{x+1}} \\ &= \frac{3x + 2}{2\sqrt{x+1}}\end{aligned}$$

$$\begin{aligned}\int_3^8 \frac{3x+2}{\sqrt{x+1}} dx &= 2 \int_3^8 \frac{3x+2}{2\sqrt{x+1}} dx \\ &= 2 \int_3^8 \frac{dy}{dx} dx \\ &= 2 \left[x\sqrt{x+1} \right]_3^8 \\ &= 2(24 - 6) \\ &= 36\end{aligned}$$

22. Given the function $y = \frac{x^2 - 1}{2x + 1}$,
find $\frac{dy}{dx}$. Hence, find $\int_0^2 \frac{x^2 + x + 1}{4x^2 + 4x + 1} dx$.

Sol.

$$\begin{aligned}\frac{dy}{dx} &= \frac{(2x+1)(2x) - (x^2-1)(2)}{(2x+1)^2} \\ &= \frac{4x^2 + 2x - 2x^2 + 2}{(2x+1)^2} \\ &= \frac{2(x^2 + x + 1)}{(2x+1)^2}\end{aligned}$$

$$\begin{aligned}\int_0^2 \frac{x^2 + x + 1}{4x^2 + 4x + 1} dx &= \frac{1}{2} \int_0^2 \frac{2(x^2 + x + 1)}{(2x+1)^2} dx \\ &= \frac{1}{2} \int_0^2 \frac{dy}{dx} dx \\ &= \frac{1}{2} \left[\frac{x^2 - 1}{2x + 1} \right]_0^2 \\ &= \frac{1}{2} \left(\frac{3}{5} - \frac{-1}{1} \right) \\ &= \frac{4}{5}\end{aligned}$$

23. Given the function $y = xe^x - e^x$, find $\frac{dy}{dx}$. Hence, find $\int_1^4 2xe^x dx$.

Sol.

$$\begin{aligned}\frac{dy}{dx} &= xe^x + e^x - e^x \\ &= xe^x\end{aligned}$$

$$\begin{aligned}\int_1^4 2xe^x dx &= 2 \int_1^4 xe^x dx \\ &= 2 \int_1^4 \frac{dy}{dx} dx \\ &= 2 \left[xe^x - e^x \right]_1^4 \\ &= 2(4e^4 - e^4 - e + e) \\ &= 6e^4\end{aligned}$$

27.3 Area

When we first introduced the concept of definite integrals, we have studied that, if $f(x) \geq 0$ in the interval $a \leq x \leq b$, then the area of the curved trapezoid bounded by the curve $y = f(x)$, the lines $x = a$ and $x = b$, the x -axis, and the y -axis is given by

$$A = \int_a^b f(x)dx$$

The applications of definite integrals are not limited to finding area of curved trapezoid. In fact, it is widely used in different fields of technologies. To solve real-life problems, the basic mindset is the steps described in the definition: splitting, approximating, finding sum, and taking limit. After understanding the concept, part of the steps can be skipped, and the related definite integrals can be written out straight away.

Let's take finding the area of the curved trapezoid bounded by the line $x = a$, $x = b$, the x -axis, and the curve $y = f(x)$ as an example and do some further elaboration. Since both sides of the curved trapezoid are the straight lines $x = a$ and $x = b$, we can split the interval $a \leq x \leq b$ into multiple smaller intervals, hence the original curved trapezoid is split into multiple smaller curved trapezoid. As shown in the diagram below, take any one of the smaller intervals, and express it as $[x, x + \Delta x]$, its corresponding smaller curved trapezoid can be approximated by a rectangle with width of Δx and height of the function value $f(x)$ at the point x . As such, the area of the smaller curved trapezoid is $\Delta A \approx f(x)\Delta x$.

Summing up the approximated area of all the smaller curved trapezoid, then find the limit of the sum when $\Delta x \rightarrow 0$, we can get the area of the original curved trapezoid, i.e.

$$\sum \Delta A \approx \sum f(x)\Delta x \xrightarrow{\Delta x \rightarrow 0} \int_a^b f(x)dx$$

The same can be applied if $f(x) \leq 0$ in the interval $a \leq x \leq b$, as shown in the diagram below. The area of the curved trapezoid bounded by the curve $y = f(x)$, the lines $x = a$ and $x = b$, the x -axis, and the y -axis is given by

$$A = - \int_a^b f(x)dx$$

If the target area A is bounded by the lines $y = c$, $y = d$, the y -axis, and the curve $x = f(y)$, where $f(y) \geq 0$ in the interval $c \leq y \leq d$, as shown in the diagram below, using the same concept, we can get the area of the region

$$A = \int_c^d f(y)dy$$

27.3.1 Practice 5

1. Find the area of the region bounded by the curve $y = x^2 - 2x + 8$ and the x -axis.

$$(x - 4)(x + 2) = 0$$

$$x = 4 \text{ or } x = -2$$

$$A = - \int_{-2}^4 (x^2 - 2x - 8) dx$$

$$= - \left[\frac{1}{3}x^3 - x^2 - 8x \right]_{-2}^4$$

$$= - \left[\frac{64}{3} - 16 - 32 + \frac{8}{3} + 4 - 16 \right] = -36$$

2. Find the area of the region bounded by the curve $x = \sqrt{y}$, the lines $y = 1$, $y = 4$, and the y -axis.

$$A = \int_1^4 \sqrt{y} dy$$

$$= \left[\frac{2}{3} y^{\frac{3}{2}} \right]_1^4$$

$$= \frac{16}{3} - \frac{2}{3}$$

$$= \frac{14}{3}$$

3. Find the area of the region bounded by the curve $x = y(y + 2)(y - 2)$ and the y -axis.

Sol.

$$A = \int_{-2}^0 y(y + 2)(y - 2) dy - \int_0^2 y(y + 2)(y - 2) dy$$

$$= \int_{-2}^0 (y^3 - 4y) dy - \int_0^2 (y^3 - 4y) dy$$

$$= \left[\frac{1}{4}y^4 - 2y^2 \right]_{-2}^0 - \left[\frac{1}{4}y^4 - 2y^2 \right]_0^2$$

$$= -(4 - 8) - (4 - 8) = 8$$

27.3.2 Practice 6

1. Find the area of the region bounded by the curve $y = x^2 - 4x + 5$ and the line $y = x + 1$.

Sol.

$$x^2 - 5x + 4 = 0$$

$$(x - 4)(x - 1) = 0$$

$$x = 4 \text{ or } x = 1$$

In the interval $1 \leq x \leq 4$, $x + 1 \geq x^2 - 4x + 5$

$$A = \int_1^4 [(x + 1) - (x^2 - 4x + 5)] dx$$

$$= \int_1^4 (-x^2 + 5x - 4) dx$$

$$= \left[-\frac{1}{3}x^3 + \frac{5}{2}x^2 - 4x \right]_1^4$$

$$= -\frac{64}{3} + 40 - 16 + \frac{1}{3} - \frac{5}{2} + 4$$

$$= \frac{9}{2}$$

2. Find the area of the region bounded by the curve $x = 4y - y^2$ and the line $x - 2y + 3 = 0$.

Sol.

$$4y - y^2 - 2y + 3 = 0$$

$$-y^2 + 2y + 3 = 0$$

$$(y - 3)(y + 1) = 0$$

In the interval $-1 \leq y \leq 3$, $4y - y^2 \geq 2y - 3$.

$$A = \int_{-1}^3 [(4y - y^2) - (2y - 3)] dy$$

$$= \int_{-1}^3 (-y^2 + 2y + 3) dy$$

$$= \left[-\frac{1}{3}y^3 + y^2 + 3y \right]_{-1}^3$$

$$= -9 + 9 + 9 - \frac{1}{3} - 1 + 3$$

$$= \frac{32}{3}$$

27.3.3 Exercise 28.3

Find the area of the region bounded by the following curves and lines:

1. $y = 3x^2$, $x = 2$, $x = 5$, and x -axis

Sol.

$$\begin{aligned} A &= \int_2^5 3x^2 dx \\ &= [x^3]_2^5 \\ &= 125 - 8 \\ &= 117 \end{aligned}$$

2. $y = (x - 1)^2$, $x = 4$, x -axis, and y -axis

Sol.

$$\begin{aligned} A &= \int_0^4 (x - 1)^2 dx \\ &= \left[\frac{1}{3}(x - 1)^3 \right]_0^4 \\ &= \frac{1}{3}(3^3 + 1) \\ &= \frac{28}{3} \end{aligned}$$

3. $y = x^2 + 4x - 21$, and x -axis

$$\begin{aligned} x^2 + 4x - 21 &= 0 \\ (x + 7)(x - 3) &= 0 \\ x &= -7 \text{ or } x = 3 \end{aligned}$$

In the interval $-7 \leq x \leq 3$, $x^2 + 4x - 21 \leq 0$.

$$\begin{aligned} A &= \left| \int_{-7}^3 (x^2 + 4x - 21) dx \right| \\ &= \left| \left[\frac{1}{3}x^3 + 2x^2 - 21x \right]_{-7}^3 \right| \\ &= \left| 9 + 18 - 63 - \left(-\frac{343}{3} + 98 + 147 \right) \right| \\ &= \frac{500}{3} \end{aligned}$$

4. $y = e^{2x}$, $x = 0$, $x = 4$, and x -axis

Sol.

$$\begin{aligned} A &= \int_0^4 e^{2x} dx \\ &= \left[\frac{1}{2}e^{2x} \right]_0^4 \\ &= \frac{1}{2}(e^8 - 1) \end{aligned}$$

5. $y = \sin \frac{x}{2}$, $0 \leq x \leq 2\pi$, and x -axis

Sol.

$$\begin{aligned} A &= \int_0^{2\pi} \sin \frac{x}{2} dx \\ &= \left[-2 \cos \frac{x}{2} \right]_0^{2\pi} \\ &= 4 \end{aligned}$$

6. $y = \cos x$, $x = 2\pi$, x -axis, and y -axis

Sol.

$$\begin{aligned} \cos x &= 0 \\ x &= \frac{\pi}{2}, \frac{3\pi}{2} \end{aligned}$$

In the interval $\frac{\pi}{2} \leq x \leq \frac{3\pi}{2}$, $\cos x \leq 0$.

$$\begin{aligned} A &= \int_0^{\frac{\pi}{2}} \cos x dx + \left| \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \cos x dx \right| + \int_{\frac{3\pi}{2}}^{2\pi} \cos x dx \\ &= \left[\sin x \right]_0^{\frac{\pi}{2}} + \left| \left[\sin x \right]_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \right| + \left[\sin x \right]_{\frac{3\pi}{2}}^{2\pi} \\ &= 1 + 1 + 1 + 1 = 4 \end{aligned}$$

7. $x = y^2$, $y = 3$, and y -axis

Sol.

$$\begin{aligned} A &= \int_0^3 y^2 dy \\ &= \left[\frac{1}{3} y^3 \right]_0^3 \\ &= 9 \end{aligned}$$

9. $y = \frac{1}{x}$, $y = \frac{1}{2}$, $y = 2$, and y -axis

Sol.

$$\begin{aligned} A &= \int_{\frac{1}{2}}^2 \frac{1}{x} dx \\ &= \left[\ln x \right]_{\frac{1}{2}}^2 \\ &= \ln 2 - \ln \frac{1}{2} \\ &= 2 \ln 2 \end{aligned}$$

11. Find the area of the region bounded by the curve $y = x^2 - 4$ and the line $y = 3x$.

Sol.

$$\begin{aligned} x^2 - 4 &= 3x \\ x^2 - 3x - 4 &= 0 \\ (x - 4)(x + 1) &= 0 \\ x &= 4 \text{ or } x = -1 \end{aligned}$$

In the interval $-1 \leq x \leq 4$, $x^2 - 4 \leq 3x$.

$$\begin{aligned} A &= \int_{-1}^4 (3x - x^2 + 4) dx \\ &= \left[\frac{3}{2} x^2 - \frac{1}{3} x^3 + 4x \right]_{-1}^4 \\ &= 24 - \frac{64}{3} + 16 - \frac{3}{2} - \frac{1}{3} + 4 \\ &= \frac{125}{6} \end{aligned}$$

8. $x = 9y - y^3$, and y -axis

Sol.

$$\begin{aligned} 9y - y^3 &= 0 \\ y(9 - y^2) &= 0 \\ y &= 0 \text{ or } y = \pm 3 \end{aligned}$$

In the interval $-3 \leq y \leq 0$, $9y - y^3 \leq 0$.

$$\begin{aligned} A &= \left| \int_{-3}^0 (9y - y^3) dy \right| + \int_0^3 (9y - y^3) dy \\ &= \left| \left[\frac{9}{2} y^2 - \frac{1}{4} y^4 \right]_{-3}^0 \right| + \left[\frac{9}{2} y^2 - \frac{1}{4} y^4 \right]_0^3 \\ &= \left| -\frac{81}{2} + \frac{81}{4} \right| + \frac{81}{2} - \frac{81}{4} \\ &= \frac{81}{2} \end{aligned}$$

10. $y^2 = x$, $x = 4$, and $x = 16$

Sol.

$$\begin{aligned} A &= 2 \int_4^{16} \sqrt{x} dx \\ &= 2 \left[\frac{2}{3} x^{\frac{3}{2}} \right]_4^{16} \\ &= 2 \left(\frac{128}{3} - \frac{16}{3} \right) \\ &= \frac{224}{3} \end{aligned}$$

12. Find the area of the region bounded by the curve $y = x^2 + 2x$ and the curve $y = 12 + 4x - x^2$.

$$\begin{aligned} x^2 + 2x &= 12 + 4x - x^2 \\ 2x^2 - 2x - 12 &= 0 \\ (x - 3)(2x + 4) &= 0 \\ x &= 3 \text{ or } x = -2 \end{aligned}$$

In the interval $-2 \leq x \leq 3$, $x^2 + 2x \leq 12 + 4x - x^2$.

$$\begin{aligned} A &= \int_{-2}^3 (12 + 4x - x^2 - x^2 - 2x) dx \\ &= \int_{-2}^3 (12 + 2x - 2x^2) dx \\ &= \left[12x + x^2 - \frac{2}{3} x^3 \right]_{-2}^3 \\ &= 36 + 9 - 18 + 24 - 4 - \frac{16}{3} \\ &= \frac{125}{3} \end{aligned}$$

13. Find the area of the region bounded by the curve $y = \sin x$ and $y = -2 \sin x$ in the interval $0 \leq x \leq \pi$.

Sol.

$$\sin x = -2 \sin x$$

$$\sin x = 0$$

$$x = 0, \pi$$

In the interval $0 \leq x \leq \pi$, $-2 \sin x \leq \sin x$.

$$\begin{aligned} A &= \int_0^{\pi} (3 \sin x) dx \\ &= \left[-3 \cos x \right]_0^{\pi} \\ &= 3 - (-3) = 6 \end{aligned}$$

14. Find the area of the region bounded by the curve $y = e^x$, $y = e^{-2x}$ and the lines $x = -2$ and $x = 4$.

Sol.

$$e^x = e^{-2x}$$

$$x = 0$$

In the interval $-2 \leq x \leq 0$, $e^x \leq e^{-2x}$.

In the interval $0 \leq x \leq 4$, $e^{-2x} \leq e^x$.

$$\begin{aligned} A &= \int_{-2}^0 (e^{-2x} - e^x) dx + \int_0^4 (e^x - e^{-2x}) dx \\ &= \left[-\frac{1}{2} e^{-2x} - e^x \right]_{-2}^0 + \left[e^x + \frac{1}{2} e^{-2x} \right]_0^4 \\ &= -\frac{1}{2} - 1 + \frac{1}{2} e^4 + e^{-2} + e^4 + \frac{1}{2} e^{-8} - 1 - \frac{1}{2} \\ &= \frac{3}{2} e^4 + e^{-2} + \frac{1}{2} e^{-8} - 3 \end{aligned}$$

15. Find the area of the region bounded by the curve $y = x^3 - 10x^2 + 28x$ and the line $y = 4x$.

Sol.

$$x^3 - 10x^2 + 28x = 4x$$

$$x^3 - 10x^2 + 24x = 0$$

$$x(x^2 - 10x + 24) = 0$$

$$x = 0 \text{ or } x = 4 \text{ or } x = 6$$

In the interval $0 \leq x \leq 4$, $x^3 - 10x^2 + 28x \geq 4x$.

In the interval $4 \leq x \leq 6$, $x^3 - 10x^2 + 28x \leq 4x$.

$$\begin{aligned} A &= \int_0^4 (x^3 - 10x^2 + 28x - 4x) dx + \int_4^6 (4x - x^3 + 10x^2 - 28x) dx \\ &= \int_0^4 (x^3 - 10x^2 + 24x) dx + \int_4^6 (-x^3 + 10x^2 - 24x) dx \\ &= \left[\frac{1}{4} x^4 - \frac{10}{3} x^3 + 12x^2 \right]_0^4 + \left[-\frac{1}{4} x^4 + \frac{10}{3} x^3 - 12x^2 \right]_4^6 \\ &= 64 - \frac{640}{3} + 192 - 324 + 720 - 432 + 64 - \frac{640}{3} + 192 \\ &= \frac{148}{3} \end{aligned}$$

16. Find the area of the region bounded by the curve $x = 8y - y^2$, $x = 16y - y^2 - 48$, and y -axis.

Sol.

$$8y - y^2 = 16y - y^2 - 48$$

$$8y = 48$$

$$y = 6$$

17. Find the area of the region bounded by the curve $x = 2y^2 - 8y + 10$ and $x = y^2 - y$.
18. Given that the curve $y = f(x)$ passes through point $(1, 0)$, and the gradient of any point on the curve (x, y) is $3x^2 - 3$. Find the area of the region bounded by the curve, $x = 2$ and x .

27.3.4 Practice 7

- Find the volume of a cylinder with radius r and height h using definite integrals.
- Shown in the diagram below is the shaded region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $a > 0$ and $b > 0$. If the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis is V_x and V_y respectively,
 - find V_x and V_y .
 - if $V_x = 3V_y$, find the value of $a : b$.

27.3.5 Practice 8

- Given that the line $x + y = a$ splits the circle $x^2 + y^2 = a^2$ into two parts, find the volume of the solid of revolution formed by rotating the smaller part of the circle about the x -axis.
- Given that a region is bounded by the curve $y^2 = 8 - x$ and $y^2 = x - 4$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.

27.3.6 Exercise 28.4

Find the volume of the solid of revolution formed by rotating the regions bounded by the following curves and lines about the x -axis (Question 1 to 7):

- $y = \sqrt{x}$, $x = 4$, $x = 9$, and x -axis
- $y = 3x$, $x = 4$, and x -axis
- $y = x(x - 2)$, and x -axis
- $x^2 + y^2 = 4$, $x = 0$, and $x = 2$
- $y = \sin x$, $x = 0$, $x = \pi$, and x -axis
- $y = e^x$, $x = -1$, $x = 1$, and x -axis
- $y = x^3 + x^2 - 2x$, and x -axis

Find the volume of the solid of revolution formed by rotating the regions bounded by the following curves and lines about the y -axis (Question 8 to 14):

- $y = x^3$, $y = 8$, and y -axis
- $x = \sqrt{y - 1}$, $y = 4$, and y -axis

10. $y^2 = x + 3$, $y = 2$, x -axis, and y -axis
11. $y^2 = x + 1$, and y -axis
12. $x^2 - y^2 = 4$, $y = 3$, and x -axis
13. $y = 1 - \sqrt{x}$, x -axis, and y -axis
14. $y = \frac{1}{x} - 1$, $y = 1$, x -axis, and y -axis
15. Given that a region is bounded by the curve $y = 4 - x^2$ and the x -axis. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
16. Given that a region is bounded by the curve $y = 5 - \sqrt{x}$, x -axis, and y -axis. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
17. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y^2 = 9x$, the line $y = 6$, and the y -axis about the y -axis.
18. Given that a region is bounded by the curve $y = x^2 + 1$, the line $x = -2$, $x = 2$, and x -axis. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
19. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y = x(6 - x)$ and the line $y = 3x$ about the x -axis.
20. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y = x^2$, the lines $x = 1$ and $y = 9$ about the y -axis.
21. Given that a region is bounded by the curve $y^2 = 8x$ and the line $y = 2x$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
22. Given that a region is bounded by the curve $y^2 = 8x$ and $y = 8x^2$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
23. Given that a region is bounded by the curve $y^2 = 2x$ and $y^2 = 12 - 4x$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
24. Shown in the diagram below is the shaded region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the line $\frac{x}{a} + \frac{y}{b} = 1$, where $a > 0$ and $b > 0$. If the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis is V_x and V_y respectively,
 - (a) find V_x and V_y .
 - (b) if $V_x = 2V_y$, find the value of $a : b$.

27.4 Revision Exercise 28

1. $\int_0^a (2x^2 - 3x + 2) dx$
2. $\int_1^3 \left(x^2 + \frac{1}{x^3} \right) dx$
3. $\int_{-\frac{\pi}{6}}^{\frac{\pi}{2}} (3 \sin \theta - 2 \cos 2\theta) d\theta$

4. $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (3 \sec^2 \theta + \tan^2 \theta) d\theta$
5. $\int_0^{\ln 2} e^{3x} dx$
6. $\int_1^3 \frac{2}{3x-1} dx$
7. $\int_1^{16} \frac{2x+3}{\sqrt{x}} dx$
8. $\int_1^4 \frac{(\sqrt{x}-1)^2}{x} dx$
9. $\int_1^2 \left(x + \frac{4}{x^2}\right)^2 dx$
10. $\int_0^1 \frac{x+1}{x^2+2x+3} dx$
11. $\int_{-1}^2 \frac{5x}{(1+x^2)^4} dx$
12. $\int_0^2 \frac{x}{\sqrt{25-4x^2}} dx$
13. $\int_2^4 \frac{3x-2}{(2x-3)^2} dx$
14. $\int_2^4 \frac{2}{x^3-x} dx$
15. $\int_1^3 \frac{1}{x^3+2x^2+x} dx$
16. $\int_0^\pi (\sin \theta + \cos \theta)^2 d\theta$
17. $\int_0^{\frac{\pi}{3}} \sec^2 \theta \tan \theta d\theta$
18. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 \theta \cos \theta d\theta$
19. $\int_0^1 \frac{e^x}{e^x+1} dx$
20. $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sec^2 \theta}{\tan \theta} d\theta$
21. Given that $\int_0^4 f(x) dx = 2$, $\int_0^3 g(x) dx = 4$, and $\int_3^8 g(x) dx = 12$. Find the value of $\int_0^8 \left[f\left(\frac{x}{2}\right) - 2g(x) \right] dx$.
22. Given the function $y = (x+3)\sqrt{2x-3}$, find $\frac{dy}{dx}$. Hence, find $\int_2^6 \frac{x}{\sqrt{2x-3}} dx$.
23. Given the function $y = x \ln x$, find $\frac{dy}{dx}$. Hence, find the following definite integrals:

(a) $\int_1^4 \ln x dx$

(b) $\int_1^4 \ln(2x) dx$

24. Find the area of the region bounded by the curve $y = \frac{1}{x+1}$, the lines $x = 1$, $x = 7$, and the x -axis.
25. Find the area of the region bounded by the curve $y = \frac{3}{x}$ and the line $y = 4 - x$.
26. Find the area of the region bounded by the curve $x = y^2 - 5y$ and the line $x + 7y = 24$.
27. Find the area of the region bounded by the curves $y = x^2$ $y^3 = x$.
28. Shown in the diagram below is the shaded region bounded by the curves $y = \ln x$, $y = \ln(2x - 1)$, and the line $y = 3$. Find the area of this region.
29. Find the area of the region bounded by the curves $x = y^3 - y$ $x = y - y^2$.
30. Shown in the diagram below is the shaded region bounded by the curves $y = \sin x$ $y = \sin 2x$ in the interval $0 \leq x \leq \pi$. Find the area of this region.
31. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y = \frac{1}{x+2}$, the line $x = 2$, and two axes about the x -axis.
32. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y = e^x - 3$ and the two axes about the x -axis.
33. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $x = y^2 - 3y$ and the y -axis about the y -axis.
34. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y = x^2$ and the line $y = x + 2$ about the x -axis.
35. Find the volume of the solid of revolution formed by rotating the region bounded by the curve $y^2 = x + 9$ and the line $y = x + 3$ about the y -axis.
36. Given that a region is bounded by the curve $y^2 = 8x$ and $y = x^2$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.
37. SHown in the diagram below is the shaded region bounded by the curve $y = 2 \cos \pi x$, the line $y = 3x$, and the y -axis.
- (a) Prove that the x -coordinate of point A is $\frac{1}{3}$.
- (b) Find the volume of the solid of revolution formed by rotating this region about the x -axis.
38. Given that a region is bounded by the curve $xy = 12$, the line $x = 4$, and $y = 6$. Find the volume of the solid of revolution formed by rotating this region about the x -axis and the y -axis respectively.