

Notes for Calculus III

Melvin Chia

November 2, 2023

Contents

1	Vector in Plane Geometry	2
2	Vector in Space	6
3	Dot Product	8
4	Projections	21
5	Direction Cosines and Direction Angles	23
6	Cross Product	25
7	Distance in Space	31
8	Lines in Space	32
9	Planes in Space	35
10	Cylindrical Coordinates	39
11	Spherical Coordinates	41
12	Vector-valued Functions	43
13	Limits of Vector-valued Functions	48
14	Derivatives and Integration of Vector-valued Functions	49
15	Velocity, Speed and Acceleration	50
16	Tangent Vectors and Normal Vectors	57


Chapter 1

Vector in Plane Geometry

If \vec{v} is a vector whose initial point is $(0, 0)$ and terminal point is (v_1, v_2) , then

$$\vec{v} = \langle v_1, v_2 \rangle$$

is the **component form** of \vec{v} . Here is the graph of the vector \vec{v} in the cartesian plane.




Since the initial point is $(0, 0)$, we say that \vec{v} is in **standard position**.

Note:

The vector with initial point and terminal point $(0, 0)$ is called the **zero vector** and is denoted by $\vec{0}$.

Consider



where P is the initial point and Q is the terminal point of the vector \vec{v} . \vec{v} can be calculated by subtracting the coordinates of the terminal point from the coordinates of the initial point. That is,

$$\vec{v} = \langle q_1 - p_1, q_2 - p_2 \rangle = \langle v_1, v_2 \rangle$$

Length / Norm / Magnitude of a Vector

The **length** or **norm** or **magnitude** of a vector \vec{v} is denoted by $\|\vec{v}\|$ and is given by

$$\|\vec{v}\| = \sqrt{v_1^2 + v_2^2}$$

If $\|\vec{v}\| = 1$, then \vec{v} is called a **unit vector**.

Example 1. Find the length of the vector $\vec{v} = \langle 1, 2 \rangle$.

$$\begin{aligned} \|\vec{v}\| &= \sqrt{v_1^2 + v_2^2} \\ &= \sqrt{1^2 + 2^2} \\ &= \sqrt{5} \end{aligned}$$

Example 2. Calculate the component form of the vector that starts with the point $P(1, 2)$ and ends with the point $Q(5, 4)$.

$$\begin{aligned} \vec{v} &= \langle q_1 - p_1, q_2 - p_2 \rangle \\ &= \langle 5 - 1, 4 - 2 \rangle \\ &= \langle 4, 2 \rangle \end{aligned}$$

By calculating the component form of the vector, we are basically just translating the vector to the origin. Hence, we can conclude that the component form of the vector is the vector that starts from the origin with the same direction and magnitude as the original vector.


If two vectors of different initial and terminal points has the same component form, then the two vectors are said to be **equivalent**.

Unit Vectors

There are two unit vectors that are commonly used in the cartesian plane. They are the **standard unit vectors** \hat{i} and \hat{j} . That is,

$$\hat{i} = \langle 1, 0 \rangle \text{ and } \hat{j} = \langle 0, 1 \rangle$$

Below is a graph of the standard unit vectors.



Given a vector $\vec{v} = \langle v_1, v_2 \rangle$, we can split it into $\langle v_1, 0 \rangle + \langle 0, v_2 \rangle$. Then, factor out the scalars to get $v_1 \langle 1, 0 \rangle + v_2 \langle 0, 1 \rangle$. Hence, we can conclude that


$$\vec{v} = v_1 \hat{i} + v_2 \hat{j}$$

where \vec{v} known as the **linear combination** of \hat{i} and \hat{j} . The scalars v_1 and v_2 are known as the **horizontal component** and **vertical component** of \vec{v} respectively.

Sometimes when we only care about the direction of the vector, we can convert any vector into a unit vector by dividing the vector by its length. That is,

$$\hat{u} = \frac{\vec{u}}{\|\vec{u}\|} \text{ where } \|\hat{u}\| = 1$$

A visual representation of this is shown below.



Example 3. Find the unit vector in the direction of $\vec{v} = \langle 1, 2 \rangle$.

$$\begin{aligned}\hat{v} &= \frac{\vec{v}}{\|\vec{v}\|} \\ &= \frac{\langle 1, 2 \rangle}{\sqrt{5}} \\ &= \left\langle \frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}} \right\rangle\end{aligned}$$

Note:

If you are asked to normalize a vector, you are asked to find the unit vector in the direction of the vector.

Example 4. Find the magnitude of the vector $\vec{v} = 2\hat{i} + 3\hat{j}$.

$$\begin{aligned}\|\vec{v}\| &= \sqrt{v_1^2 + v_2^2} \\ &= \sqrt{2^2 + 3^2} \\ &= \sqrt{13}\end{aligned}$$

Example 5. Find the vector \vec{u} with magnitude 4 and same direction as $\vec{v} = \langle 0, 3 \rangle$.

First, we find the unit vector in the direction of \vec{v} .

$$\begin{aligned}\hat{v} &= \frac{\vec{v}}{\|\vec{v}\|} \\ &= \frac{\langle 0, 3 \rangle}{\sqrt{0^2 + 3^2}} \\ &= \langle 0, 1 \rangle\end{aligned}$$

Then, we multiply the unit vector by the magnitude of the target vector.


$$\begin{aligned}\vec{u} &= 4\hat{v} \\ &= 4\langle 0, 1 \rangle \\ &= \langle 0, 4 \rangle\end{aligned}$$

Chapter 2

Vector in Space

Space Coordinates

The cartesian plane that we are used to is a two-dimensional plane. However, if we extend the dimension further into the third dimension, we get the **three-dimensional space**, which is also known as **Euclidean space**. Below is a graph of the three-dimensional space.



A point in the three-dimensional space is represented by an ordered triple (x, y, z) , where x is the horizontal component, y is the vertical component and z is the depth component.

Example 1. Find the magnitude of the vector $\vec{v} = \langle 3, -2, 1 \rangle$.

$$\begin{aligned}\|\vec{v}\| &= \sqrt{v_1^2 + v_2^2 + v_3^2} \\ &= \sqrt{3^2 + (-2)^2 + 1^2} \\ &= \sqrt{14}\end{aligned}$$

Example 2. Find the magnitude of the vector $\vec{v} = -2\hat{i} + 3\hat{j} + 4\hat{k}$.

$$\begin{aligned}\|\vec{v}\| &= \sqrt{v_1^2 + v_2^2 + v_3^2} \\ &= \sqrt{(-2)^2 + 3^2 + 4^2} \\ &= \sqrt{29}\end{aligned}$$

To find the magnitude of a vector $\vec{v} = \langle v_1, v_2, v_3 \rangle$, we can use the formula

$$\|\vec{v}\| = \sqrt{v_1^2 + v_2^2 + v_3^2}$$

Note:

For vector in any dimension, the magnitude of the vector is given by

$$\|\vec{v}\| = \sqrt{\sum_{i=1}^n v_i^2}$$

where n is the dimension of the vector.

Example 3. Find the vector \vec{u} with magnitude 6 and same direction as $\vec{v} = \langle -6, 4, 0 \rangle$.

$$\begin{aligned}\|\vec{v}\| &= \sqrt{v_1^2 + v_2^2 + v_3^2} \\ &= \sqrt{(-6)^2 + 4^2 + 0^2} \\ &= \sqrt{52} \\ &= 2\sqrt{13}\end{aligned}$$

$$\begin{aligned}\hat{v} &= \frac{\vec{v}}{\|\vec{v}\|} \\ &= \frac{\langle -6, 4, 0 \rangle}{2\sqrt{13}} \\ &= \left\langle -\frac{3}{\sqrt{13}}, \frac{2}{\sqrt{13}}, 0 \right\rangle\end{aligned}$$

$$\begin{aligned}\vec{u} &= 6\hat{v} \\ &= \left\langle -\frac{18}{\sqrt{13}}, \frac{12}{\sqrt{13}}, 0 \right\rangle\end{aligned}$$

Chapter 3

Dot Product

Let $\vec{v} = \langle v_1, v_2, v_3 \rangle$ and $\vec{u} = \langle u_1, u_2, u_3 \rangle$ be two vectors in \mathbb{R}^3 (three-dimensional space), then the dot product of \vec{v} and \vec{u} is given by

$$\vec{v} \cdot \vec{u} = v_1 u_1 + v_2 u_2 + v_3 u_3$$

The final result of the dot product is a number, hence the name **scalar product**. The dot product is also known as the **inner product** of two vectors.

Example 1. Calculate the dot product of the vectors $\vec{v} = \langle 1, 2 \rangle$ and $\vec{u} = \langle -2, 1 \rangle$.

$$\begin{aligned}\vec{v} \cdot \vec{u} &= v_1 u_1 + v_2 u_2 \\ &= 1(-2) + 2(1) \\ &= 0\end{aligned}$$

Note:

If $\vec{v} \cdot \vec{u} = 0$, then \vec{v} and \vec{u} are perpendicular to each other, or they are **orthogonal**. That is,

$$\vec{v} \cdot \vec{u} = 0 \iff \vec{v} \perp \vec{u}$$

Properties of Dot Product

1. $\vec{v} \cdot \vec{u} = \vec{u} \cdot \vec{v}$
2. $\vec{v} \cdot (\vec{u} + \vec{w}) = \vec{v} \cdot \vec{u} + \vec{v} \cdot \vec{w}$
3. $(c\vec{v}) \cdot \vec{u} = c(\vec{v} \cdot \vec{u})$, where c is a scalar
4. $\vec{0} \cdot \vec{v} = 0$, where $\vec{0} = \langle 0, 0 \rangle$
5. $\vec{v} \cdot \vec{v} = \|\vec{v}\|^2$
6. The angle θ between two vectors \vec{v} and \vec{u} is given by

$$\cos \theta = \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} \quad \text{where} \quad 0 \leq \theta \leq \pi$$

From this, we can easily derive that

$$7. \vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta$$

Example 1. Let $\vec{v} = \langle 1, 2 \rangle$, $\vec{u} = \langle 3, 4 \rangle$. Calculate $\vec{u} \cdot (5\vec{v})$.

$$\begin{aligned} \vec{u} \cdot (5\vec{v}) &= 5(\vec{u} \cdot \vec{v}) \\ &= 5(1 \cdot 3 + 2 \cdot 4) \\ &= 55 \end{aligned}$$

Example 2. Prove property 5.

Let $\vec{v} = \langle v_1, v_2 \rangle$, then

$$\begin{aligned} \vec{v} \cdot \vec{v} &= \langle v_1, v_2 \rangle \cdot \langle v_1, v_2 \rangle \\ &= v_1 v_1 + v_2 v_2 \\ &= v_1^2 + v_2^2 \\ &= \left(\sqrt{v_1^2 + v_2^2} \right)^2 \\ &= \|\vec{v}\|^2 \quad \blacksquare \end{aligned}$$

Example 3. Calculate the dot product of the vectors $\vec{v} = \langle 1, 2, 4 \rangle$ and $\vec{u} = \langle -3, -2, 0 \rangle$.

$$\begin{aligned} \vec{v} \cdot \vec{v} &= \langle 1, 2, 4 \rangle \cdot \langle -3, -2, 0 \rangle \\ &= 1(-3) + 2(-2) + 4(0) \\ &= -7 \end{aligned}$$

Example 4. Calculate the dot product of the vectors $\vec{v} = 3i + 2j + 4k$ and $\vec{u} = -i + j$.

$$\begin{aligned}\vec{v} \cdot \vec{u} &= (3i + 2j + 4k) \cdot (-i + j) \\ &= 3(-1) + 2(1) + 4(0) \\ &= -1\end{aligned}$$

Example 5. Determine whether the following vectors are orthogonal.

$$\vec{v} = -2i + 3j + k \quad \vec{u} = 2i + j + k$$

$$\begin{aligned}\vec{v} \cdot \vec{u} &= (-2i + 3j + k) \cdot (2i + j + k) \\ &= -2(2) + 3(1) + 1(1) \\ &= 0\end{aligned}$$

Since $\vec{v} \cdot \vec{u} = 0$, \vec{v} and \vec{u} are orthogonal.

Example 6. Find the angle between the vectors $\vec{v} = \langle 3, 1 \rangle$ and $\vec{u} = \langle 2, -1 \rangle$.

$$\begin{aligned}\cos \theta &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} \\ &= \frac{3(2) + 1(-1)}{\sqrt{3^2 + 1^2} \sqrt{2^2 + (-1)^2}} \\ &= \frac{5}{\sqrt{10} \sqrt{5}} \\ &= \frac{1}{\sqrt{2}} \\ \theta &= \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) \\ &= \frac{\pi}{4}\end{aligned}$$

Example 7. Find the angle between the vectors $\vec{v} = i - 3j + k$ and $\vec{u} = 2i + 4j - k$.


$$\begin{aligned}\cos \theta &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} \\ &= \frac{1(2) + (-3)(4) + 1(-1)}{\sqrt{1^2 + (-3)^2 + 1^2} \sqrt{2^2 + 4^2 + (-1)^2}} \\ &= \frac{-11}{\sqrt{231}} \\ \theta &= 2.38 \text{ rad}\end{aligned}$$

Selected Exercises

Source: *Thomas' Calculus 14th Ed. Exercise 11.3*

Theory and Examples

19. **Sums and differences** In the accompanying figure, it looks as if $\mathbf{v}_1 + \mathbf{v}_2$ and $\mathbf{v}_1 - \mathbf{v}_2$ are orthogonal. Is this mere coincidence, or are there circumstances under which we may expect the sum of two vectors to be orthogonal to their difference? Give reasons for your answer.




Solution.

$$\begin{aligned}
 (v_1 + v_2) \cdot (v_1 - v_2) &= v_1 \cdot v_1 - v_1 \cdot v_2 + v_2 \cdot v_1 - v_2 \cdot v_2 \\
 &= \|v_1\|^2 - \|v_2\|^2 \\
 &= 0 \quad (v_1 = v_2)
 \end{aligned}$$

Therefore, the sum of two vectors is orthogonal to their difference **if and only if the two vectors are in equal length.** ■

20. **Orthogonality on a circle** Suppose that AB is the diameter of a circle with center O and that C is a point on one of the arcs joining A and B . Show that \overrightarrow{CA} and \overrightarrow{CB} are orthogonal.




Solution. Since $\|u\| = \|v\| = \text{radius of the circle}$, $\overrightarrow{CA} = \overrightarrow{CO} + \overrightarrow{OA} = -u - v$ and $\overrightarrow{CB} = \overrightarrow{CO} + \overrightarrow{OB} = u - v$, we have

$$\begin{aligned}\overrightarrow{CA} \cdot \overrightarrow{CB} &= (-u - v) \cdot (u - v) \\ &= -u \cdot u - u \cdot (-v) - v \cdot u - v \cdot (-v) \\ &= -\|u\|^2 + \|v\|^2 \\ &= 0\end{aligned}$$

Therefore, \overrightarrow{CA} and \overrightarrow{CB} are orthogonal. ■

21. **Diagonals of a rhombus** Show that the diagonals of a rhombus (parallelogram with sides of equal length) are perpendicular.

Solution.




Since $d_1 = u + v$ and $d_2 = -v + u$, we have

$$\begin{aligned}d_1 \cdot d_2 &= (u + v) \cdot (-v + u) \\ &= u \cdot u - u \cdot v + v \cdot u - v \cdot v \\ &= \|u\|^2 - \|v\|^2 \\ &= 0\end{aligned}$$

Therefore, the diagonals of a rhombus are perpendicular. ■

24. **Diagonal of parallelogram** Show that the indicated diagonal of the parallelogram determined by vectors \mathbf{u} and \mathbf{v} bisects the angle between \mathbf{u} and \mathbf{v} if $|\mathbf{u}| = |\mathbf{v}|$.



Solution.


$$\begin{aligned}
 (u + v) \cdot u &= u \cdot u + u \cdot v \\
 &= \|u\|^2 + u \cdot v \\
 &= \|v\|^2 + u \cdot v \\
 &= v \cdot v + u \cdot v \\
 &= (u + v) \cdot v
 \end{aligned}$$

Hence, the angle $\arccos\left(\frac{(u + v) \cdot u}{\|u + v\| \|u\|}\right)$ between the diagonal and \vec{u} is equal to the angle $\arccos\left(\frac{(u + v) \cdot v}{\|u + v\| \|v\|}\right)$ between the diagonal and \vec{v} because the inverse cosine function is one-to-one.

Therefore, the indicated diagonal of the parallelogram determined by vectors \mathbf{u} and \mathbf{v} bisects the angle between \mathbf{u} and \mathbf{v} if $|\mathbf{u}| = |\mathbf{v}|$. ■

25. **Projectile motion** A gun with muzzle velocity of 400 m/s is fired at an angle of 8° above the horizontal. Find the horizontal and vertical components of the velocity.

Solution. Let $v + u$ be the velocity of the bullet, where \vec{v} is the vertical component and \vec{u} is the horizontal component.




$$\begin{aligned}
 \frac{\|u\|}{\|u + v\|} &= \cos 8^\circ \\
 \|u\| &= 400 \cos 8^\circ \\
 &= 396.107 \text{ m/s}
 \end{aligned}$$


$$\begin{aligned}
 \frac{\|v\|}{\|u + v\|} &= \sin 8^\circ \\
 \|v\| &= 400 \sin 8^\circ \\
 &= 55.669 \text{ m/s}
 \end{aligned}$$

■

26. **Inclined plane** Suppose that a box is being towed up an inclined plane as shown in the figure. Find the force w needed to make the component of the force parallel to the inclined plane equal to 2.5 N.



Solution.



$$\begin{aligned}\|w\| \cos(33^\circ - 15^\circ) &= 2.5 \\ \|w\| &= \frac{2.5}{\cos 18^\circ}\end{aligned}$$

$$\begin{aligned}w &= \|w\| \langle \cos 33^\circ, \sin 33^\circ \rangle \\ &= \frac{2.5}{\cos 18^\circ} \langle \cos 33^\circ, \sin 33^\circ \rangle \\ &= \langle 2.205, 1.432 \rangle\end{aligned}$$

■

29. **Orthogonal unit vectors** If \mathbf{u}_1 and \mathbf{u}_2 are orthogonal unit vectors and $\mathbf{v} = a\mathbf{u}_1 + b\mathbf{u}_2$, find $\mathbf{v} \cdot \mathbf{u}_1$.

Solution.

$$\begin{aligned}\mathbf{v} \cdot \mathbf{u}_1 &= (a\mathbf{u}_1 + b\mathbf{u}_2) \cdot \mathbf{u}_1 \\ &= a\mathbf{u}_1 \cdot \mathbf{u}_1 + b\mathbf{u}_2 \cdot \mathbf{u}_1 \\ &= a\|\mathbf{u}_1\|^2 + b\mathbf{u}_2 \cdot \mathbf{u}_1 \\ &= a(1) + b(0) \\ &= a\end{aligned}$$

■

30. A force $\mathbf{F} = 2\mathbf{i} + \mathbf{j} - 3\mathbf{k}$ is applied to a spacecraft with velocity vector $\mathbf{v} = 3\mathbf{i} - \mathbf{j}$. Express \mathbf{F} as a sum of a vector parallel to \mathbf{v} and a vector orthogonal to \mathbf{v} .

Solution. The vector parallel to \mathbf{v} is

$$\begin{aligned}\text{proj}_{\mathbf{v}} \mathbf{F} &= \left(\frac{\mathbf{F} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \left(\frac{5}{10} \right) (3\mathbf{i} - \mathbf{j}) \\ &= \left(\frac{3}{2} \right) \mathbf{i} - \left(\frac{1}{2} \right) \mathbf{j}\end{aligned}$$

The vector orthogonal to \mathbf{v} is

$$\begin{aligned}\mathbf{F} - \text{proj}_{\mathbf{v}} \mathbf{F} &= (2\mathbf{i} + \mathbf{j} - 3\mathbf{k}) - \left(\frac{3}{2} \right) \mathbf{i} + \left(\frac{1}{2} \right) \mathbf{j} \\ &= \left(\frac{1}{2} \right) \mathbf{i} + \left(\frac{3}{2} \right) \mathbf{j} - 3\mathbf{k}\end{aligned}$$

■

Equations for Lines in the Plane

33. **Line perpendicular to a vector** Show that $\mathbf{v} = a\mathbf{i} + b\mathbf{j}$ is perpendicular to the line $ax + by = c$ by establishing that the slope of the vector \mathbf{v} is the negative reciprocal of the slope of the given line.

Solution. Let $P(x_1, y_1)$ and $Q(x_2, y_2)$ be two points on the line $ax + by = c$. Then,

$$\begin{aligned}P(x_1, y_1) &= \left(x_1, \frac{c - ax_1}{b} \right) \\ Q(x_2, y_2) &= \left(x_2, \frac{c - ax_2}{b} \right)\end{aligned}$$

$$\begin{aligned}
\overrightarrow{PQ} &= \langle x_2 - x_1, y_2 - y_1 \rangle \\
&= \left\langle x_2 - x_1, \frac{c - ax_2}{b} - \frac{c - ax_1}{b} \right\rangle \\
&= \left\langle x_2 - x_1, \frac{a(x_1 - x_2)}{b} \right\rangle
\end{aligned}$$

$$\begin{aligned}
\overrightarrow{PQ} \cdot v &= \left\langle x_2 - x_1, \frac{a(x_1 - x_2)}{b} \right\rangle \cdot \langle a, b \rangle \\
&= a(x_2 - x_1) + \frac{a(x_1 - x_2)}{b} \cdot b \\
&= a(x_2 - x_1) - a(x_2 - x_1) \\
&= 0
\end{aligned}$$

Therefore, \mathbf{v} is perpendicular to the line $ax + by = c$. ■

Alternatively, the slope of the vector \mathbf{v} is $\frac{b}{a}$, while the slope of the line $ax + by = c$ is $-\frac{a}{b}$. Therefore, the vector \mathbf{v} is perpendicular to the line $ax + by = c$, since slope of the vector \mathbf{v} is the negative reciprocal of the slope of the given line. ■

34. **Line parallel to a vector** Show that the vector $\mathbf{v} = a\mathbf{i} + b\mathbf{j}$ is parallel to the line $bx - ay = c$ by establishing that the slope of the line segment representing \mathbf{v} is the same as the slope of the given line.

Solution. The slope of the vector v is $\frac{b}{a}$, while the slope of the given line is also $\frac{b}{a}$. Therefore, the vector \mathbf{v} is parallel to the line $bx - ay = c$. ■

Work

43. **Work along a line** Find the work done by a force $\mathbf{F} = 5\mathbf{i}$ (magnitude 5 N) in moving an object along the line from the origin to the point $(1, 1)$ (distance in meters).

Solution. Let $\vec{D} = \langle 1, 1 \rangle$ be the displacement vector of the object.

$$\begin{aligned}
W &= \vec{F} \cdot \vec{D} \\
&= \langle 5, 0 \rangle \cdot \langle 1, 1 \rangle \\
&= 5 \text{ J}
\end{aligned}$$

44. **Locomotive The Union Pacific's Big Boy** locomotive could pull 6000-tonne trains with a tractive effort (pull) of 602,148 N. At this level of effort, about how much work did Big Boy do on the (approximately straight) 605 – km journey from San Francisco to Los Angeles?

Solution.

$$\begin{aligned}
 W &= \vec{F} \cdot \vec{D} \\
 &= \|\vec{F}\| \|\vec{D}\| \cos \theta \\
 &= 602148 \cdot 605000 \cdot \cos 0^\circ \\
 &= 3.6429954 \times 10^{11} \text{ J}
 \end{aligned}$$

■


45. **Inclined plane** How much work does it take to slide a crate 20 m along a loading dock by pulling on it with a 200N force at an angle of 30° from the horizontal?

Solution.

$$\begin{aligned}
 W &= \vec{F} \cdot \vec{D} \\
 &= \|\vec{F}\| \|\vec{D}\| \cos \theta \\
 &= 200 \cdot 20 \cdot \cos 30^\circ \\
 &= 3464.102 \text{ J}
 \end{aligned}$$

■

46. **Sailboat** The wind passing over a boat's sail exerted a 1000 N magnitude force \mathbf{F} as shown here. How much work did the wind perform in moving the boat forward 1 km? Answer in joules.



Solution.

$$\begin{aligned}
 W &= \vec{F} \cdot \vec{D} \\
 &= \|\vec{F}\| \|\vec{D}\| \cos \theta \\
 &= 1000 \cdot 1000 \cdot \cos 60^\circ \\
 &= 500000 \text{ J}
 \end{aligned}$$

■

Selected Exercises

Source: *Larson Calculus 11th Ed. Exercise 11.3*

55. **Finding an Angle** Find the angle between a cube's diagonal and one of its edges.

Solution. Let a be the length of the cube's edge, then the length of the cube's diagonal is $\sqrt{3}a$. Let one side of the cube $\vec{u} = \langle a, 0, 0 \rangle$ and the diagonal of the cube $\vec{v} = \langle a, a, a \rangle$, then

$$\begin{aligned}\cos \theta &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} \\ &= \frac{a^2 + 0 + 0}{a\sqrt{3}a} \\ &= \frac{1}{\sqrt{3}} \\ \theta &= \arccos\left(\frac{1}{\sqrt{3}}\right) \\ &= 54.74^\circ\end{aligned}$$

■

56. **Finding an Angle** Find the angle between the diagonal of a cube and the diagonal of one of its sides.

Solution. Let a be the length of the cube's edge, then the length of the cube's diagonal is $\sqrt{3}a$, and the length of the diagonal of one of its sides is $\sqrt{2}a$. Let the diagonal of the cube $\vec{u} = \langle a, a, a \rangle$ and the diagonal of one of its sides $\vec{v} = \langle a, a, 0 \rangle$, then


$$\begin{aligned}\cos \theta &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} \\ &= \frac{a^2 + a^2 + 0}{\sqrt{3}a\sqrt{2}a} \\ &= \frac{2}{\sqrt{6}} \\ \theta &= \arccos\left(\frac{2}{\sqrt{6}}\right) \\ &= 35.26^\circ\end{aligned}$$

■

57. **Braking Load** A truck weighing 22 metric tons is parked on a 10° slope (see figure). Assume the only force to overcome is that due to gravity. Find (a) the force required to keep the truck from rolling down the hill and (b) the force perpendicular to the hill.

Solution.

(a)



Let \vec{F} be the force required for the truck to roll down the hill, then \vec{F} is the projection of the force due to gravity $\vec{u} = \langle 0, -22000 \rangle$ onto the hill $\vec{v} = \langle \cos 10^\circ, \sin 10^\circ \rangle$.

$$\begin{aligned}
 \vec{F} &= \text{proj}_{\vec{v}} \vec{u} \\
 &= \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v} \\
 &= \frac{(-22000)(\sin 10^\circ)}{1} \cdot \langle \cos 10^\circ, \sin 10^\circ \rangle \\
 &= -3820.26 \langle \cos 10^\circ, \sin 10^\circ \rangle
 \end{aligned}$$


Since a force of $\|\vec{F}\| = 3.82$ metric tons is required for the truck to roll down the hill, the same force of 3.82 metric tons is required to keep the truck from rolling down the hill. ■

- (b) Let \vec{w} be the force perpendicular to the hill, then

$$\begin{aligned}
 \vec{u} &= \vec{F} + \vec{w} \\
 \vec{w} &= \vec{u} - \vec{F} \\
 &= \langle 0, -22000 \rangle - ((-22000)(\sin 10^\circ) \langle \cos 10^\circ, \sin 10^\circ \rangle) \\
 &= \langle 22000(\sin 10^\circ) \cos 10^\circ, -22000 + 22000 \sin 10^\circ \sin 10^\circ \rangle \\
 &= \langle 3762.222, -21336.619 \rangle \\
 \|\vec{w}\| &= \sqrt{(3762.222)^2 + (-21336.619)^2} \\
 &= 21.67 \text{ metric tons}
 \end{aligned}$$

■

59. **Work** An object is pulled 10 meters across a floor using a force of 85 newtons. The direction of the force is 60° above the horizontal (see figure). Find the work done.



Solution. Let $\vec{F} = \langle 85 \cos 60^\circ, 85 \sin 60^\circ \rangle$ be the force applied to the object, and $\vec{D} = \langle 10, 0 \rangle$ be the displacement vector of the object, then

$$\begin{aligned} W &= \vec{F} \cdot \vec{D} \\ &= \langle 85 \cos 60^\circ, 85 \sin 60^\circ \rangle \cdot \langle 10, 0 \rangle \\ &= 425 \text{ joules} \end{aligned}$$

61. **Work** A car is towed using a force of 1600 newtons. The chain used to pull the car makes a 25° angle with the horizontal. Find the work done in towing the car 2 kilometers.



Solution. Let $\vec{F} = \langle 1600 \cos 25^\circ, 1600 \sin 25^\circ \rangle$ be the force applied to the car, and $\vec{D} = \langle 2000, 0 \rangle$ be the displacement vector of the car, then

$$\begin{aligned} W &= \vec{F} \cdot \vec{D} \\ &= \langle 1600 \cos 25^\circ, 1600 \sin 25^\circ \rangle \cdot \langle 2000, 0 \rangle \\ &= 2900 \text{ kJ} \end{aligned}$$


■

Chapter 4

Projections

Given two vectors \vec{v} and \vec{u} . Construct a line from the terminal point of \vec{u} perpendicular to \vec{v} . The vector that starts from the initial point of \vec{u} and ends at the intersection of the line and \vec{v} is called the **projection of \vec{u} onto \vec{v}** , which is also known as the **vector component of \vec{u} along \vec{v}** .

Construct another vector from the initial point of \vec{u} that is orthogonal to \vec{v} , and the projection of \vec{u} onto that vector is called the **vector component of \vec{u} orthogonal to \vec{v}** . The visual representation is shown below.



From the diagram, it is not hard to see that

$$\vec{u} = \vec{w}_1 + \vec{w}_2$$

Hence, the vector component of \vec{u} orthogonal to \vec{v} is given by

$$\vec{w}_2 = \vec{u} - \vec{w}_1$$

The projection of \vec{u} onto \vec{v} is given by

$$\vec{w}_1 = \text{proj}_{\vec{v}} \vec{u} = \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

Example 1. Find the projection of $\vec{u} = \langle 6, 7 \rangle$ onto $\vec{v} = \langle 1, 4 \rangle$. Hence, find the vector component of \vec{u} orthogonal to \vec{v} .

$$\begin{aligned} \text{proj}_{\vec{v}} \vec{u} &= \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v} \\ &= \left(\frac{6(1) + 7(4)}{1^2 + 4^2} \right) \langle 1, 4 \rangle \\ &= \left(\frac{34}{17} \right) \langle 1, 4 \rangle \\ &= \langle 2, 8 \rangle \end{aligned}$$

$$\begin{aligned} \vec{w}_2 &= \vec{u} - \vec{w}_1 \\ &= \langle 6, 7 \rangle - \langle 2, 8 \rangle \\ &= \langle 4, -1 \rangle \end{aligned}$$

Example 2. Find the projection of $\vec{u} = 2i + 3j$ onto $\vec{v} = 5i + j$. Hence, find the vector component of \vec{u} orthogonal to \vec{v} .


$$\begin{aligned} \text{proj}_{\vec{v}} \vec{u} &= \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v} \\ &= \left(\frac{2(5) + 3(1)}{5^2 + 1^2} \right) (5i + j) \\ &= \left(\frac{13}{26} \right) (5i + j) \\ &= \left(\frac{5}{2} \right) i + \left(\frac{1}{2} \right) j \end{aligned}$$

$$\begin{aligned} \vec{w}_2 &= \vec{u} - \vec{w}_1 \\ &= (2i + 3j) - \left(\left(\frac{5}{2} \right) i + \left(\frac{1}{2} \right) j \right) \\ &= \left(-\frac{1}{2} \right) i + \left(\frac{5}{2} \right) j \end{aligned}$$

Notes: Selected exercises are mixed in the exercises of the previous chapters.

Chapter 5

Direction Cosines and Direction Angles



Direction angles are angles that a vector makes with the unit vectors \hat{i} , \hat{j} and \hat{k} , denoted by α , β and γ respectively. In other words, *alpha*, β and γ are the angles between the vector and the x , y and z axis respectively.

Recall the formula for the dot product of two vectors \vec{v} and \vec{u}

$$\vec{v} \cdot \vec{u} = \|\vec{v}\| \|\vec{u}\| \cos \theta$$

where θ is the angle between the two vectors.

Let's calculate the dot product of \vec{v} and \hat{i} .

$$\vec{v} \cdot \hat{i} = \|\vec{v}\| \|\hat{i}\| \cos \alpha$$

$$\langle v_1, v_2, v_3 \rangle \cdot \langle 1, 0, 0 \rangle = \|\vec{v}\| \times 1 \times \cos \alpha$$

$$v_1 = \|\vec{v}\| \cos \alpha$$

$$\cos \alpha = \frac{v_1}{\|\vec{v}\|}$$

Similarly, we can calculate the dot product of \vec{v} and \hat{j} and \hat{k} in the same way. Hence, we can conclude that

$$\cos \alpha = \frac{v_1}{\|\vec{v}\|} \quad \cos \beta = \frac{v_2}{\|\vec{v}\|} \quad \cos \gamma = \frac{v_3}{\|\vec{v}\|}$$

These are called the **direction cosines** of \vec{v} .

We can also express any unit vector \hat{v} in terms of its direction cosines.

$$\begin{aligned} \vec{v} &= v_1\hat{i} + v_2\hat{j} + v_3\hat{k} \\ \frac{\vec{v}}{\|\vec{v}\|} &= \frac{v_1}{\|\vec{v}\|}\hat{i} + \frac{v_2}{\|\vec{v}\|}\hat{j} + \frac{v_3}{\|\vec{v}\|}\hat{k} \\ \hat{v} &= \cos \alpha \hat{i} + \cos \beta \hat{j} + \cos \gamma \hat{k} \end{aligned}$$

If we take the magnitude of \hat{v} , we get

$$\|\hat{v}\| = \sqrt{\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma} = 1$$

Squaring both sides, we get

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

Example 1. Find the direction cosines and direction angles of the vector $\vec{u} = \hat{i} + 8\hat{j} + 4\hat{k}$.

$$\begin{aligned} \|\vec{u}\| &= \sqrt{1^2 + 8^2 + 4^2} \\ &= \sqrt{81} \\ &= 9 \end{aligned}$$

$$\begin{aligned} \cos \alpha &= \frac{v_1}{\|\vec{v}\|} = \frac{1}{9} & \alpha &= \cos^{-1} \left(\frac{1}{9} \right) \approx 1.459 \text{ rad} \\ \cos \beta &= \frac{v_2}{\|\vec{v}\|} = \frac{8}{9} & \beta &= \cos^{-1} \left(\frac{8}{9} \right) \approx 0.476 \text{ rad} \\ \cos \gamma &= \frac{v_3}{\|\vec{v}\|} = \frac{4}{9} & \gamma &= \cos^{-1} \left(\frac{4}{9} \right) \approx 1.110 \text{ rad} \end{aligned}$$

Chapter 6

Cross Product

Let $\vec{v} = \langle v_1, v_2, v_3 \rangle$ and $\vec{u} = \langle u_1, u_2, u_3 \rangle$ be two vectors in \mathbb{R}^3 (three-dimensional space), then the cross product of \vec{v} and \vec{u} is given by


$$\vec{v} \times \vec{u} = (v_2u_3 - v_3u_2)\hat{i} + (v_3u_1 - v_1u_3)\hat{j} + (v_1u_2 - v_2u_1)\hat{k}$$

The final result of the cross product is a vector, hence the name **vector product**.

It can also be calculated using the determinant of a 3×3 matrix as shown below.

$$\begin{aligned} \vec{v} \times \vec{u} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ v_1 & v_2 & v_3 \\ u_1 & u_2 & u_3 \end{vmatrix} \\ &= \begin{vmatrix} v_2 & v_3 \\ u_2 & u_3 \end{vmatrix} \hat{i} - \begin{vmatrix} v_1 & v_3 \\ u_1 & u_3 \end{vmatrix} \hat{j} + \begin{vmatrix} v_1 & v_2 \\ u_1 & u_2 \end{vmatrix} \hat{k} \\ &= (v_2u_3 - v_3u_2)\hat{i} - (v_1u_3 - v_3u_1)\hat{j} + (v_1u_2 - v_2u_1)\hat{k} \end{aligned}$$

Geometrically speaking, the cross product of two vectors \vec{v} and \vec{u} is a vector that is orthogonal to both \vec{v} and \vec{u} , and its direction is given by the right-hand rule. That is, $\vec{v} \times \vec{u} \neq \vec{u} \times \vec{v}$.



Another property of the cross product is that the magnitude of the cross product of two vectors \vec{v} and \vec{u} is given by


$$\|\vec{v} \times \vec{u}\| = \|\vec{v}\| \|\vec{u}\| \sin \theta$$

where θ is the angle between the two vectors.

The magnitude of the cross product of two vectors \vec{v} and \vec{u} is the area of the parallelogram formed by \vec{v} and \vec{u} .

Proof. Note that $\cos \theta = \frac{\vec{v} \cdot \vec{u}}{\|\vec{v}\| \|\vec{u}\|}$, hence

$$\begin{aligned}
 \|\vec{v} \times \vec{u}\| &= \|\vec{v}\| \|\vec{u}\| \sin \theta \\
 &= \|\vec{v}\| \|\vec{u}\| \sqrt{1 - \cos^2 \theta} \\
 &= \|\vec{v}\| \|\vec{u}\| \sqrt{1 - \left(\frac{\vec{v} \cdot \vec{u}}{\|\vec{v}\| \|\vec{u}\|} \right)^2} \\
 &= \sqrt{\|\vec{v}\|^2 \|\vec{u}\|^2 - (\vec{v} \cdot \vec{u})^2} \\
 &= \sqrt{(v_1^2 + v_2^2 + v_3^2)(u_1^2 + u_2^2 + u_3^2) - (v_1 u_1 + v_2 u_2 + v_3 u_3)^2} \\
 &= \sqrt{(v_2 u_3 - v_3 u_2)^2 + (v_3 u_1 - v_1 u_3)^2 + (v_1 u_2 - v_2 u_1)^2} \\
 &= \|\vec{v} \times \vec{u}\|
 \end{aligned}$$



Since the height of the parallelogram is $\|\vec{u}\| \sin \theta$ and the base of the parallelogram is $\|\vec{v}\|$, the area of the parallelogram is given By

$$\begin{aligned}
 \text{Area} &= (\text{base})(\text{height}) \\
 &= \|\vec{v}\| \|\vec{u}\| \sin \theta \\
 &= \|\vec{v} \times \vec{u}\| \quad \blacksquare
 \end{aligned}$$

Example 1. Find the cross product of the vectors $\vec{v} = \langle 1, 2, 3 \rangle$ and $\vec{u} = \langle -1, 0, 4 \rangle$.

$$\begin{aligned}
 \vec{v} \times \vec{u} &= \begin{vmatrix} i & j & k \\ 1 & 2 & 3 \\ -1 & 0 & 4 \end{vmatrix} \\
 &= \begin{vmatrix} 2 & 3 \\ 0 & 4 \end{vmatrix} i - \begin{vmatrix} 1 & 3 \\ -1 & 4 \end{vmatrix} j + \begin{vmatrix} 1 & 2 \\ -1 & 0 \end{vmatrix} k \\
 &= (2(4) - 3(0))i - (1(4) - 3(-1))j + (1(0) - 2(-1))k \\
 &= 8i - 7j + 2k
 \end{aligned}$$

Example 2. Find the unit vector that is orthogonal to both $\vec{v} = 2i - 3j + k$ and $\vec{u} = i + 2j - k$.

$$\begin{aligned}
 \vec{v} \times \vec{u} &= \begin{vmatrix} i & j & k \\ 2 & -3 & 1 \\ 1 & 2 & -1 \end{vmatrix} \\
 &= \begin{vmatrix} -3 & 1 \\ 2 & -1 \end{vmatrix} i - \begin{vmatrix} 2 & 1 \\ 1 & -1 \end{vmatrix} j + \begin{vmatrix} 2 & -3 \\ 1 & 2 \end{vmatrix} k \\
 &= (-3(-1) - 1(2))i - (2(-1) - 1(1))j + (2(2) - 1(-3))k \\
 &= i + 3j + 7k
 \end{aligned}$$

$$\begin{aligned}
 \|\vec{v} \times \vec{u}\| &= \sqrt{1^2 + 3^2 + 7^2} \\
 &= \sqrt{59}
 \end{aligned}$$

$$\begin{aligned}
 \text{Unit vector} &= \frac{\vec{v} \times \vec{u}}{\|\vec{v} \times \vec{u}\|} \\
 &= \frac{1}{\sqrt{59}}i + \frac{3}{\sqrt{59}}j + \frac{7}{\sqrt{59}}k
 \end{aligned}$$

Selected Exercises

Source: *Larson Calculus 11th Ed. Exercise 11.4*

Area In Exercises and 28, find the area of the triangle with the given vertices. (Hint: $\frac{1}{2}\|u \times v\|$ is the area of the triangle having u and v as adjacent sides.)

25. $A(0, 0, 0), B(1, 0, 3), C(-3, 2, 0)$

Solution.


$$\begin{aligned}\vec{u} &= \langle 1 - 0, 0 - 0, 3 - 0 \rangle = \langle 1, 0, 3 \rangle \\ \vec{v} &= \langle -3 - 0, 2 - 0, 0 - 0 \rangle = \langle -3, 2, 0 \rangle \\ \vec{u} \times \vec{v} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & 3 \\ -3 & 2 & 0 \end{vmatrix} \\ &= \hat{i} \begin{vmatrix} 0 & 3 \\ 2 & 0 \end{vmatrix} - \hat{j} \begin{vmatrix} 1 & 3 \\ -3 & 0 \end{vmatrix} + \hat{k} \begin{vmatrix} 1 & 0 \\ -3 & 2 \end{vmatrix} \\ &= \hat{i}(-6) - \hat{j}(9) + \hat{k}(2) \\ &= \langle -6, -9, 2 \rangle\end{aligned}$$

Hence, the area of the triangle is

$$\frac{1}{2} \|\vec{u} \times \vec{v}\| = \frac{1}{2} \sqrt{(-6)^2 + (-9)^2 + 2^2} = \frac{1}{2} \sqrt{121} = \frac{11}{2} \text{ sq. units}$$

■

27. **Torque** The brakes on a bicycle are applied using a downward force of 100 newtons on the pedal when the crank makes a 40° angle with the horizontal (see figure). The crank is 17 centimeters in length. Find the torque at P .




Solution. Let the force $\vec{F} = \langle 0, 0, -100 \rangle$ and the position vector of the pedal $\vec{r} = \langle 0, 0.17 \cos 40^\circ, 0.17 \sin 40^\circ \rangle$.

$$\begin{aligned}
 \vec{\tau} &= \vec{r} \times \vec{F} \\
 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 0.17 \cos 40^\circ & 0.17 \sin 40^\circ \\ 0 & 0 & -100 \end{vmatrix} \\
 &= \hat{i} \begin{vmatrix} 0.17 \cos 40^\circ & 0.17 \sin 40^\circ \\ 0 & -100 \end{vmatrix} - \hat{j} \begin{vmatrix} 0 & 0.17 \sin 40^\circ \\ 0 & -100 \end{vmatrix} + \hat{k} \begin{vmatrix} 0 & 0.17 \cos 40^\circ \\ 0 & 0 \end{vmatrix} \\
 &= \hat{i}(-17 \cos 40^\circ) - \hat{j}(0) + \hat{k}(0) \\
 &= \langle -17 \cos 40^\circ, 0, 0 \rangle \\
 \|\vec{\tau}\| &= \sqrt{(-17 \cos 40^\circ)^2 + 0^2 + 0^2} = 17 \cos 40^\circ \approx 13.023 \text{ J}
 \end{aligned}$$

■

29. **Optimization** A force of 180 newtons acts on the bracket shown in the figure.



- (a) Determine the vector \overrightarrow{AB} and the vector \mathbf{F} representing the force. (\mathbf{F} will be in terms of θ .)

Solution. The horizontal and vertical distances between A and B are -40cm and 32cm respectively. Hence, $\overrightarrow{AB} = -0.4j + 0.32k$. ■

- (b) Find the magnitude of the moment about A by evaluating $\|\vec{AB} \times \mathbf{F}\|$.

Solution.

$$\begin{aligned}\vec{AB} \times \mathbf{F} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & -0.4 & 0.32 \\ 0 & -180 \cos \theta & -180 \sin \theta \end{vmatrix} \\ &= \hat{i} \begin{vmatrix} -0.4 & 0.32 \\ -180 \cos \theta & -180 \sin \theta \end{vmatrix} - \hat{j} \begin{vmatrix} 0 & 0.32 \\ 0 & -180 \sin \theta \end{vmatrix} + \hat{k} \begin{vmatrix} 0 & -0.4 \\ 0 & -180 \cos \theta \end{vmatrix} \\ &= \hat{i}(72 \sin \theta + 57.6 \cos \theta) - \hat{j}(0) + \hat{k}(0) \\ &= \langle 72 \sin \theta + 57.6 \cos \theta, 0, 0 \rangle\end{aligned}$$

$$\begin{aligned}\|\vec{AB} \times \mathbf{F}\| &= \sqrt{(72 \sin \theta + 57.6 \cos \theta)^2 + 0^2 + 0^2} \\ &= |72 \sin \theta + 57.6 \cos \theta|\end{aligned}$$

■

- (c) Use the result of part (b) to determine the magnitude of the moment when $\theta = 30^\circ$.

Solution. When $\theta = 30^\circ$,

$$\|\vec{AB} \times \mathbf{F}\| = |72 \sin 30^\circ + 57.6 \cos 30^\circ| = 85.883 \text{ J}$$

- (d) Use the result of part (b) to determine the angle θ when the magnitude of the moment is maximum. At that angle, what is the relationship between the vectors \mathbf{F} and \vec{AB} ? $\frac{d}{d\theta}(|72 \sin \theta + 57.6 \cos \theta|) = 0$. Hence,

$$\frac{d}{d\theta}(|72 \sin \theta + 57.6 \cos \theta|) = \frac{72 \cos \theta - 57.6 \sin \theta}{|72 \sin \theta + 57.6 \cos \theta|} = 0$$

$$72 \cos \theta = 57.6 \sin \theta$$

$$\tan \theta = \frac{72}{57.6} = \frac{5}{4}$$

$$\theta = \tan^{-1}\left(\frac{5}{4}\right) \approx 51.340^\circ$$


$$\begin{aligned}F \cdot \vec{AB} &= 180 \cos \theta \times -0.4 + 180 \sin \theta \times 0.32 \\ &= -72 \cos\end{aligned}$$

When $\theta = 51.340^\circ$, \mathbf{F} and \vec{AB} are perpendicular to each other.

■

Chapter 7

Distance in Space



The distance between two points $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$ in space is essentially the magnitude of the vector $\overrightarrow{P_1P_2}$, which is given by


$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Example 1. Find the distance between the points $P_1(-3, 2, 5)$ and $P_2(4, 0, 8)$ in space.

$$\begin{aligned} d &= \sqrt{(4 - (-3))^2 + (0 - 2)^2 + (8 - 5)^2} \\ &= \sqrt{7^2 + (-2)^2 + 3^2} \\ &= \sqrt{62} \end{aligned}$$

Chapter 8

Lines in Space



To find the equation of a line in space, we need a point $P(x_1, y_1, z_1)$ on the line and a vector $\vec{v} = \langle a, b, c \rangle$ that is parallel to the line. The vector \vec{v} is called the **direction vector** of the line, while a , b and c are called the **direction numbers**.

Since \vec{v} is parallel to L , the vector \overrightarrow{PQ} is also parallel to L , where $Q(x, y, z)$ is any point on L . Hence, \overrightarrow{PQ} is a scalar multiple of \vec{v} , that is,

$$\begin{aligned}\overrightarrow{PQ} &= t\vec{v} \\ \langle x - x_1, y - y_1, z - z_1 \rangle &= t\langle a, b, c \rangle \\ &= \langle at, bt, ct \rangle\end{aligned}$$

Comparing both sides, we get

$$x - x_1 = at \quad y - y_1 = bt \quad z - z_1 = ct$$

Rearranging the equations, we get the **parametric equations** of the line L .

$$x = x_1 + at \quad y = y_1 + bt \quad z = z_1 + ct$$

If $a, b, c \neq 0$, we can solve for t for each of the parametric equations to get the **symmetric equations** of the line L .

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$$

Example 1. Find the equation of the line that passes through the point $P(-1, 4, 5)$ and is parallel to the vector $\vec{v} = 4i - j$.

$$\begin{aligned} x &= -1 + 4t \\ y &= 4 - t \\ z &= 5 \end{aligned}$$

Note that it is impossible to find the symmetric equation of the line since $c = 0$.

Example 2. L passes through the point $P(2, 7, 1)$ and is parallel to the vector $\vec{v} = \langle -2, -4, 6 \rangle$. Find the parametric and symmetric equations of L .

$$\begin{aligned} x &= 2 - 2t \\ y &= 7 - 4t \\ z &= 1 + 6t \end{aligned}$$

$$\frac{x - 2}{-2} = \frac{y - 7}{-4} = \frac{z - 1}{6}$$

Example 3. Find the equation of the line passing through the point $(1, 0, 1)$ and parallel to the line given by the parametric equations

$$x = 3 + 3t \quad y = 5 - 2t \quad z = -7 + t$$

The line is parallel to the vector $\vec{v} = \langle 3, -2, 1 \rangle$. Hence, the equation of the line is given by

$$x = 1 + 3t \quad y = -2t \quad z = 1 + t$$

Also, the symmetric equations of the line is given by

$$\frac{x - 1}{3} = \frac{y}{-2} = \frac{z - 1}{1}$$

Example 4. Find the equation of the line passing through points $(7, -2, 6)$ and $(-3, 0, 6)$.

$$\begin{aligned}\vec{v} &= \langle -3 - 7, 0 - (-2), 6 - 6 \rangle \\ &= \langle -10, 2, 0 \rangle\end{aligned}$$


The equation of the line is given by

$$x = 7 - 10t \quad y = -2 + 2t \quad z = 6$$

There is no symmetric equation of the line since $c = 0$.

Chapter 9

Planes in Space



To find the equation of a plane in space, we need a point $P(x_1, y_1, z_1)$ on the plane and a vector $\vec{n} = \langle a, b, c \rangle$ that is orthogonal to the plane, called the **normal vector** of the plane.

For any point $Q(x, y, z)$ on the plane, the vector \overrightarrow{PQ} is orthogonal to \vec{n} , that is,

$$\overrightarrow{PQ} \cdot \vec{n} = 0$$

$$\langle x - x_1, y - y_1, z - z_1 \rangle \cdot \langle a, b, c \rangle = 0$$

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

This equation is called the **standard form** of the equation of the plane. Regrouping the terms, we obtain the **general form** of equation of the plane.

$$ax + by + cz + d = 0$$

where $d = -ax_1 - by_1 - cz_1$.

Example 1. Find the equation of the plane that passes through the point $P(4, 5, -7)$ and is perpendicular to the vector $\vec{n} = \hat{j}$.

$$\begin{aligned}\vec{n} &= \langle 0, 1, 0 \rangle \\ 0(x - 4) + 1(y - 5) + 0(z + 7) &= 0 \\ y - 5 &= 0 \\ y &= 5\end{aligned}$$

Example 2. Find the equation of the plane that passes through the point $P(0, 7, 0)$ and is perpendicular to the vector $\vec{n} = 3\hat{i} + 8\hat{k}$.

$$\begin{aligned}\vec{n} &= \langle 3, 0, 8 \rangle \\ 3(x - 0) + 0(y - 7) + 8(z - 0) &= 0 \\ 3x + 8z &= 0\end{aligned}$$

Example 3. Given three points $(0, 0, 0)$, $(2, 0, 7)$, and $(-2, -1, 7)$ in space, find the equation of the plane that passes through these points.

$$\begin{aligned}\vec{u} &= \langle 2 - 0, 0 - 0, 7 - 0 \rangle \\ &= \langle 2, 0, 7 \rangle \\ \vec{v} &= \langle -2 - 0, -1 - 0, 7 - 0 \rangle \\ &= \langle -2, -1, 7 \rangle \\ \vec{n} &= \vec{u} \times \vec{v} \\ &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & 7 \\ -2 & -1 & 7 \end{vmatrix} \\ &= \begin{vmatrix} 0 & 7 \\ -1 & 7 \end{vmatrix} \hat{i} - \begin{vmatrix} 2 & 7 \\ -2 & 7 \end{vmatrix} \hat{j} + \begin{vmatrix} 2 & 0 \\ -2 & -1 \end{vmatrix} \hat{k} \\ &= (0(7) - 7(-1))\hat{i} - (2(7) - (-2)(7))\hat{j} + (2(-1) - (-2)(0))\hat{k} \\ &= 7\hat{i} - 28\hat{j} - 2\hat{k} \\ &= \langle 7, -28, -2 \rangle\end{aligned}$$

$$\begin{aligned}7(x - 0) - 28(y - 0) - 2(z - 0) &= 0 \\ 7x - 28y - 2z &= 0\end{aligned}$$

Example 4. Find the equation of the plane that passes through $(4, 2, 1)$, $(-1, 8, 8)$ and is parallel to z -axis.

$$\begin{aligned}
 \vec{v} &= \langle -1 - 4, 8 - 2, 8 - 1 \rangle \\
 &= \langle -5, 6, 7 \rangle \\
 \vec{n} &= \vec{v} \times \hat{k} \\
 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -5 & 6 & 7 \\ 0 & 0 & 1 \end{vmatrix} \\
 &= \begin{vmatrix} 6 & 7 \\ 0 & 1 \end{vmatrix} \hat{i} - \begin{vmatrix} -5 & 7 \\ 0 & 1 \end{vmatrix} \hat{j} + \begin{vmatrix} -5 & 6 \\ 0 & 0 \end{vmatrix} \hat{k} \\
 &= (6(1) - 7(0))\hat{i} - (-5(1) - 7(0))\hat{j} + (-5(0) - 6(0))\hat{k} \\
 &= 6\hat{i} + 5\hat{j} \\
 &= \langle 6, 5, 0 \rangle \\
 6(x - 4) + 5(y - 2) + 0(z - 1) &= 0 \\
 6x + 5y - 34 &= 0
 \end{aligned}$$

Example 5. Find the equation of the plane such that the point $(2, 0, 1)$ and the line $\frac{x}{2} = \frac{y - 4}{-1} = \frac{z}{1}$ is on the plane.
 When $x = 0$, $y = 4$ and $z = 0$, hence the point $(0, 4, 0)$ is on the plane.

$$\begin{aligned}
 \vec{v} &= \langle 2 - 0, 0 - 4, 1 - 0 \rangle \\
 &= \langle 2, -4, 1 \rangle \\
 \vec{n} &= \langle 2, -1, 1 \rangle \times \langle 2, -4, 1 \rangle \\
 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 1 \\ 2 & -4 & 1 \end{vmatrix} \\
 &= \begin{vmatrix} -1 & 1 \\ -4 & 1 \end{vmatrix} \hat{i} - \begin{vmatrix} 2 & 1 \\ 2 & 1 \end{vmatrix} \hat{j} + \begin{vmatrix} 2 & -1 \\ 2 & -4 \end{vmatrix} \hat{k} \\
 &= (-1(1) - 1(-4))\hat{i} - (2(1) - 2(1))\hat{j} + (2(-4) - 2(-1))\hat{k} \\
 &= -3\hat{i} + 0\hat{j} + (-6)\hat{k} \\
 &= \langle -3, 0, -6 \rangle \\
 -3(x - 2) + 0(y - 0) - 6(z - 1) &= 0 \\
 -3x - 6z + 12 &= 0
 \end{aligned}$$

Example 6. Find the equation of the plane that passes through the points $(3, 4, 1)$ and $(3, 1, -7)$ and is perpendicular to the plane $8x + 9y + 3z = 13$.


The normal vector of the plane is $\langle 8, 9, 3 \rangle$. Since the target plane is perpendicular to the given plane, the normal vector of the given plane is parallel to the target plane.

$$\begin{aligned}
 \vec{u} &= \langle 8, 9, 3 \rangle \\
 \vec{v} &= \langle 3 - 3, 1 - 4, -7 - 1 \rangle \\
 &= \langle 0, -3, -8 \rangle \\
 \vec{n} &= \vec{u} \times \vec{v} \\
 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 8 & 9 & 3 \\ 0 & -3 & -8 \end{vmatrix} \\
 &= \begin{vmatrix} 9 & 3 \\ -3 & -8 \end{vmatrix} \hat{i} - \begin{vmatrix} 8 & 3 \\ 0 & -8 \end{vmatrix} \hat{j} + \begin{vmatrix} 8 & 9 \\ 0 & -3 \end{vmatrix} \hat{k} \\
 &= (9(-8) - 3(-3))\hat{i} - (8(-8) - 3(0))\hat{j} + (8(-3) - 9(0))\hat{k} \\
 &= -63\hat{i} + 64\hat{j} - 24\hat{k} \\
 &= \langle -63, 64, -24 \rangle \\
 &\quad -63(x - 3) + 64(y - 4) - 24(z - 1) = 0 \\
 &\quad \quad \quad -63x + 64y - 24z - 43 = 0
 \end{aligned}$$

Chapter 10

Cylindrical Coordinates

Cylindrical coordinates are an extension of polar coordinates into three dimensions. A point $P(x, y, z)$ in space is represented by the ordered triple (r, θ, z) , called the **cylindrical coordinates** of P . Here, (r, θ) are the polar representation of the projection of P in the xy -plane, and z is the direct distance from the (r, θ) to P .



To convert from Cartesian coordinates to cylindrical coordinates, we use the following equations.

$$r^2 = x^2 + y^2 \quad \tan \theta = \frac{y}{x} \quad z = z$$

To convert from cylindrical coordinates to Cartesian coordinates, we use the following equations.

$$x = r \cos \theta \quad y = r \sin \theta \quad z = z$$

Example 1. Convert $\left(4, \frac{5\pi}{6}, 3\right)$ from cylindrical coordinates to Cartesian coordinates.

$$\begin{aligned}x &= r \cos \theta \\&= 4 \cos \left(\frac{5\pi}{6}\right) \\&= -2\sqrt{3} \\y &= r \sin \theta \\&= 4 \sin \left(\frac{5\pi}{6}\right) \\&= 2 \\z &= z \\&= 3\end{aligned}$$

Hence, $(-2\sqrt{3}, 2, 3)$ is the Cartesian coordinates of $\left(4, \frac{5\pi}{6}, 3\right)$.

Example 2. Convert $(1, \sqrt{3}, 2)$ from Cartesian coordinates to cylindrical coordinates.


$$\begin{aligned}r &= \sqrt{x^2 + y^2} \\&= \sqrt{1^2 + \sqrt{3}^2} \\&= 2 \\ \theta &= \tan^{-1} \left(\frac{y}{x}\right) \\&= \tan^{-1} \left(\frac{\sqrt{3}}{1}\right) \\&= \frac{\pi}{3} \\z &= z \\&= 2\end{aligned}$$

Hence, $\left(2, \frac{\pi}{3}, 2\right)$ is the cylindrical coordinates of $(1, \sqrt{3}, 2)$.

Chapter 11

Spherical Coordinates

Spherical coordinates are an ordered triple $P = (\rho, \theta, \phi)$, where ρ is the distance between the origin and P ($\rho \geq 0$ since it is a distance), θ is the angle from cylindrical coordinates where $r \geq 0$ and ϕ is the angle between the positive z -axis and the line segment \overrightarrow{OP} ($0 \leq \phi \leq \pi$).



To convert from Cartesian coordinates to spherical coordinates, we use the following equations.

$$\rho^2 = x^2 + y^2 + z^2 \quad \tan \theta = \frac{y}{x} \quad \cos \phi = \frac{z}{\rho}$$

To convert from spherical coordinates to Cartesian coordinates, we use the following equations.

$$x = \rho \sin \phi \cos \theta \quad y = \rho \sin \phi \sin \theta \quad z = \rho \cos \phi$$

Example 1. Convert $\left(10, \frac{\pi}{6}, \frac{\pi}{4}\right)$ from spherical coordinates to Cartesian coordinates.

$$\begin{aligned}
 x &= \rho \sin \phi \cos \theta \\
 &= 10 \sin \left(\frac{\pi}{4}\right) \cos \left(\frac{\pi}{6}\right) \\
 &= \frac{5\sqrt{6}}{2} \\
 y &= \rho \sin \phi \sin \theta \\
 &= 10 \sin \left(\frac{\pi}{4}\right) \sin \left(\frac{\pi}{6}\right) \\
 &= \frac{5\sqrt{2}}{2} \\
 z &= \rho \cos \phi \\
 &= 10 \cos \left(\frac{\pi}{4}\right) \\
 &= 5\sqrt{2}
 \end{aligned}$$

Hence, $\left(\frac{5\sqrt{6}}{2}, \frac{5\sqrt{2}}{2}, 5\sqrt{2}\right)$ is the Cartesian coordinates of $\left(10, \frac{\pi}{6}, \frac{\pi}{4}\right)$.

Example 2. Convert $(-8, -8, \sqrt{19})$ from Cartesian coordinates to spherical coordinates.

$$\begin{aligned}
 \rho &= \sqrt{x^2 + y^2 + z^2} \\
 &= \sqrt{(-8)^2 + (-8)^2 + (\sqrt{19})^2} \\
 &= 7\sqrt{3} \\
 \theta &= \tan^{-1} \left(\frac{y}{x}\right) \\
 &= \tan^{-1} \left(\frac{-8}{-8}\right) \\
 &= \frac{5\pi}{4} \text{ (3rd quadrant since } x \text{ and } y \text{ are negative)} \\
 \phi &= \arccos \left(\frac{z}{\rho}\right) \\
 &= \arccos \left(\frac{\sqrt{19}}{7\sqrt{3}}\right)
 \end{aligned}$$

Hence, $\left(7\sqrt{3}, \frac{5\pi}{4}, \arccos \left(\frac{\sqrt{19}}{7\sqrt{3}}\right)\right)$ is the spherical coordinates of $(-8, -8, \sqrt{19})$.

Chapter 12

Vector-valued Functions

A vector valued-function is a function that maps a real number to a vector. In the plane, a vector-valued function is given by

$$\vec{r}(t) = \langle x(t), y(t) \rangle$$

and in space, a vector-valued function is given by

$$\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$$

Note that different functions can give the same curve. For example,

$$\vec{r}(t) = \langle \cos t, \sin t \rangle \quad t \in [0, 2\pi]$$

and

$$\vec{r}(t) = \langle \cos 2t, \sin 2t \rangle \quad t \in [0, 2\pi]$$

both give the unit circle.

The domain of a vector-valued function \vec{r} is the intersection of the domains of the component functions $x(t)$, $y(t)$ and $z(t)$. For example, given a vector-valued function $\vec{r}(t) = \frac{1}{t}\hat{i} + \frac{1}{t-1}\hat{j} + \frac{1}{\cos t}\hat{k}$ is $(-\infty, 0) \cup (0, 1) \cup (1, \infty)$. The domain of each component function is

$$D_{x(t)} = (-\infty, 0) \cup (0, \infty) \quad D_{y(t)} = (-\infty, 1) \cup (1, \infty) \quad D_{z(t)} = \mathbb{R}$$

Combining the domains of the component functions, we get the domain of the vector-valued function $\vec{r}(t)$

$$(D_{x(t)} \cap D_{y(t)} \cap D_{z(t)}) = (-\infty, 0) \cup (0, 1) \cup (1, \infty)$$

Example 1. Sketch the vector-valued function $\vec{r}(t) = \langle 2 \cos t, -3 \sin t \rangle$.

$$\begin{aligned} x &= 2 \cos t \\ \cos t &= \frac{x}{2} \\ y &= -3 \sin t \\ \sin t &= -\frac{y}{3} \\ \cos^2 t + \sin^2 t &= 1 \\ \left(\frac{x}{2}\right)^2 + \left(-\frac{y}{3}\right)^2 &= 1 \\ \frac{x^2}{4} + \frac{y^2}{9} &= 1 \end{aligned}$$


Hence, the graph of the vector-valued function $\vec{r}(t) = \langle 2 \cos t, -3 \sin t \rangle$ is an ellipse with major axis of length 6 and minor axis of length 4.

To find the orientation of the function, we can plot points in increasing values of t .

When $t = 0$, $\vec{r}(0) = \langle 2 \cos 0, -3 \sin 0 \rangle = \langle 2, 0 \rangle$.

When $t = \frac{\pi}{2}$, $\vec{r}\left(\frac{\pi}{2}\right) = \langle 2 \cos \frac{\pi}{2}, -3 \sin \frac{\pi}{2} \rangle = \langle 0, -3 \rangle$.

Hence, the orientation of the function is clockwise.



Example 2. Sketch the vector-valued function $\vec{r}(t) = \frac{t}{8}\hat{i} + (t-1)\hat{j}$.

$$x = \frac{t}{8}$$

$$t = 8x$$

$$y = t - 1$$

$$t = y + 1$$

$$8x = y + 1$$


$$y = 8x - 1$$

Hence, the graph of the vector-valued function $\vec{r}(t) = \frac{t}{8}\hat{i} + (t-1)\hat{j}$ is a line with y -intercept of -1 and slope of 8 .

When $t = 0$, $\vec{r}(0) = \frac{0}{8}\hat{i} + (0-1)\hat{j} = \langle 0, -1 \rangle$.

When $t = 8$, $\vec{r}(8) = \frac{8}{8}\hat{i} + (8-1)\hat{j} = \langle 1, 7 \rangle$.

Hence, the orientation of the function is going up.



Example 3. Represent $y = x + 9$ as vector-valued function.

$$\begin{aligned}x &= t \\y &= t + 9 \\\vec{r}(t) &= x(t)\hat{i} + y(t)\hat{j} \\&= t\hat{i} + (t + 9)\hat{j}\end{aligned}$$

Example 4. Represent $x^2 + y^2 = 64$ as vector-valued function.

$$\begin{aligned}x &= 8 \cos t \\y &= 8 \sin t \\\vec{r}(t) &= x(t)\hat{i} + y(t)\hat{j} \\&= 8 \cos t \hat{i} + 8 \sin t \hat{j}\end{aligned}$$

Example 5. Represent $(x - 2)^2 + (y + 1)^2 = 4$ as vector-valued function.

$$\begin{aligned}2 \cos t &= x - 2 \\x &= 2 \cos t + 2 \\2 \sin t &= y + 1 \\y &= 2 \sin t - 1 \\\vec{r}(t) &= x(t)\hat{i} + y(t)\hat{j} \\&= (2 \cos t + 2)\hat{i} + (2 \sin t - 1)\hat{j}\end{aligned}$$

Hence, we can conclude that the vector-valued function of a circle with radius r and centre (h, k) is

$$\vec{r}(t) = (h + r \cos t)\hat{i} + (k + r \sin t)\hat{j}$$

Example 6. Represent $\frac{x^2}{9} + \frac{y^2}{4} = 1$ as vector-valued function.

$$\begin{aligned}x &= 3 \cos t \\y &= 2 \sin t \\\vec{r}(t) &= x(t)\hat{i} + y(t)\hat{j} \\&= 3 \cos t \hat{i} + 2 \sin t \hat{j}\end{aligned}$$

Example 7. Represent $\frac{(x-1)^2}{4} + \frac{(y+2)^2}{25} = 1$ as vector-valued function.

$$x - 1 = 2 \cos t$$

$$x = 2 \cos t + 1$$

$$y + 2 = 5 \sin t$$

$$y = 5 \sin t - 2$$

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j}$$

$$= (2 \cos t + 1)\hat{i} + (5 \sin t - 2)\hat{j}$$

Hence, we can conclude that the vector-valued function of an ellipse with major axis of length $2a$ and minor axis of length $2b$ is

$$\vec{r}(t) = (h + a \cos t)\hat{i} + (k + b \sin t)\hat{j}$$

Example 8. Represent $\frac{x^2}{25} - \frac{y^2}{16} = 1$ as vector-valued function.

$$x = 5 \cosh t$$

$$y = 4 \sinh t$$

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j}$$

$$= 5 \cosh t \hat{i} + 4 \sinh t \hat{j}$$

Example 9. Represent $\frac{(x-1)^2}{4} - \frac{(y+7)^2}{9} = 1$ as vector-valued function.

$$x - 1 = 2 \cosh t$$

$$x = 2 \cosh t + 1$$

$$y + 7 = 3 \sinh t$$

$$y = 3 \sinh t - 7$$

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j}$$

$$= (2 \cosh t + 1)\hat{i} + (3 \sinh t - 7)\hat{j}$$

Hence, we can conclude that the vector-valued function of a hyperbola with centre (h, k) is

$$\vec{r}(t) = (h + a \cosh t)\hat{i} + (k + b \sinh t)\hat{j}$$

Chapter 13

Limits of Vector-valued Functions

The limit of a vector-valued function $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ as t approaches a is

$$\begin{aligned}\lim_{t \rightarrow a} \vec{r}(t) &= \langle \lim_{t \rightarrow a} x(t), \lim_{t \rightarrow a} y(t), \lim_{t \rightarrow a} z(t) \rangle \\ &= \lim_{t \rightarrow a} x(t)\hat{i} + \lim_{t \rightarrow a} y(t)\hat{j} + \lim_{t \rightarrow a} z(t)\hat{k}\end{aligned}$$

Example 1. Find $\lim_{t \rightarrow \pi} (t\hat{i} + \cos t\hat{j} + \sin t\hat{k})$.

$$\begin{aligned}\lim_{t \rightarrow \pi} (t\hat{i} + \cos t\hat{j} + \sin t\hat{k}) &= \lim_{t \rightarrow \pi} t\hat{i} + \lim_{t \rightarrow \pi} \cos t\hat{j} + \lim_{t \rightarrow \pi} \sin t\hat{k} \\ &= \pi\hat{i} + \cos \pi\hat{j} + \sin \pi\hat{k} \\ &= \pi\hat{i} - \hat{j}\end{aligned}$$

Example 2. Find $\lim_{t \rightarrow 0} (e^{6t}\hat{i} + \frac{\sin 2t}{2t}\hat{j} + e^{-5t}\hat{k})$.

$$\begin{aligned}\lim_{t \rightarrow 0} (e^{6t}\hat{i} + \frac{\sin 2t}{2t}\hat{j} + e^{-5t}\hat{k}) &= \lim_{t \rightarrow 0} e^{6t}\hat{i} + \lim_{t \rightarrow 0} \frac{\sin 2t}{2t}\hat{j} + \lim_{t \rightarrow 0} e^{-5t}\hat{k} \\ &= 1\hat{i} + 1\hat{j} + 1\hat{k} \\ &= \hat{i} + \hat{j} + \hat{k}\end{aligned}$$

Example 3. Find $\lim_{t \rightarrow \infty} (e^{-t}\hat{i} + \frac{1}{t^2 + 7}\hat{j} + e^{\arctan t}\hat{k})$.

$$\begin{aligned}\lim_{t \rightarrow \infty} (e^{-t}\hat{i} + \frac{1}{t^2 + 7}\hat{j} + e^{\arctan t}\hat{k}) &= \lim_{t \rightarrow \infty} e^{-t}\hat{i} + \lim_{t \rightarrow \infty} \frac{1}{t^2 + 7}\hat{j} + \lim_{t \rightarrow \infty} e^{\arctan t}\hat{k} \\ &= 0\hat{i} + 0\hat{j} + e^{\frac{\pi}{2}}\hat{k} \\ &= e^{\frac{\pi}{2}}\hat{k}\end{aligned}$$

Chapter 14

Derivatives and Integration of Vector-valued Functions

The derivative of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is

$$\vec{r}'(t) = x'(t)\hat{i} + y'(t)\hat{j} + z'(t)\hat{k}$$

The indefinite integral of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is

$$\int \vec{r}(t) dt = \int x(t)\hat{i} + \int y(t)\hat{j} + \int z(t)\hat{k} + C$$

The definite integral of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ from a to b is

$$\int_a^b \vec{r}(t) dt = \int_a^b x(t)\hat{i} + \int_a^b y(t)\hat{j} + \int_a^b z(t)\hat{k}$$

Since this topic is relatively straightforward, there will be no examples. :)

Chapter 15

Velocity, Speed and Acceleration

If $x(t)$, $y(t)$ and $z(t)$ are differentiable functions, then the velocity of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is

$$\vec{v}(t) = (\vec{r})'(t) = x'(t)\hat{i} + y'(t)\hat{j} + z'(t)\hat{k}$$

The speed of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is

$$|\vec{v}(t)| = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2}$$

The acceleration of a vector-valued function $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ is

$$\vec{a}(t) = (\vec{v})'(t) = x''(t)\hat{i} + y''(t)\hat{j} + z''(t)\hat{k}$$

The path of a projectile launched from an initial height h with initial speed v_0 at an angle of elevation θ is given by

$$\vec{r}(t) = v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \right] \hat{j}$$

Since this topic is relatively straightforward also, there will be no examples. :)

Selected Exercises

Source: *Larson Calculus 11th Ed. Exercise 12.3*

Projectile Motion

In Exercises 27-32, use the model for projectile motion, assuming there is no air resistance and $g = 9.8$ meters per second per second.

27. A baseball is hit from a height of 1 meter above the ground with an initial speed of 40 feet per second and at an angle of 22° above the horizontal. Find the maximum height reached by the baseball. Determine whether it will clear a 3-meters-high fence located 105 meters from home plate.

Solution. Given that $h = 1$, $v_0 = 40$, $\theta = 22^\circ$ and $g = 9.8$,

$$\begin{aligned}\vec{r}(t) &= v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \right] \hat{j} \\ &= 40 \cos 22^\circ t \hat{i} + \left[1 + (40 \sin 22^\circ)t - \frac{1}{2}(9.8)t^2 \right] \hat{j}\end{aligned}$$

The velocity vector is

$$\vec{v}(t) = (\vec{r})'(t) = 40 \cos 22^\circ \hat{i} + (40 \sin 22^\circ - 9.8t) \hat{j}$$

The maximum height is reached when the vertical component of the velocity is 0.

$$40 \sin 22^\circ - 9.8t = 0 \implies t = \frac{40 \sin 22^\circ}{9.8} \approx 1.53 \text{ seconds}$$

The maximum height is

$$\begin{aligned}y &= 1 + (40 \sin 22^\circ) \left(\frac{40 \sin 22^\circ}{9.8} \right) - \frac{1}{2}(9.8) \left(\frac{40 \sin 22^\circ}{9.8} \right)^2 \\ &\approx 12.46 \text{ meters}\end{aligned}$$

The ball is 105 meters from home plate when $x(t) = 105$.

$$40 \cos 22^\circ t = 105 \implies t = \frac{105}{40 \cos 22^\circ} \approx 2.83 \text{ seconds}$$

At this time, the height of the ball is

$$\begin{aligned}y &= 1 + (40 \sin 22^\circ) \left(\frac{105}{40 \cos 22^\circ} \right) - \frac{1}{2}(9.8) \left(\frac{105}{40 \cos 22^\circ} \right)^2 \\ &\approx 4.15 \text{ meters}\end{aligned}$$

Hence, the ball will clear the fence. ■

28. Determine the maximum height and range of a projectile fired at a height of 2 meters above the ground with an initial speed of 300 meters per second and at angle of 45° above the horizontal.

Solution. Given that $h = 2$, $v_0 = 300$, $\theta = 45^\circ$ and $g = 9.8$,

$$\begin{aligned}\vec{r}(t) &= v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \right] \hat{j} \\ &= 300 \cos 45^\circ t \hat{i} + \left[2 + (300 \sin 45^\circ)t - \frac{1}{2}(9.8)t^2 \right] \hat{j} \\ &= 150\sqrt{2}t \hat{i} + \left[2 + 150\sqrt{2}t - 4.9t^2 \right] \hat{j}\end{aligned}$$

The velocity vector is

$$\begin{aligned}\vec{v}(t) &= (\vec{r})'(t) \\ &= 150\sqrt{2}\hat{i} + (150\sqrt{2} - 9.8t)\hat{j}\end{aligned}$$

The maximum height is reached when the vertical component of the velocity is 0.

$$\begin{aligned}150\sqrt{2} - 9.8t &= 0 \\ t &= \frac{150\sqrt{2}}{9.8} \\ &\approx 21.65 \text{ seconds}\end{aligned}$$

The maximum height is

$$\begin{aligned}y &= 2 + (150\sqrt{2})\left(\frac{150\sqrt{2}}{9.8}\right) - 4.9\left(\frac{150\sqrt{2}}{9.8}\right)^2 \\ &\approx 2,297.92 \text{ meters}\end{aligned}$$

The projectile hits the ground when $y(t) = 0$.

$$\begin{aligned}2 + (150\sqrt{2})t - 4.9t^2 &= 0 \\ t &= \frac{-150\sqrt{2} - \sqrt{150^2(2) - 4(-4.9)(2)}}{2(-4.9)} \\ &\approx 43.302 \text{ seconds}\end{aligned}$$

Hence, the range is

$$x = 150\sqrt{2} \left(\frac{-150\sqrt{2} - \sqrt{150^2(2) - 4(-4.9)(2)}}{2(-4.9)} \right) \approx 9185.67 \text{ meters}$$

■

29. A baseball, hit 1 meter above the ground, leaves the bat at an angle of 45° and is caught by an outfielder 1 feet above the ground and 100 feet from home plate. What is the initial speed of the ball, and how high does it rise?

Solution. Given that $h = 1$, $\theta = 45^\circ$ and $g = 9.8$,

$$\begin{aligned}\vec{r}(t) &= v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta) t - \frac{1}{2} g t^2 \right] \hat{j} \\ &= v_0 \cos 45^\circ t \hat{i} + \left[1 + (v_0 \sin 45^\circ) t - \frac{1}{2} (9.8) t^2 \right] \hat{j} \\ &= \frac{v_0}{\sqrt{2}} t \hat{i} + \left[1 + \frac{v_0}{\sqrt{2}} t - 4.9 t^2 \right] \hat{j}\end{aligned}$$

When the ball is caught, $x(t) = 100$ and $y(t) = 1$.

$$\frac{v_0}{\sqrt{2}} t = 100 \implies t = \frac{100\sqrt{2}}{v_0} \dots (1)$$

$$1 + \frac{v_0}{\sqrt{2}} t - 4.9 t^2 = 1$$

$$4.9 t^2 - \frac{v_0}{\sqrt{2}} t = 0$$

$$t(4.9 t - \frac{v_0}{\sqrt{2}}) = 0$$

$$t = \frac{v_0}{4.9\sqrt{2}} \dots (2)$$

Equating (1) and (2),

$$\frac{100\sqrt{2}}{v_0} = \frac{v_0}{4.9\sqrt{2}}$$

$$v_0^2 = 980$$

$$v_0 = 14\sqrt{5} \text{ meters per second } (v_0 > 0)$$

Hence,

$$\begin{aligned}\vec{r}(t) &= \frac{14\sqrt{5}}{\sqrt{2}} t \hat{i} + \left[1 + \frac{14\sqrt{5}}{\sqrt{2}} t - 4.9 t^2 \right] \hat{j} \\ &= 7\sqrt{10} t \hat{i} + \left[1 + 7\sqrt{10} t - 4.9 t^2 \right] \hat{j}\end{aligned}$$

The maximum height is reached when the vertical component of the velocity is 0.

$$7\sqrt{10} - 9.8 t = 0$$

$$t = \frac{7\sqrt{10}}{9.8}$$

$$\approx 2.26 \text{ seconds}$$

The maximum height is

$$\begin{aligned} y &= 1 + (7\sqrt{10}) \left(\frac{7\sqrt{10}}{9.8} \right) - 4.9 \left(\frac{7\sqrt{10}}{9.8} \right)^2 \\ &= 26 \text{ meters} \end{aligned}$$

■

30. A baseball player at second base throws a ball 28 meters to the player at first base. The ball is released at a point 1.5 meters above the ground with an initial speed of 80 kilometres per hour and at an angle of 15° above the horizontal. At what height does the player at first base catch the ball?

Solution. Given that $h = 1.5$, $v_0 = 80\text{km/h} = 80 \times \frac{1000}{3600} = 22\frac{2}{9}\text{m/s}$, $\theta = 15^\circ$ and $g = 9.8$,

$$\begin{aligned} \vec{r}(t) &= v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta) t - \frac{1}{2} g t^2 \right] \hat{j} \\ &= 22\frac{2}{9} \cos 15^\circ t \hat{i} + \left[1.5 + (22\frac{2}{9} \sin 15^\circ) t - \frac{1}{2} (9.8) t^2 \right] \hat{j} \\ &= 22\frac{2}{9} \cos 15^\circ t \hat{i} + \left[1.5 + 22\frac{2}{9} \sin 15^\circ t - 4.9 t^2 \right] \hat{j} \end{aligned}$$

When the ball is caught, $x(t) = 28$

$$22\frac{2}{9} \cos 15^\circ t = 28 \implies t = \frac{28}{22\frac{2}{9} \cos 15^\circ} \approx 1.305 \text{ seconds}$$

The height of the ball is

$$\begin{aligned} y &= 1.5 + 22\frac{2}{9} \sin 15^\circ \left(\frac{28}{22\frac{2}{9} \cos 15^\circ} \right) - 4.9 \left(\frac{28}{22\frac{2}{9} \cos 15^\circ} \right)^2 \\ &\approx 0.66 \text{ meters} \end{aligned}$$

■

31. Eliminate the parameter t from the position vector for the motion of a projectile to show that the rectangular equation is

$$y = -\frac{g \sec^2 \theta}{2v_0^2} x^2 + (\tan \theta)x + h$$

Solution.

$$\vec{r}(t) = v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \right] \hat{j}$$

Let $x = v_0 \cos \theta t$ and $y = h + (v_0 \sin \theta)t - \frac{1}{2}gt^2$.

$$\begin{aligned} x &= v_0 \cos \theta t \\ t &= \frac{x}{v_0 \cos \theta} \\ y &= h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \\ &= h + (v_0 \sin \theta) \left(\frac{x}{v_0 \cos \theta} \right) - \frac{1}{2}g \left(\frac{x}{v_0 \cos \theta} \right)^2 \\ &= h + \frac{v_0 \sin \theta}{v_0 \cos \theta} x - \frac{1}{2} \frac{g}{v_0^2 \cos^2 \theta} x^2 \\ &= h + (\tan \theta)x - \frac{1}{2} \frac{g}{v_0^2 \cos^2 \theta} x^2 \\ &= -\frac{g \sec^2 \theta}{2v_0^2} x^2 + (\tan \theta)x + h \end{aligned}$$

■

32. The path of a ball is given by the rectangular equation

$$y = x - 0.0245x^2.$$

Use the result of Exercise 31 to find the position vector. Then find the speed and direction of the ball at the point at which it has travelled 15 meters horizontally.

Solution. Comparing the equation with the equation in Exercise 31,

$$\begin{aligned}
 \tan \theta &= 1 \\
 \theta &= 45^\circ \\
 -\frac{g \sec^2 \theta}{2v_0^2} &= -0.0245 \\
 \frac{9.8}{v_0^2} &= 0.0245 \\
 v_0^2 &= 400 \\
 v_0 &= 20 \text{ meters per second } (v_0 > 0)
 \end{aligned}$$

Hence, the position vector is

$$\begin{aligned}
 \vec{r}(t) &= v_0 \cos \theta t \hat{i} + \left[h + (v_0 \sin \theta)t - \frac{1}{2}gt^2 \right] \hat{j} \\
 &= 20 \cos 45^\circ t \hat{i} + \left[0 + (20 \sin 45^\circ)t - \frac{1}{2}(9.8)t^2 \right] \hat{j} \\
 &= 10\sqrt{2}t \hat{i} + \left[10\sqrt{2}t - 4.9t^2 \right] \hat{j}
 \end{aligned}$$

The time taken to travel 15 meters horizontally is

$$10\sqrt{2}t = 15 \implies t = \frac{3\sqrt{2}}{2} \approx 2.12 \text{ seconds}$$

The speed vector is

$$\vec{v}(t) = (\vec{r})'(t) = 10\sqrt{2}\hat{i} + (10\sqrt{2} - 9.8t)\hat{j}$$

The direction of the ball is

$$\begin{aligned}
 \vec{v}\left(\frac{3\sqrt{2}}{2}\right) &= 10\sqrt{2}\hat{i} + (10\sqrt{2} - 9.8\left(\frac{3\sqrt{2}}{2}\right))\hat{j} \\
 &= 10\sqrt{2}\hat{i} - 14.7\sqrt{2}\hat{j}
 \end{aligned}$$

Hence, the speed of the ball is

$$\begin{aligned}
 \left\| \vec{v}\left(\frac{3\sqrt{2}}{2}\right) \right\| &= \sqrt{(10\sqrt{2})^2 + (-14.7\sqrt{2})^2} \\
 &= 25.14 \text{ meters per second}
 \end{aligned}$$

■

Chapter 16

Tangent Vectors and Normal Vectors

Let C be a smooth curve represented by $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ on an open interval I , the unit tangent vector at $t = a$ is a vector that is tangent to the curve at $t = a$ and has a magnitude of 1. The unit tangent vector is given by

$$\vec{T}(t) = \frac{(\vec{r})'(t)}{\|(\vec{r})'(t)\|} = \frac{x'(t)\hat{i} + y'(t)\hat{j} + z'(t)\hat{k}}{\sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2}}, \quad (\vec{r})'(t) \neq \vec{0}$$