Waterlogging caused by drainage issues

"WATERLOGGING DUE TO DRAINAGE" *Generated by LeapSpace*

Waterlogging Due to Drainage: Quick Insights

Key Findings Table

Key finding	Main settings/populations	Evidence
Waterlogging due to drainage occurs when water inputs exceed the capacity or functioning of surface or subsurface drainage systems.	Agricultural land, irrigated regions, urban and peri-urban drainage systems	[1] [2] [3] [4] [5] [6] [7]
The main hydrologic signature is prolonged near-surface saturation, shallow water tables, and saturation-excess ponding rather than infiltration failure alone.	Poorly drained fields, drained agricultural plots, restrictive-soil landscapes	[8] [9] [10] [11] [12] [13]
Soil type and landscape position strongly control whether drainage failure becomes chronic waterlogging.	Clay soils, shallow or restrictive profiles, depressions, convergent slopes	[12] [13] [14] [15] [16] [17]
Maintenance failures such as blockage, siltation, weed growth, and pump or outlet problems can sharply reduce drainage performance.	Irrigation schemes, city drains, stormwater inlets, subsurface drainage networks	[6] [18] [19] [20] [21]
Crop damage arises mainly through root-zone oxygen stress, nutrient disruption, and higher disease risk, causing substantial yield loss.	Maize, wheat, cotton, other major crops on waterlogged soils	[22] [23] [24] [25] [26] [27]
Drainage improvements usually reduce saturated days and improve trafficability and yield, but the best response is often an integrated package rather than a single intervention.	Waterlogged farmland, saline/waterlogged land, slow-permeability soils	[28] [29] [30] [31] [32] [33]
Faster drainage is not always optimal: free drainage can increase water and nutrient losses, while controlled drainage can conserve water and nutrients but may worsen wetness under already wet conditions.	Humid and irrigated agricultural systems	[34] [35] [36] [37] [38] [39]

Key finding	Main settings/populations	Evidence
Climate extremes increasingly overwhelm drainage systems, amplifying waterlogging risk.	Urban systems, peri-urban fields, compound flood settings	[7] [40] [41] [42] [43] [44]

Direct Answer

Waterlogging due to drainage is the condition in which rainfall, irrigation recharge, seepage, or runoff cannot be removed fast enough because the drainage system is inadequate, blocked, poorly maintained, undersized, or poorly matched to local soils and topography. The immediate result is shallow groundwater, ponding, and prolonged soil saturation; the practical consequences are reduced land usability, crop root stress, nutrient disruption, disease risk, lower yields, and in urban settings longer-lasting inundation and infrastructure problems [1] [3] [7] [22] [23] [45] [46] [47].

A practical way to assess the problem is to test four linked factors: **water inputs**, **soil transmissivity**, **drainage-network capacity**, and **maintenance condition**. Taken together, the evidence suggests that waterlogging is most likely where high inputs coincide with low-permeability or restrictive soils, low-lying topography, and drains whose spacing, depth, outlets, or upkeep are inadequate for present conditions [14] [15] [17] [21] [44] [48] [49].

Confidence level: High - The evidence is consistent across hydrologic, agronomic, engineering, and urban drainage studies [1] [6] [8] [22] [30] [40].

Study Scope

This report synthesizes field studies, drainage design studies, calibrated hydrologic modeling, irrigation and urban drainage case studies, and agronomic experiments spanning roughly 1990-2025, with strongest coverage in agricultural drainage and moderate coverage in urban/peri-urban drainage failure under storm conditions [8] [19] [27] [40] [50] [51] [52].

Assumptions & Limitations

- The evidence strongly explains mechanisms and design effects, but there are fewer direct before-after field studies of real blockage or maintenance failure sequences than of modeled drainage scenarios [8] [20] [21] [50] [53].
- No single universal threshold defines “inadequate drainage”; critical limits vary by soil hydraulic conductivity, topography, rainfall intensity, outlet capacity, crop, and management [7] [17] [49].
- The supplied evidence does not provide global prevalence estimates or a region-by-region ranking of drainage-caused waterlogging [1] [42].

Suggested Further Research

- Direct before-after measurements of blockage, siltation, pump failure, and weed growth on groundwater depth, runoff, and ponding duration [20] [21].

- Maintenance thresholds that identify how much obstruction causes major loss of drainage function [19] [54].
- Shared catchment studies linking urban overflow and agricultural waterlogging in peri-urban landscapes [42] [55].
- Crop-specific damage functions based on duration, depth, and timing of drainage-caused waterlogging [26] [27].
- Evaluation of hybrid mitigation packages combining grading, subsurface drainage, controlled drainage, pumping, and soil improvement under climate extremes [28] [29] [30] [44].

Full Report: Waterlogging Due to Drainage Failure

Foundational Context

Waterlogging is not simply “too much rain.” In the drainage literature, it is better understood as a **soil-water imbalance** in which incoming water exceeds the landscape’s ability to transmit, store, or remove that water through surface and subsurface pathways [1] [2] [3]. Drainage systems exist to control that imbalance by lowering water tables, conveying excess water, and limiting the duration of root-zone saturation [2] [56] [57].

Three linked concepts are central:

1. **Water inputs**: precipitation, irrigation recharge, canal seepage, wastewater/stormwater overflow, and upslope runoff all add water to soils or drainage networks [3] [55] [58].
2. **Landscape transmission and storage**: soil hydraulic conductivity, restrictive layers, soil depth, and local topography determine whether water infiltrates, perches, moves laterally, or ponds [12] [13] [14] [49].
3. **Drainage removal capacity**: drain spacing, drain depth, outlet size, slope, roughness, pumping capacity, and maintenance condition control how fast excess water can leave the site [4] [5] [21] [48].

When any part of this chain fails, water tables rise, drainage discharge becomes insufficient, and ponding persists [6] [10] [11]. In agriculture this reduces aeration and rooting; in urban and peri-urban environments it prolongs surface inundation and worsens infrastructure impacts [40] [45] [47].

Confidence level: High - Multiple source types converge on the same conceptual model [1] [2] [4] [14].

1. Why drainage-related waterlogging occurs

The literature consistently treats waterlogging as a **drainage-capacity and drainage-function problem**. Excess rainfall alone does not automatically produce chronic waterlogging; persistent wetness emerges when water inputs exceed what the drainage system and soil profile can transmit or evacuate [1] [58] [59] [60]. This can happen through several pathways:

- **Undersized drainage design**, including insufficient drain spacing, shallow drains, weak outlet capacity, or low drainage coefficients [4] [5] [48].
- **Poor maintenance**, including weed growth, sediment deposition, bio-clogging, irregular cleaning, and blocked inlets or pipes [6] [18] [19] [21].
- **External recharge**, such as irrigation return flow, canal seepage, or perched groundwater sustained by restrictive layers [3] [58] [61].
- **Hydroclimatic overload**, where intense storms or compound flood conditions exceed the design assumptions of the network [40] [42] [44].

Taken together, these findings suggest that drainage-related waterlogging is usually not a single-cause problem; it arises from interaction between water supply, soil resistance, and network failure or underperformance [7] [14] [20].

2. Hydrological mechanisms of drainage-related waterlogging

2.1 Soil water balance changes under drainage failure or inadequacy

Field and modeling studies show that drainage affects the partitioning of water among runoff, infiltration, subsurface drainage, groundwater storage, and deep seepage [11] [53] [62]. A calibrated DRAINMOD study reported surface runoff equal to 1-10% of annual precipitation and subsurface drainage equal to 26-45%, while ponding events were identified as saturation-excess rather than infiltration-excess [8]. Another field study found high partitioning to surface drainage and tile drainage, yet also showed that a large share of infiltrated water was not captured by tiles, demonstrating that drainage performance is only one part of field water balance behavior [9].

This matters because waterlogging often develops not from rainfall failing to enter the soil, but from water entering and then remaining too close to the surface for too long [8] [11]. In that sense, drainage failure changes both **how much water leaves** and **how long water remains** in root-zone or near-surface storage [10] [50].

2.2 Groundwater table response and ponding

Subsurface drainage is designed to control water-table depth, but its performance depends on system geometry and the surrounding soil [2] [56] [57]. Modeling studies show that design changes alter water-table fluctuations and surface runoff; for example, shallow 75-cm drains increased surface runoff in DRAINMOD simulations [10]. Studies also indicate that differences in drainage management can be represented as changes in infiltration, runoff, drainage outflow, deep seepage, and water-table dynamics [50] [53].

The strongest recurring hydrologic signature is **prolonged shallow water tables and saturation-excess ponding** [8] [10] [12]. In restrictive landscapes, excess water-table metrics directly quantify how long water remains above a drainage threshold, linking poor drainage to the duration—not just occurrence—of wet conditions [13].

Confidence level: High - Process evidence is strong and consistent, though much of it comes from calibrated modeling rather than observed blockage events [8] [50] [53].

3. Drainage design, capacity, and maintenance failures

3.1 Design controls on drainage performance

Drainage performance is strongly governed by **drain spacing, depth, hydraulic conductivity, drainage coefficient, and outlet capacity** [4] [48] [51]. One DRAINMOD study found that maintaining water tables around 1.0-1.5 m required substantially narrower spacing in clay than in clay loam, showing how soil permeability shifts design requirements [48]. Hydraulic conductivity is especially critical because low or vertically variable conductivity can leave water tables shallower than intended and increase both crop and salinity risk [49] [63].

Drainage coefficients define the hydraulic removal capacity at the outlet and must be distinguished from general drainage intensity [4]. Even when in-field drainage exists, outlet works, slope, roughness, and pumping capacity can constrain actual discharge [5] [64]. This means that a field may appear “drained” on paper but still remain vulnerable if downstream conveyance is inadequate [4] [7].

3.2 Blockage, siltation, and maintenance neglect

Maintenance failure is a recurring cause of real-world waterlogging. In one irrigation scheme, weed and aquatic grass blockage rendered both surface and subsurface drains non-functional, contributing to waterlogging [18]. In Indian cities, blocked drains, heavy siltation, and irregular cleaning were reported to reduce hydraulic capacity and prolong ponding [6]. In flat urban catchments, blocked inlets triggered flooding even before sewer capacity was exceeded, linking many flood events directly to insufficient inlet maintenance [19].

Agricultural systems show a similar pattern. Neglected ditch and tile cleaning gradually reduces conveyance over time [20], while sediment deposition and bio-clogging progressively reduce subsurface drain conductance [21]. Pipe siltation also reduces effective cross-sectional area and redistributes hydraulic loads, worsening inundation under storm events [54] [65].

Taken together, these studies imply that many drainage failures are operational rather than purely structural: maintenance can determine whether a designed system remains functional [18] [21].

Confidence level: High - Evidence spans both agricultural and urban cases and identifies specific failure modes [6] [18] [19] [21].

4. Soil, topography, and climate controls

4.1 Soil and profile controls

Drainage inadequacy does not produce the same outcome everywhere. Clay-rich soils, shallow topsoils, restrictive subsurface layers, and vertically variable conductivity all promote perched water tables and slow drainage [13] [14] [16] [63]. Heavy or restrictive soils may require closer drains or additional soil improvement to achieve target performance [17].

In claypan and similarly restrictive profiles, water status is better described by overall profile wetness and stratification than by a single soil depth, because post-rain redistribution remains shallow and persistent [66]. This helps explain why some fields remain waterlogged long after rainfall has ended [13].

4.2 Topographic and landscape controls

Topography controls where water accumulates and how long it remains. Bowl-shaped, low-lying, and convergent positions stay wetter; convex or upslope positions dry more quickly [12] [15]. Small depressions can enlarge saturated zones and sustain runoff or waterlogged patches [12] [67]. Slope position may matter as much as slope orientation because lateral redistribution and soil depth vary across hillslopes [68] [69].

This means drainage design should be landscape-specific: a uniform spacing or outlet assumption may underperform in depressional terrain or in areas with strong convergence [15] [48].

4.3 Rainfall extremes and compound flooding

Storm intensification increasingly overwhelms drainage systems [40] [41] [70]. Urban studies report more overflow points, ponding, and flood-prone locations under future rainfall extremes [40] [71]. Seasonal peaks and midsummer flood-season rainfall are also associated with higher waterlogging risk [72]. In peri-urban agricultural settings, sudden heavy rainfall can cause mixed stormwater and wastewater/irrigation drains to overflow into fields, producing long-duration inundation and crop damage [55].

Compound events are especially serious: when intense precipitation coincides with upstream flooding and drainage constraints, inundation area and depth can far exceed rainfall-only scenarios [42]. This is a strong indication that historical design standards may be increasingly misaligned with present extremes [43] [44].

Confidence level: High - Strong cross-source agreement on soil and topographic mediation; climate-risk evidence is also strong, especially for overload of drainage capacity [14] [15] [40] [42].

5. Agricultural impacts of waterlogging due to drainage

5.1 Root stress and nutrient disruption

Waterlogging fills soil pores with water and reduces oxygen availability, causing root-zone hypoxia or anoxia [23] [73] [74]. This suppresses root respiration, ATP production, root growth, and overall root function [23] [75]. Nutrient uptake then declines, especially for nitrogen, because waterlogging disrupts N availability, absorption, assimilation, and broader carbon-nitrogen metabolism [24] [76] [77] [78].

Other nutrient imbalances also occur. Studies report effects on K, P, Ca, and Na balance, along with leaching and denitrification losses that further reduce crop performance [76] [79] [80] [81]. Waterlogging also raises disease and pest susceptibility, including root rot and harmful microbial conditions [25] [75] [82].

5.2 Yield loss

Yield penalties are substantial and consistent across crops. Early-stage waterlogging reduced maize yield by 58.8-69.8% in reported studies [83]. Cotton showed large yield losses and was identified in a meta-analysis as having among the highest row-crop reductions [26]. Winter wheat field studies also reported reduced dry matter and final yield under winter waterlogging [27].

Taken together, these findings suggest that duration and timing of waterlogging are critical: early or prolonged saturation during sensitive growth periods can produce severe losses even if plants survive [27] [83].

Confidence level: High - Mechanisms and yield effects are repeatedly documented across crop studies [22] [23] [24] [27].

6. Mitigation and management responses

6.1 Drainage and field management responses

Drainage improvement generally reduces waterlogging impacts. Subsurface drainage increased rice and rapeseed yields in waterlogged settings, and simulations showed wheat yield recovery from 65% to 78-82% of potential after drainage improvement [30] [32]. On drained saline land, yields increased by 18.8-27.6% while undrained areas declined [33].

Drainage also improves field usability. Studies found subsurface drainage made fields trafficable 2-4 weeks earlier in silty clay loam and up to 5 weeks earlier in muck, while long-term simulations linked drainage to fewer waterlogged days and less planting delay [31] [32]. Narrower drain spacing or optimized design reduced groundwater table depth and soil water content [30].

Raised beds, ridge tillage, and similar management practices can partially restore aeration and nutrient uptake where full drainage reconstruction is not feasible [23] [27] [77].

6.2 Why integrated solutions often work best

Evidence favors combinations of interventions rather than one measure alone, especially where waterlogging co-occurs with salinity, irrigation recharge, or slowly permeable soils [28] [29] [84]. Integrated packages may include:

- subsurface drainage,
- surface grading or surface drains,
- raised beds or ridge systems,
- pumping or outlet upgrades,
- soil improvement where conductivity is limiting [17] [29] [30].

This pattern implies that chronic waterlogging usually reflects multiple bottlenecks, so removing only one may leave the system underperforming [21] [28] [49].

Confidence level: High - Consistent evidence supports drainage improvement, with the strongest gains in complex settings coming from combined approaches [28] [29] [30] [85].

7. Controlled drainage tradeoffs

Controlled drainage (CD) differs from free drainage by intentionally regulating outflow. The evidence shows a real tradeoff.

Benefits

CD consistently reduced drainage outflow and nutrient losses compared with free drainage, including lower discharge volumes and reduced nitrate or phosphorus export [35] [36] [52] [86] [87] [88] [89]. In irrigated and arid settings, it also helped maintain root-zone groundwater availability and improved later-season water supply and water-use efficiency [88] [90] [91]. Some studies found improved crop N uptake and maintained or higher yields, even with lower fertilizer rates [90] [92] [93].

Risks and limits

Yield response to CD depends on weather, crop, and field condition. Yield gains are more common in dry years or drought-prone settings, while wet seasons may show no benefit or even lower yields; paddy rice also showed yield declines in a meta-analysis [36] [37] [94] [95] [96]. CD can also create partially waterlogged conditions when drainage is very shallow or regulation is prolonged under already wet conditions [37] [38] [39] [97].

So the evidence does not support “more retained water is always better.” Rather, optimal drainage is context-specific: enough removal to prevent waterlogging, but not so much that water and nutrients are unnecessarily lost [10] [34] [39].

Confidence level: High - The tradeoff is directly documented across comparative CD studies [35] [36] [37] [39].

8. Practical diagnostic framework

A practical evidence-based diagnostic sequence for drainage-related waterlogging is:

1. **Quantify water inputs**: rainfall peaks, irrigation recharge, seepage, overflow, upslope contributions [3] [55] [58].
2. **Check saturation pattern**: depth to water table, ponding duration, whether wetness is localized in depressions or widespread [11] [13] [15].
3. **Evaluate soils**: hydraulic conductivity, restrictive layers, shallow profiles, compaction [14] [16] [49].
4. **Inspect drainage design**: spacing, depth, drainage coefficient, outlet and pump capacity [4] [5] [48] [51].
5. **Inspect maintenance**: weed blockage, inlet obstruction, siltation, bio-clogging, irregular cleaning [6] [18] [19] [21].
6. **Match intervention to failure mode**: redesign if capacity is inadequate; rehabilitate if obstruction dominates; integrate surface and subsurface measures if soils/topography create persistent saturation [28] [29] [30].

This framework is an inference drawn from the combined evidence base rather than a single published protocol [21] [32] [48].
Confidence level: Medium - Strongly supported by synthesis across studies, but not presented as a standardized framework in one source [1] [18] [28].

Conclusion

Waterlogging due to drainage is best understood as the outcome of **excess water meeting insufficient or degraded drainage performance** [1] [3] [4]. Its clearest hydrologic expressions are shallow water tables, prolonged saturation, and saturation-excess ponding [8] [10] [12]. Whether the problem becomes chronic depends heavily on soil permeability, restrictive layers, topographic convergence, and the adequacy and maintenance of drainage networks [14] [15] [21]. In agriculture, the main damage pathway is root oxygen stress plus nutrient disruption, leading to disease risk and yield loss [23] [24] [25] [27]. Drainage improvements usually help, but the best solutions are site-specific and often integrated; aggressive free drainage may increase losses, while controlled drainage can conserve water and nutrients yet worsen wetness under already saturated conditions [30] [34] [35] [39]. Climate extremes make both design adequacy and maintenance more critical than before [40] [42] [44].

Overall confidence level: High - The central mechanisms, impacts, and major mitigation principles are strongly and repeatedly supported, though direct before-after blockage evidence remains thinner than process and modeling evidence [7] [8] [20] [21].

References

1. P Pais, I.P., Moreira, R., Semedo, J.N., (...), Scotti-Campos, P. (2023) Wheat Crop under Waterlogging: Potential Soil and Plant Effects. *Plants*. https://doi.org/10.3390/plants12010149
2. (1982) Design of subsurface drains. *Developments in Water Science*. https://doi.org/10.1016/S0167-5648(08)70578-1
3. Deng, C., Bailey, R.T. (2020) Assessing causes and identifying solutions for high groundwater levels in a highly managed irrigated region. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2020.106329
4. Wayne Skaggs, R. (2016) Quantifying subsurface drainage rates for research sites. *10th International Drainage Symposium 2016*. https://doi.org/10.13031/ids.20162493719
5. Wayne Skaggs, R. (2017) Coefficients for quantifying subsurface drainage rates. *Applied Engineering in Agriculture*. https://doi.org/10.13031/aea.12302
6. Singh, H., Nielsen, M., Greatrex, H. (2023) Causes, impacts, and mitigation strategies of urban pluvial floods in India: A systematic review. *International Journal of Disaster Risk Reduction*. https://doi.org/10.1016/j.ijdrr.2023.103751
7. Liwanag, F., Mostrales, D., Sanchez, K., (...), Ignacio, M.T. (2018) GIS-based estimation of catchment basin parameters and maximum discharge calculation using rational method of Luinab Catchment in Iligan City. *Philippine Journal of Science*. https://www.scopus.com/pages/publications/85066398463
8. Saadat, S., Frankenberger, J., Bowling, L., Srinivasulu, S. (2020) Evaluation of surface ponding and runoff generation in a seasonally frozen drained agricultural field. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2020.124985

9. Gaillot, A., Delbart, C., Salvador-Blanes, S., (...), Cerdan, O. (2023) Analysis of seasonal variation in the hydrological behaviour of a field combining surface and tile drainage. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2023.108329
10. Ghane, E., Askar, M.H. (2021) Predicting the effect of drain depth on profitability and hydrology of subsurface drainage systems across the eastern USA. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2021.107072
11. Golmohammadi, G., Prasher, S.O., Madani, A., (...), Youssef, M.A. (2016) SWATDRAIN, a new model to simulate the hydrology of agricultural Lands, model development and evaluation. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2015.11.003
12. Van Tol, J.J., Roux, P.A.L.L., Hensley, M. (2010) Soil indicators of hillslope hydrology in the bedford catchments, South Africa. *South African Journal of Plant and Soil*. https://doi.org/10.1080/02571862.2010.10639993
13. Nibler, J.C. (1975) Drainage: Numerically evaluated for humid regions.. https://www.scopus.com/pages/publications/0016449239
14. Cox, J.W., McFarlane, D.J. (1995) The causes of waterlogging in shallow soils and their drainage in southwestern Australia. *Journal of Hydrology*. https://doi.org/10.1016/0022-1694(94)02614-H
15. Reid, I., Parkinson, R.J. (1984) The nature of the tile-drain outfall hydrograph in heavy clay soils. *Journal of Hydrology*. https://doi.org/10.1016/0022-1694(84)90086-6
16. Swarowsky, A., Dahlgren, R.A., O'Geen, A.T. (2012) Linking subsurface lateral flowpath activity with streamflow characteristics in a semiarid headwater catchment. *Soil Science Society of America Journal*. https://doi.org/10.2136/sssaj2011.0061
17. Trafford, B.D. (1975) Improving the design of practical field drainage. *Soil Science*. https://doi.org/10.1097/00010694-197505000-00002
18. Chikwanda, O., Chisadza, B., Mpala, C. (2025) An assessment of the drainage system and waterlogging at the Silalabuhwa irrigation scheme. *Physics and Chemistry of the Earth*. https://doi.org/10.1016/j.pce.2024.103837
19. Wang, L., Li, J., Deng, Z., Gong, Q. (2021) Spotting strategic storm drain inlets in flat urban catchments. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2021.126504
20. Schleyer, C. (2017) Avoiding Floods in Spring and Droughts in Summer-Water Regulation Strategies in Germany and Poland. *Competition for Water Resources: Experiences and Management Approaches in the US and Europe*. https://doi.org/10.1016/B978-0-12-803237-4.00021-5
21. https://www.sciencedirect.com/science/article/pii/S236362112600096X
22. Huawei, L., Shouwei, H., Weibao, Y., (...), Fahong, W. (2022) Crop exposure to waterlogging stress: responses to physiological, biochemical, and molecular levels. *Sustainable Crop Productivity and Quality under Climate Change: Responses of Crop Plants to Climate Change*. https://doi.org/10.1016/B978-0-323-85449-8.00007-5

23. Li, T., Cui, L., Filipović, V., (...), Dang, Y.P. (2025) From soil health to agricultural productivity: The critical role of soil constraint management. *Catena*. https://doi.org/10.1016/j.catena.2025.108776

24. Kashyap, Suryakanta, Biswal, Bisworanjita, Bhakuni, Kanika, (...), Kumar, Rakesh (2024) Amelioration of abiotic stresses in forage crop production using microbial stimulants: an overview. *Microbial Biostimulants for Plant Growth and Abiotic Stress Amelioration*. https://doi.org/10.1016/B978-0-443-13318-3.00019-7

25. Gantait, Saikat, Subrahmanyeswari, Tsama, Kumar, Gunti Vinod, (...), Suprasanna, Penna (2024) Advances in understanding and engineering plant root system architecture to alleviate abiotic stress. *Current Omics Advancement in Plant Abiotic Stress Biology*. https://doi.org/10.1016/B978-0-443-21625-1.00011-7

26. Wu, Lianhai, Blackwell, Martin, Dunham, Sarah, (...), McGrath, Steve P. (2019) Simulation of Phosphorus Chemistry, Uptake and Utilisation by Winter Wheat. *Plants*. https://doi.org/10.3390/plants8100404

27. Beegum, Sahila, Truong, Van, Bheemanahalli, Raju, (...), Reddy, Kambham Raja (2023) Developing functional relationships between waterlogging and cotton growth and physiology-towards waterlogging modeling. *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2023.1174682

28. Gupta, S.K. (2016) Integrated drainage solutions for waterlogged saline lands. *Innovative Saline Agriculture*. https://doi.org/10.1007/978-81-322-2770-0_9

29. Singh, A. (2018) Salinization of agricultural lands due to poor drainage: A viewpoint. *Ecological Indicators*. https://doi.org/10.1016/j.ecolind.2018.07.037

30. Qi, B., Zheng, X., Junejo, A.R., (...), Liu, H. (2026) Synergistic water management enhances rice-rapeseed rotation yields in waterlogged areas: field and modeling evidence. *Irrigation Science*. https://doi.org/10.1007/s00271-026-01114-4

31. Paul, C.L., de Vries, J. (1983) Soil trafficability in spring 2. Prediction and the effect of subsurface drainage.. *Canadian Journal of Soil Science*. https://doi.org/10.4141/cjss83-003

32. Jin, C.-X., Sands, G.R. (2004) Simulation of spring wheat response to subsurface drainage in northwest Minnesota. *ASAE Annual International Meeting 2004*. https://www.scopus.com/pages/publications/30044446838

33. Sharma, D.P., Gupta, S.K. (2006) Subsurface drainage for reversing degradation of waterlogged saline lands. *Land Degradation and Development*. https://doi.org/10.1002/ldr.737

34. Luo, W., Sands, G.R., Youssef, M., (...), Canelón, D. (2010) Modeling the impact of alternative drainage practices in the northern Corn-belt with DRAINMOD-NII. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2009.10.009

35. Ayars, J.E., Christen, E.W., Hornbuckle, J.W. (2006) Controlled drainage for improved water management in arid regions irrigated agriculture. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2006.07.004

36. Wang, Z., Shao, G., Lu, J., (...), Ding, J. (2020) Effects of controlled drainage on crop yield, drainage water quantity and quality: A meta-analysis. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2020.106253

37. Youssef, M.A., Strock, J., Bagheri, E., (...), Pease, L. (2023) Impact of controlled drainage on corn yield under varying precipitation patterns: A synthesis of studies across the U.S. Midwest and Southeast. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2022.107993

38. Deichmann, M.M., Neumann Andersen, M.N., Thomsen, I.K., Borgesen, C.D. (2019) Impacts of controlled drainage during winter on the physiology and yield of winter wheat in Denmark. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2019.01.013

39. Fouss, James L., Rogers, James S. (1992) ET from shallow groundwater maintained by controlled-drainage/subirrigation system. *Irrigation and Drainage: Saving a Threatened Resource - In Search of Solutions, Proceedings of the Irrigation and Drainage Sessions at Water Forum '92*. https://www.scopus.com/pages/publications/0027035678

40. Lameche, E.-K., Boutaghane, H., Saber, M., (...), Sumi, T. (2026) Climate-adaptive urban drainage modeling: Assessing future flood hazards in Algiers, Algeria. *Urban Climate*. https://doi.org/10.1016/j.uclim.2026.102843

41. Wasko, C., Nathan, R., Stein, L., O'Shea, D. (2021) Evidence of shorter more extreme rainfalls and increased flood variability under climate change. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2021.126994

42. Zhao, Z., Huo, A., Liu, Q., (...), Elbeltagi, A. (2024) Assessment of urban inundation and prediction of combined flood disaster in the middle reaches of Yellow river basin under extreme precipitation. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2024.131707

43. Andrews, T.D., Felzer, B.S. (2015) Very-heavy precipitation in the greater New York City region and widespread drought alleviation tied to Western US agriculture. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0144416

44. Ragno, E., Aghakouchak, A., Love, C.A., (...), Lima, C.H.R. (2018) Quantifying Changes in Future Intensity-Duration-Frequency Curves Using Multimodel Ensemble Simulations. *Water Resources Research*. https://doi.org/10.1002/2017WR021975

45. Birgisson, B., Ruth, B.E. (2003) Improving Performance Through Consideration of Terrain Conditions: Soils, Drainage, and Climate. *Transportation Research Record*. https://doi.org/10.3141/1819b-47

46. Cox, J.W., McFarlane, D.J., Skaggs, R.W. (1994) Field evaluation of drainmod for predicting waterlogging intensity and drain performance in south-western australia. *Australian Journal of Soil Research*. https://doi.org/10.1071/SR9940653

47. Stancu, M., Cheveresan, M., Zaharia, V., Poienariu, T. (2017) Climate change adaptation in urban areas. Case study for the Tineretului area in Bucharest. *Procedia Engineering*. https://doi.org/10.1016/j.proeng.2017.11.146

48. Malota, M., Senzanje, A. (2015) Modelling mid-span water table depth and drainage discharge dynamics using drainmod 6.1 in a sugarcane field in Pongola, South Africa. *Water SA*. https://doi.org/10.4314/wsa.v41i3.04

49. Gallichand, J. (1994) Numerical simulations of steady-state subsurface drainage with vertically decreasing hydraulic conductivity. *Irrigation and Drainage Systems*. https://doi.org/10.1007/BF00880794

50. Luo, W., Sands, G.R., Youssef, M., (...), Canelón, D. (2010) Modeling the impact of alternative drainage practices in the northern Corn-belt with DRAINMOD-NII. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2009.10.009

51. Willardson, L.S. (1982) Drainage coefficients.. *TRANS. AM. SOC. AGRIC. ENGRS. (GEN. EDN.)*. https://www.scopus.com/pages/publications/0019926197

52. Fouss, J.L., Skaggs, R.W., Fausey, N.R., Pitts, D.J. (2004) Implementing controlled-drainage technology to reduce nitrate loss in drainage water. *Proceedings of the 8th International Drainage Symposium - Drainage VIII*. https://www.scopus.com/pages/publications/27844514844

53. Negm, L.M., Youssef, M.A., Chescheir, G.M., Skaggs, R.W. (2016) DRAINMOD-based tools for quantifying reductions in annual drainage flow and nitrate losses resulting from drainage water management on croplands in eastern North Carolina. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2015.12.014

54. Zhang, J., Fang, H., Zhang, Z., Di, D. (2026) Deep learning-based rapid prediction of urban emergency response capability: A perspective considering the coupled effect of rainfall and drainage pipeline siltation. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2025.134555

55. Narain, V. (2025) Climate Extremes and Peri-Urban Flooding: The Case of Rawta Village, India. *Lecture Notes in Civil Engineering*. https://doi.org/10.1007/978-981-96-2030-2_2

56. Oosterbaan, R.J. (1988) Agricultural criteria for subsurface drainage: A systems analysis. *Agricultural Water Management*. https://doi.org/10.1016/0378-3774(88)90063-7

57. Strock, J., Fausey, N. (2023) Drainage, soil water storage, buffering and filtering. *Encyclopedia of Soils in the Environment, Second Edition*. https://doi.org/10.1016/B978-0-12-822974-3.00215-9

58. Sallam, G.A.H. (2019) The Assessment of Egypt's Subsurface Drainage System. *Handbook of Environmental Chemistry*. https://doi.org/10.1007/698_2017_45

59. Valipour, M. (2014) Drainage, waterlogging, and salinity. *Archives of Agronomy and Soil Science*. https://doi.org/10.1080/03650340.2014.905676

60. Singh, A. (2017) Waterlogging and salinity management for sustainable irrigated agriculture. II: Engineering measures and biodrainage. *Journal of Irrigation and Drainage Engineering*. https://doi.org/10.1061/(ASCE)IR.1943-4774.0001227

61. Pazira, E., Homaee, M. (2010) Salt leaching efficiency of subsurface drainage systems at presence of diffusing saline water table boundary: A case study in Khuzestan Plains, Iran. *ASABE - 9th International Drainage Symposium 2010, Held Jointly with CIGR and CSBE/SCGAB*. https://www.scopus.com/pages/publications/79956082200

62. Youssef, M.A., Abdelbaki, A.M., Negm, L.M., (...), Jaynes, D.B. (2018) DRAINMOD-simulated performance of controlled drainage across the U.S. Midwest. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2017.11.012

63. Memon, N.A., Broughton, R.S., Madramootoo, C.A., (...), von Hoyningen Huene, B. (1986) A method of designing subsurface irrigation/ drainage systems to maximize net benefits. *Canadian Water Resources Journal*. https://doi.org/10.4296/cwrj1104046

64. Ahmadi, M.Z. (1995) A field approach to estimation of humid area drainage coefficients. *Agricultural Water Management*. https://doi.org/10.1016/0378-3774(95)01186-2

65. Li, J., Di, D., Jia, H., (...), Xu, J. (2026) Surrogate-assisted gray–green optimization for urban pluvial flood mitigation under rainfall variability and pipe siltation. *Journal of Water Process Engineering*. https://doi.org/10.1016/j.jwpe.2026.110940

66. Nimantha, H.H.P., Davis, M.P., Udawatta, R.P., (...), Righi, C.A. (2026) Land-use effects on soil moisture profile dynamics in a restrictive claypan soil. *Agrosystems, Geosciences and Environment*. https://doi.org/10.1002/agg2.70453

67. Appels, W.M., Bogaart, P.W., van der Zee, S.E.A.T.M. (2017) Feedbacks Between Shallow Groundwater Dynamics and Surface Topography on Runoff Generation in Flat Fields. *Water Resources Research*. https://doi.org/10.1002/2017WR020727

68. Metzen, D., Sheridan, G.J., Benyon, R.G., (...), Lane, P.N.J. (2019) Spatio-temporal transpiration patterns reflect vegetation structure in complex upland terrain. *Science of the Total Environment*. https://doi.org/10.1016/j.scitotenv.2019.07.357

69. Alan Yeakley, J.A., Swank, W.T., Swift, L.W., (...), Shugart, H.H. (1998) Soil moisture gradients and controls on a southern Appalachian hillslope from drought through recharge. *Hydrology and Earth System Sciences*. https://doi.org/10.5194/hess-2-41-1998

70. Bao, Z., Wu, Y., He, W., (...), Li, Z. (2025) Intensified Rainfall, Growing Floods: Projecting Urban Drainage Challenges in South-Central China Under Climate Change Scenarios. *Applied Sciences (Switzerland)*. https://doi.org/10.3390/app152111577

71. Ding, X., Liao, W., Lei, X., (...), Wang, H. (2022) Assessment of the impact of climate change on urban flooding: A case study of Beijing, China. *Journal of Water and Climate Change*. https://doi.org/10.2166/wcc.2022.224

72. Hu, H., Dong, Q., Xin, C., (...), Wen, J. (2024) Spatial-temporal distribution of extreme rainfall thresholds and assessment of urban waterlogging impact during flood seasons in Shanghai. *Journal of Natural Disasters*. https://doi.org/10.13577/j.jnd.2024.0303

73. Evans, D.E. (2003) WATER RELATIONS OF PLANTS | Plant Responses to Waterlogging. *Encyclopedia of Applied Plant Sciences*. https://doi.org/10.1016/B0-12-227050-9/00113-7
74. Evans, D.E., Gladish, D.K. (2016) Plant Responses to Waterlogging. *Encyclopedia of Applied Plant Sciences*. https://doi.org/10.1016/B978-0-12-394807-6.00083-6
75. Basavaraj, PS, Jangid, Krishna Kumar, Babar, Rohit, (...), Prabhakar, M (2024) Non-invasive measurements to identify mungbean genotypes for waterlogging tolerance. *PeerJ*. https://doi.org/10.7717/peerj.16872
76. Yang, Rui, Li, Murong, Harrison, Matthew Tom, (...), Wang, Xiaoyan (2022) *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2022.890083
77. Ren, Baizhao, Hu, Juan, Liu, Peng, (...), Zhang, Jiwang (2021) Responses of nitrogen efficiency and antioxidant system of summer maize to waterlogging stress under different tillage. *PeerJ*. https://doi.org/10.7717/peerj.11834
78. Pan, Rui, He, Dongli, Xu, Le, (...), Zhang, Wenying (2019) *BMC Genomics*. https://doi.org/10.1186/s12864-018-5405-3
79. https://www.sciencedirect.com/science/article/pii/S1573503622001474
80. Johansson, Eva, Kuktaite, Ramune, Labuschagne, Maryke, (...), Vazquez, Daniel (2023) Adaptation to abiotic stress factors and their effects on cereal and pseudocereal grain quality. *Developing Sustainable and Health Promoting Cereals and Pseudocereals*. https://doi.org/10.1016/B978-0-323-90566-4.00001-1
81. Zhang, Y., Xu, S., Liu, G., (...), Dong, H. (2023) Ridge intertillage alters rhizosphere bacterial communities and plant physiology to reduce yield loss of waterlogged cotton. *Field Crops Research*. https://doi.org/10.1016/j.fcr.2023.108849
82. Talero-Sarmiento, L.H., Lamos-Díaz, H., Marquez-Gonzalez, J.D. (2025) Optimizing cocoa biomass density through integrated irrigation and drainage management under water stress: A linear programming approach. *Ecological Informatics*. https://doi.org/10.1016/j.ecoinf.2025.103262
83. Rahman, Khalil Ur, Ali, Kashmala, Rauf, Mamoona, Arif, Muhammad (2023) *Microorganisms*. https://doi.org/10.3390/microorganisms11082025
84. Manik, S.M.N., Pengilley, G., Dean, G., (...), Zhou, M. (2019) Soil and crop management practices to minimize the impact of waterlogging on crop productivity. *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2019.00140
85. Kamra, S.K. (2015) An overview of subsurface drainage for management of waterlogged saline soils of india. *Water and Energy International*. https://www.scopus.com/pages/publications/84944058473
86. Shokrana, M.S.B., Ghane, E., AbdalAal, Y., Nejadhashemi, A.P. (2023) Predicting the effect of weir management on the discharge of a controlled drainage system in a changing climate. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2023.108534

87. Shedekar, V.S., King, K.W., Fausey, N.R., (...), Brown, L.C. (2021) Exploring the effectiveness of drainage water management on water budgets and nitrate loss using three evaluation approaches. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2020.106501

88. Javani, H.J., Liaghat, A., Hassanoghli, A., Ritzema, R. (2018) Managing controlled drainage in irrigated farmers' fields: A case study in the Moghan plain, Iran. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2018.06.037

89. Sunohara, M., Youssef, M.A., Topp, E., Lapen, D.R. (2010) Measured effect of agricultural drainage water management on hydrology, water quality, and crop yield. *ASABE - 9th International Drainage Symposium 2010, Held Jointly with CIGR and CSBE/SCGAB*. https://www.scopus.com/pages/publications/79956126166

90. Dou, X., Shi, H., Li, R., (...), Wang, B. (2021) Effects of controlled drainage on the content change and migration of moisture, nutrients, and salts in soil and the yield of oilseed sunflower in the hetao irrigation district. *Sustainability (Switzerland)*. https://doi.org/10.3390/su13179835

91. Dukhovny, V., Kenjabaev, S., Yakubov, S., Umirzakov, G. (2018) Controlled Subsurface Drainage as a Strategy for Improved Water Management in Irrigated Agriculture of Uzbekistan. *Irrigation and Drainage*. https://doi.org/10.1002/ird.2259

92. Dou, X., Shi, H., Li, R., (...), Tian, F. (2023) Evaluating the Effects of Controlled Drainage on Nitrogen Uptake, Utilization, Leaching, and Loss in Farmland Soil. *Agronomy*. https://doi.org/10.3390/agronomy13122936

93. Qi, D., Zhu, J. (2023) Controlled Drainage Mediates Cotton Yield at Reduced Nitrogen Rates by Improving Soil Nitrogen and Water Contents. *Journal of Soil Science and Plant Nutrition*. https://doi.org/10.1007/s42729-023-01285-5

94. Baird, A., Frankenberger, J., Bowling, L., Kladivko, E. (2024) IMPACT OF CONTROLLED DRAINAGE ON CROP YIELD INCLUDING WITHIN-FIELD VARIABILITY. *Journal of the ASABE*. https://doi.org/10.13031/ja.15520

95. Poole, C.A., Skaggs, R.W., Cheschier, G.M., (...), Crozier, C.R. (2009) Effects of drainage water management on crop yields. *American Society of Agricultural and Biological Engineers Annual International Meeting 2009, ASABE 2009*. https://www.scopus.com/pages/publications/77649219342

96. Negm, L.M., Youssef, M.A., Jaynes, D.B. (2017) Evaluation of DRAINMOD-DSSAT simulated effects of controlled drainage on crop yield, water balance, and water quality for a corn-soybean cropping system in central Iowa. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2017.03.010

97. Han, X., Zhu, Y., Wang, X., (...), Tang, R. (2025) Assessing the effects of controlled drainage on regional hydrological cycle and crop waterlogging and drought based on a coupled agro-hydrological model. *Agricultural Water Management*. https://doi.org/10.1016/j.agwat.2025.109881