

Equation 1: $\partial_x^2 f(x, y) + \partial_y^2 f(x, y) = 0$

Timothée Chauvin
Jean-Stanislas Denain

February 2019

1 Equation

In this part we consider the Laplace equation:

$$\partial_x^2 f(x, y) + \partial_y^2 f(x, y) = 0$$

with the following Dirichlet border conditions:

$$\forall (x, y) \in \partial\mathbb{D}, f(x, y) = \phi(x, y)$$

where:

- $\mathbb{D} = [0; 1]^2$ is the square of side length 1.
- $\partial\mathbb{D} = \{0; 1\} \times [0; 1] \cup [0; 1] \times \{0; 1\}$ is the border of \mathbb{D} .
- ϕ is a function $\partial\mathbb{D} \rightarrow \mathbb{R}$.

Let $L \in \mathbb{N}^*$: our goal is to build $u : \{0, \dots, L-1\} \times \{0, \dots, L-1\} \rightarrow \mathbb{R}$ such that for every $0 \leq i < L$ and $0 \leq j < L$, $u(i, j)$ approaches $f(\frac{i}{L}, \frac{j}{L})$. We call:

- \mathcal{D} the lattice $\{0, \dots, L\}^2$
- $\partial\mathcal{D} = \{0, L\} \times \{0, \dots, L\} \sqcup \{0, \dots, L\} \times \{0, L\}$ its border

2 Random Walk

Let $(i, j) \in \mathcal{D}$, and $\alpha = \gamma/L^2$: L must be chosen so that $0 \leq \alpha \leq 1$. We define a random walk $(X_n)_{n \in \mathbb{N}}$ on \mathcal{D} . For every $n \in \mathbb{N}$:

- If $X_n = (k, l)$ is **on the border** $\partial\mathcal{D}$ of the lattice, *i.e.* $k \in \{0, L\}$ or $l \in \{0, L\}$, then it **stays put** with probability 1:

$$\mathbb{P}[X_{n+1} = (k, l) | X_n = (k, l)] = 1$$

- If $X_n = (k, l)$ is **inside the lattice**, *i.e.* $0 < k < L$ and $0 < l < L$, it **moves** with equal probability $\frac{1}{4}(1 - \alpha)$ to one of the 4 neighbours of (k, l) : for every neighbour (k', l') of (k, l) ,

$$\mathbb{P}[X_{n+1} = (k', l') | X_n = (k, l)] = \frac{1}{4}$$

3 Approximation

3.1 Defining $u(i, j)$

We now introduce the stopping time $\tau = \inf\{n \in \mathbb{N} | X_n \in \partial\mathcal{D}\}$, the time the **border is reached**. $u(i, j)$ is then defined as the expectation $\mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)] = \mathbb{E}[\tilde{\phi}(X_\tau)\mathbf{1}_{\tau < T} | X_0 = (i, j)]$, where $\tilde{\phi}$ sends $(i, j) \in \partial\mathcal{D}$ to $\phi(i/L, j/L)$.

3.2 What equation does $u(i, j)$ follow?

Let's take a closer look at the $u(i, j)$ defined as above:

$$\begin{aligned} u(i, j) &= \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)] \\ &= \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i+1,j)}] \\ &\quad + \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i-1,j)}] \\ &\quad + \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i,j-1)}] \\ &\quad + \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i,j+1)}] \end{aligned}$$

Now let's focus on $\mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i+1,j)}]$, the first term:

$$\begin{aligned} \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau)\mathbf{1}_{X_1=(i+1,j)}] &= \mathbb{P}_{(i,j)}[X_1 = (i+1, j)] \cdot \mathbb{E}_{(i,j)}[\tilde{\phi}(X_\tau) | X_1 = (i+1, j)] \\ &= \mathbb{P}[X_1 = (i+1, j) | X_0 = (i, j)] \cdot \mathbb{E}[\tilde{\phi}(X_\tau) | X_1 = (i+1, j), X_0 = (i, j)] \\ &= \frac{1}{4} \cdot \mathbb{E}[\tilde{\phi}(X_\tau) | X_1 = (i+1, j)] \\ &= \frac{1}{4} \cdot \mathbb{E}_{(i+1,j)}[\tilde{\phi}(X_\tau)] \\ &= \frac{1}{4} u(i+1, j) \end{aligned}$$

Proceeding similarly with the four other terms, we end up with:

$$u(i, j) = \frac{1}{4} [u(i+1, j) + u(i-1, j) + u(i, j+1) + u(i, j-1)] \quad (*)$$

This can be rewritten as $\tilde{\Delta}u(i, j) = 0$, where $\tilde{\Delta}$ is the discretised Laplacian, sending every function $u : \mathcal{D} \rightarrow \mathbb{R}$ to the function

$$\tilde{\Delta}u : (i, j) \mapsto \frac{1}{4} \sum_{(k,l) \sim (i,j)} [u(k, l) - u(i, j)]$$

3.3 Back to continuous functions

Let's now show how all of this relates to the original partial differential equation: define $f : \mathbb{D} \rightarrow \mathbb{R}$ by $f(x, y) = u(\lfloor xL \rfloor, \lfloor yL \rfloor)$. Then for every $(x, y) \in \mathbb{D}$:

$$f(x+1/L, y) + f(x-1/L, y) - 2f(x, y) = u(\lfloor xL \rfloor + 1, \lfloor yL \rfloor) + u(\lfloor xL \rfloor - 1, \lfloor yL \rfloor) - 2u(\lfloor xL \rfloor, \lfloor yL \rfloor)$$

$$f(x, y+1/L) + f(x, y-1/L) - 2f(x, y) = u(\lfloor xL \rfloor, \lfloor yL \rfloor + 1) + u(\lfloor xL \rfloor, \lfloor yL \rfloor - 1) - 2u(\lfloor xL \rfloor, \lfloor yL \rfloor)$$

Therefore:

$$\frac{1}{4(1/L)^2} [f(x+1/L, y) + f(x-1/L, y) + f(x, y+1/L) + f(x, y-1/L) - 4f(x, y)f(x, y)] = 0$$

Remember, $\alpha = \gamma/L^2$, so that:

$$\frac{1}{4(1/L)^2} [f(x+1/L, y) + f(x-1/L, y) + f(x, y+1/L) + f(x, y-1/L) - 4f(x, y)f(x, y)] = 0$$

When $L \rightarrow \infty$, the left hand side converges to $\Delta f(x, y)$, hence:

$$\boxed{\forall (x, y) \in \mathbb{D}, \Delta f(x, y) = 0}$$

3.4 Border conditions

Let $(x, y) \in \partial\mathbb{D}$: then $(\lfloor xL \rfloor, \lfloor yL \rfloor) \in \partial\mathcal{D}$. Let $i = \lfloor xL \rfloor$ and $j = \lfloor yL \rfloor$, and consider once again the random walk $(X_n)_{n \in \mathbb{N}}$ initially in (i, j) defined above. Since $X_0 = (i, j) \in \partial\mathcal{D}$, the border is reached immediately, therefore the expectation $\mathbb{E}_{(i, j)}[\tilde{\phi}(X_\tau)]$ is simply the value $\tilde{\phi}(X_0) = \tilde{\phi}(i, j) = \phi(x, y)$.

Since this is true for every $L \in \mathbb{N}^*$, by taking the limit as $L \rightarrow \infty$ we also have:

$$\boxed{\forall (x, y) \in \partial\mathbb{D}, f(x, y) = \phi(x, y)}$$

4 Watching the process unfold through time

4.1 Defining $v_n(i, j)$

Define for every $n \in \mathbb{N}$ the function $v_n : \mathcal{D} \rightarrow \mathbb{R}$ sending every $(i, j) \in \mathcal{D}$ to:

$$v_n(i, j) = \mathbb{E}_{(i, j)}[\tilde{\phi}(X_n)]$$

4.2 What equation does $v_n(i, j)$ follow?

Since the walk is symmetric a calculation very similar to the one above leads to:

$$\begin{aligned} v_{n+1}(i, j) &= \mathbb{E}_{(i, j)}[\tilde{\phi}(X_{n+1})] \\ &= \mathbb{E}_{(i, j)}[\tilde{\phi}(X_{n+1})\mathbf{1}_{X_1=(i+1, j)}] \\ &\quad + \mathbb{E}_{(i, j)}[\tilde{\phi}(X_{n+1})\mathbf{1}_{X_1=(i-1, j)}] \\ &\quad + \mathbb{E}_{(i, j)}[\tilde{\phi}(X_{n+1})\mathbf{1}_{X_1=(i, j-1)}] \\ &\quad + \mathbb{E}_{(i, j)}[\tilde{\phi}(X_{n+1})\mathbf{1}_{X_1=(i, j+1)}] \end{aligned}$$

Therefore:

$$v_{n+1}(i, j) - v_n(i, j) = \frac{1}{4} [v_n(i+1, j) + v_n(i, j+1) + v_n(i-1, j) + v_n(i, j-1) - 4v_n(i, j)] \quad (**)$$

4.3 Back to continuous functions

Define $f : \mathbb{R}_+^* \times \mathbb{D} \rightarrow \mathbb{R}$ by $f(t, x, y) = u_{\lfloor nL^2 \rfloor}(\lfloor xL \rfloor, \lfloor yL \rfloor)$. Then for every $(t, x, y) \in \mathbb{R}_+^* \times \mathbb{D}$, multiplying both sides of $(**)$ by L^2 yields:

$$\frac{f(t + 1/L^2, x, y) - f(t, x, y)}{(1/L^2)} = \frac{f(t, x + 1/L, y) + f(t, x - 1/L, y) + f(t, x, y + 1/L) + f(t, x, y - 1/L) - 4f(t, x, y)}{4(1/L)^2}$$

Therefore when $L \rightarrow \infty$:

$$\boxed{\forall (t, (x, y)) \in \mathbb{R}_+^* \times \mathbb{D}, \partial_t f(t, x, y) = \Delta f(t, x, y)}$$

The border conditions are also verified:

$$\boxed{\forall (t, (x, y)) \in \mathbb{R}_+^* \times \partial\mathbb{D}, f(t, x, y) = \phi(x, y)}$$