

# The EIS Filter

David N. DeJong  
University of Pittsburgh

Spring 2008

The **EIS Filter** (DeJong, Dharmarajan, Liesenfeld, Richard, 2007) extends the concept of efficient importance sampling (Richard & Zhang, 2007 *J. of Econometrics*) to applications involving state space representations.

The application of EIS in this context is non-trivial, due to the absence of an analytical expression for the targeted integrand.

Recall the critical elements of the state space representation:

$$f(s_t | Y_t) = \frac{f(y_t, s_t | Y_{t-1})}{f(y_t | Y_{t-1})} = \frac{f(y_t | s_t, Y_{t-1}) f(s_t | Y_{t-1})}{f(y_t | Y_{t-1})}, \quad (1)$$

$$f(s_t | Y_{t-1}) = \int f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1}) ds_{t-1}, \quad (2)$$

$$f(y_t | Y_{t-1}) = \int f(y_t | s_t, Y_{t-1}) f(s_t | Y_{t-1}) ds_t. \quad (3)$$

## Issue:

We lack an analytical expression for  $f(s_{t-1} | Y_{t-1})$  in (2), and for  $f(s_t | Y_{t-1})$  in (3).

# Introduction, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

We overcome the lack of an analytical expression for  $f(s_{t-1}|Y_{t-1})$  using an EIS approximation constructed in the previous period.

We overcome the lack of an analytical expression for  $f(s_t|Y_{t-1})$  by approximating

$$\int f(s_t|s_{t-1}, Y_{t-1}) f(s_{t-1}|Y_{t-1}) ds_{t-1}$$

via an additional EIS calculation (“inner EIS”).

# The EIS Filter

EIS Filter

DND

## Objective:

$$\begin{aligned} f(y_t | Y_{t-1}) &= \int f(y_t | s_t, Y_{t-1}) f(s_t | Y_{t-1}) ds_t \\ &= \int \varphi(s_t) ds_t \end{aligned}$$

**Approach to approximation:** Introduce Importance Sampler

$$f(y_t | Y_{t-1}) = \int \frac{\varphi(s_t)}{g(s_t | a_t)} g(s_t | a_t) ds_t,$$

with  $a_t$  determined from

$$(\hat{a}_t^{l+1}, \hat{c}_t^{l+1}) = \arg \min_{a_t, c_t} \sum_{i=1}^R [\ln \varphi_t(s_{t,l}^i) - c_t - \ln k(s_{t,l}^i; a_t)]^2.$$

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

# The EIS Filter, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

**Resulting approximation:**

$$\hat{f}_N(y_t | Y_{t-1}) = \frac{1}{N} \sum_{i=1}^N \frac{f(y_t | s_t^i, Y_{t-1}) f(s_t^i | Y_{t-1})}{g(s_t^i; \hat{a}_t)}.$$

# The EIS Filter, cont.

## Complication:

- ▶  $f(s_t | Y_{t-1})$  must be evaluated for each MC drawing  $s_t^i$  in constructing  $\hat{f}_N(y_t | Y_{t-1})$ .
- ▶ But as noted, we lack an analytical expression for  $f(s_t | Y_{t-1})$ .

**Resolution (Inner EIS):** weighted-sum approximation of  $f(s_t | Y_{t-1})$ .

# The EIS Filter, cont.

EIS Filter

DND

Write  $f(s_t|Y_{t-1})$  as

$$\begin{aligned} f(s_t|Y_{t-1}) &= \int f(s_t|s_{t-1}, Y_{t-1}) f(s_{t-1}|Y_{t-1}) ds_{t-1} \\ &= \frac{\int f(s_t|s_{t-1}, Y_{t-1}) f(y_{t-1}|s_{t-1}, Y_{t-2}) f(s_{t-1}|Y_{t-2}) ds_{t-1}}{f(y_{t-1}|Y_{t-2})} \\ &= \frac{\int f(s_t|s_{t-1}, Y_{t-1}) f(y_{t-1}|s_{t-1}, Y_{t-2}) f(s_{t-1}|Y_{t-2}) ds_{t-1}}{\int f(y_{t-1}|s_{t-1}, Y_{t-2}) f(s_{t-1}|Y_{t-2}) ds_{t-1}} \end{aligned}$$

**Note:** We already have an optimized sampler for the denominator constructed in the previous period:

$$g(s_{t-1}; a_{t-1}).$$

**Idea:** Use this to approximate both the numerator and denominator.

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS  
Dirac Distributions

Constructing  
Effective Initial  
Samplers, Inner  
EIS

RBC Example



# The EIS Filter, cont.

## Resulting approximation:

$$\hat{f}_N(s_t | Y_{t-1}) = \frac{\sum_{i=1}^S f(s_t | s_{t-1}^{0,i}, Y_{t-1}) \cdot \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})}{\sum_{i=1}^S \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})},$$
$$\omega(s_{t-1}; \hat{a}_{t-1}) = \frac{f(y_{t-1} | s_{t-1}, Y_{t-2}) f(s_{t-1} | Y_{t-2})}{g(s_{t-1} | \hat{a}_{t-1})}.$$

# The EIS Filter, cont.

## Implementation algorithm:

- Inherit  $\{s_{t-1}^{0,i}, \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})\}_{i=1}^N$  from previous period

# The EIS Filter, cont.

EIS Filter

DND

## Implementation algorithm:

- ▶ Inherit  $\{s_{t-1}^{0,i}, \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})\}_{i=1}^N$  from previous period
- ▶ Obtain  $\hat{a}_t$  using

$$(\hat{a}_t^{l+1}, \hat{c}_t^{l+1}) = \arg \min_{a_t, c_t} \sum_{i=1}^R [\ln \varphi_t(s_{t,l}^i) - c_t - \ln k(s_{t,l}^i; a_t)]^2.$$

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

# The EIS Filter, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

## Implementation algorithm:

- ▶ Inherit  $\{s_{t-1}^{0,i}, \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})\}_{i=1}^N$  from previous period
- ▶ Obtain  $\hat{a}_t$  using

$$(\hat{a}_t^{l+1}, \hat{c}_t^{l+1}) = \arg \min_{a_t, c_t} \sum_{i=1}^R [\ln \varphi_t(s_{t,l}^i) - c_t - \ln k(s_{t,l}^i; a_t)]^2.$$

- ▶ Using  $\hat{a}_t$ , generate  $\{s_t^{0,i}\}_{i=1}^N$ , and use to approximate

$$\hat{f}_N(y_t | Y_{t-1}) = \frac{1}{N} \sum_{i=1}^N \frac{f(y_t | s_t^i, Y_{t-1}) f(s_t^i | Y_{t-1})}{g(s_t^i; a_t)}.$$

# The EIS Filter, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

## Implementation algorithm:

- ▶ Inherit  $\{s_{t-1}^{0,i}, \omega(s_{t-1}^{0,i}; \hat{a}_{t-1})\}_{i=1}^N$  from previous period
- ▶ Obtain  $\hat{a}_t$  using

$$(\hat{a}_t^{l+1}, \hat{c}_t^{l+1}) = \arg \min_{a_t, c_t} \sum_{i=1}^R [\ln \varphi_t(s_{t,l}^i) - c_t - \ln k(s_{t,l}^i; a_t)]^2.$$

- ▶ Using  $\hat{a}_t$ , generate  $\{s_t^{0,i}\}_{i=1}^N$ , and use to approximate

$$\hat{f}_N(y_t | Y_{t-1}) = \frac{1}{N} \sum_{i=1}^N \frac{f(y_t | s_t^i, Y_{t-1}) f(s_t^i | Y_{t-1})}{g(s_t^i; a_t)}.$$

- ▶ Pass  $\{s_t^{0,i}, \omega(s_t^{0,i}; \hat{a}_t)\}_{i=1}^N$  to the next period

# Constructing Initial Samplers, Inner EIS

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

(Actually, begin with outer EIS) In establishing an initial sampler for

$$f(y_t | Y_{t-1}) = \int f(y_t | s_t, Y_{t-1}) f(s_t | Y_{t-1}) ds_t,$$

disconnect between  $f(y_t | s_t, Y_{t-1})$  and  $f(s_t | Y_{t-1})$  can cause the EIS algorithm to “crash” (e.g., deliver non-positive-definite VCV matrices).

This very problem is the root of “sample impoverishment” associated with the Particle Filter.

# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

A similar challenge arises given disconnect between  $f(s_t | s_{t-1}, Y_{t-1})$  and  $f(s_{t-1} | Y_{t-1})$  in establishing an initial sampler for

$$f(s_t | Y_{t-1}) = \int f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1}) ds_{t-1}.$$

That is, for the inner EIS, the sampler established for  $f(s_{t-1} | Y_{t-1})$  in period  $t - 1$  may not be effective in delivering an accurate approximation for  $f(s_t | Y_{t-1})$ .

# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Proposition: address these challenges by working with linear-Normal approximations to targeted densities.

(This is the subject of ongoing research; initial results appear highly promising in applications involving DSGE models.)



# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Return to the inner EIS problem

$$f(s_t | Y_{t-1}) = \int f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1}) ds_{t-1}.$$

From period  $t - 1$ , we have a normal approximation (constructed via EIS) for  $f(s_{t-1} | Y_{t-1})$  :

$$f^*(s_{t-1} | Y_{t-1}) \sim N_k(\mu, \Omega).$$

# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

In turn,  $f(s_t | s_{t-1}, Y_{t-1})$  is either linear-normal, or may be approximated as such (e.g., using the log-linear approximation of the model). Then it may be written as

$$f^*(s_t | s_{t-1}, Y_{t-1}) \sim N_k(\Phi s_{t-1}, \Psi).$$

Combining the distributions yields

$$\begin{aligned} f^* \begin{pmatrix} s_t \\ s_{t-1} \end{pmatrix} &\sim N_{2k} \left( \begin{bmatrix} \Phi_k \\ I_k \end{bmatrix} \mu, \begin{bmatrix} \Psi_{k \times k} & 0_{k \times k} \\ 0_{k \times k} & 0_{k \times k} \end{bmatrix} + \begin{bmatrix} \Phi \\ I \end{bmatrix} \Omega \begin{bmatrix} \Phi \\ I \end{bmatrix}' \right) \\ &= N_{2k}(m, \Sigma). \end{aligned}$$

# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Our goal is to implement  $f^* \left( \begin{smallmatrix} s_t \\ s_{t-1} \end{smallmatrix} \right) \equiv f^* (s_t, s_{t-1})$  as a sampler for evaluating

$$f(s_t | Y_{t-1}) = \int f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1}) ds_{t-1}.$$

Recall that we must do this conditionally for each value of  $s_t$  generated in the outer EIS integral. Therefore a final step is to factor  $f^* (s_t, s_{t-1})$  as

$$f^* (s_t, s_{t-1}) = f^* (s_t) f^* (s_{t-1} | s_t).$$

# Constructing Initial Samplers, cont.

EIS Filter

DND

Partitioning:

$$m = \begin{pmatrix} m_1 \\ m_2 \end{pmatrix}, \quad \Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix},$$

with  $m_i$   $k \times 1$  and  $V_{ij}$   $k \times k$ , the distributions we seek are

$$\begin{aligned} f^*(s_t) &\sim N_k(m_1, \Sigma_{11}), \\ f^*(s_{t-1}|s_t) &\sim N_k(m_{2\cdot 1} + \Delta_{21}s_t, \Sigma_{22\cdot 1}), \end{aligned}$$

where

$$\begin{aligned} \Delta_{21} &= \Sigma_{21}\Sigma_{11}^{-1}, \\ m_{2\cdot 1} &= m_2 - \Delta_{21}m_1, \\ \Sigma_{22\cdot 1} &= \Sigma_{22} - \Sigma_{21}\Sigma_{11}^{-1}\Sigma_{12}. \end{aligned}$$

(e.g., see Bauwens, Lubrano, Richard, 1999, OUP).

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

# Constructing Initial Samplers, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Having partitioned,

$$\begin{aligned} f(s_t | Y_{t-1}) &= \int f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1}) ds_{t-1} \\ &= \int \frac{f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1})}{f^*(s_t, s_{t-1})} f^*(s_t, s_{t-1}) ds_{t-1} \\ &= \int \frac{f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1})}{f^*(s_{t-1} | s_t) f^*(s_t)} f^*(s_{t-1} | s_t) f^*(s_t) ds_{t-1} \\ &= f^*(s_t) \int \frac{f(s_t | s_{t-1}, Y_{t-1}) f(s_{t-1} | Y_{t-1})}{f^*(s_{t-1} | s_t) f^*(s_t)} f^*(s_{t-1} | s_t) ds_{t-1} \end{aligned}$$

# Constructing Initial Samplers, cont.

EIS Filter

DND

Implementation: Obtain drawings  $\{s_{t-1}^i\}$  from  $f^*(s_{t-1}|s_t)$ , approximate  $f(s_t|Y_{t-1})$  as

$$\begin{aligned} f(s_t|Y_{t-1}) &\approx f^*(s_t) \frac{1}{R} \sum_{i=1}^R \frac{f(s_t|s_{t-1}^i, Y_{t-1}) f(s_{t-1}^i|Y_{t-1})}{f^*(s_{t-1}^i|s_t) f^*(s_t)} \\ &\equiv f^*(s_t) \frac{1}{R} \sum_{i=1}^R w(s_{t-1}^i). \end{aligned}$$

If  $f^*(s_t, s_{t-1})$  is a good approximation for  $f(s_t|s_{t-1}, Y_{t-1}) f(s_{t-1}|Y_{t-1})$ , then the  $w(s_{t-1}^i)$ 's will be roughly 1's, and

$$f(s_t|Y_{t-1}) \approx f^*(s_t).$$

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

It is often the case that  $f(s_t | s_{t-1}, Y_{t-1})$  is degenerate along certain dimensions.

For example, in the RBC model, the transition equation

$$\left(1 + \frac{g}{1 - \alpha}\right) k_t = i(k_{t-1}, z_{t-1}) + (1 - \delta) k_{t-1}$$

implies degeneracy in the  $k_t$  dimension.

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

In this case, implementation of inner EIS requires an additional step:

the sampler  $f^*(s_t, s_{t-1})$  must be altered so that conditional on  $s_t$ , draws of  $s_{t-1}$  are obtained that satisfy equality restrictions implied by the degeneracy of  $f(s_t | s_{t-1}, Y_{t-1})$ .



# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Let  $s_t$  partition into  $s_t = (p_t, q_t)$ , so that the transition equations consist of two parts:

- ▶ a proper transition density  $f(p_t | s_{t-1}, Y_{t-1})$  for  $p_t$
- ▶ an identity for  $q_t | p_t, s_{t-1}$ :

$$q_t \equiv \phi(p_t, p_{t-1}, q_{t-1}) = \phi(p_t, s_{t-1}).$$

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Assuming that  $\phi(p_t, s_{t-1})$  is differentiable and strictly monotone in  $q_{t-1}$ , with inverse

$$q_{t-1} = \psi(p_t, q_t, p_{t-1}) = \psi(s_t, p_{t-1}),$$

the predictive density becomes

$$f(s_t | Y_{t-1}) = \int J(s_t, p_{t-1}) f(p_t | s_{t-1}, Y_{t-1}) \cdot f(p_{t-1}, q_{t-1} | Y_{t-1})|_{q_{t-1}=\psi(s_t, p_{t-1})} dp_{t-1},$$

where

$$J(s_t, p_{t-1}) = \frac{\partial}{\partial q_t} \psi(s_t, p_{t-1}).$$

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Consider the RBC model. We continue to have a normal approximation (constructed via EIS) for  $f(s_{t-1}|Y_{t-1})$  :

$$f^*(s_{t-1}|Y_{t-1}) \sim N_2(\mu, \Omega).$$

In addition, we have a proper transition density for  $\ln z_t$ , and a degenerate transition for  $k_t$  :

$$f(z_t|s_{t-1}) \sim N_1\left(\begin{bmatrix} 0 & \rho \end{bmatrix} s_{t-1}, \sigma_\varepsilon^2\right).$$

# Dirac Distributions, cont.

Combining these densities yields

$$f^* \left( \begin{array}{c} z_t \\ s_{t-1} \end{array} \right) \sim N_3 \left( \begin{bmatrix} 0 & \rho \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \mu, V \right),$$
$$V = \sigma_\varepsilon^2 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & \rho \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \Omega \begin{bmatrix} 0 & \rho \\ 1 & 0 \\ 0 & 1 \end{bmatrix}'$$

(Compare with  $f^*(s_t, s_{t-1})$  above.)

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Having obtained  $f^* \left( \begin{smallmatrix} z_t \\ s_{t-1} \end{smallmatrix} \right) \equiv f^* (z_t, s_{t-1})$ , we next seek a transformation into  $f^* (s_t, z_{t-1})$ .

Recall why: we must compute the inner EIS for each candidate  $s_t$  to be evaluated in the outer EIS step.

## Dirac Distributions, cont.

To accomplish the transformation, we exploit the linear approximation (in order to retain a linear-normal sampler) of the degenerate state-transition equation

$$k_t = f_{kk} k_{t-1} + f_{kz} z_{t-1},$$

which yields the mapping

$$\begin{aligned} \begin{bmatrix} s_t \\ z_{t-1} \end{bmatrix} &= A \begin{bmatrix} z_t \\ s_{t-1} \end{bmatrix}, \\ A &= \begin{bmatrix} 0 & f_{kk} & f_{kz} \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \\ |A| &= |f_{kk}| \end{aligned}$$

(Note:  $(z, k)$  are understood here to represent logged deviations from ss.)

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Using this mapping,  $f^*(z_t, s_{t-1})$  transforms into

$$f^*(s_t, z_{t-1}) \sim N_{2k}(m, \Sigma),$$

$$m = B\mu,$$

$$B = A \begin{bmatrix} 0 & \rho \\ 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$\Sigma = AVA'$$

Note that since

$$|\Sigma|^{-1} = |f_{kk}| |V|^{-1},$$

the multiplicative term  $|f_{kk}|$  will enter into  $f^*(s_t, z_{t-1})$  (applying the usual change-of-variables formula).

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Having established  $f^*(s_t, z_{t-1})$ , we partition as in the non-degenerate case. Specifically, letting

$$m = \begin{pmatrix} m_1 \\ m_2 \end{pmatrix}, \quad \Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \sigma_\varepsilon^2 \end{pmatrix},$$

with  $m_1$   $2 \times 1$ ,  $m_2$   $1 \times 1$ , etc., we have

$$f^*(s_t, z_{t-1}) = f^*(s_t) f^*(z_{t-1} | s_t),$$

with



# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

$$\begin{aligned}f^*(s_t) &\sim N_k(m_1, \Sigma_{11}), \\f^*(s_{t-1}|s_t) &\sim N_k(m_{2\cdot 1} + \Delta_{21}s_t, \Sigma_{22\cdot 1}),\end{aligned}$$

where

$$\begin{aligned}\Delta_{21} &= \Sigma_{21}\Sigma_{11}^{-1}, \\m_{2\cdot 1} &= m_2 - \Delta_{21}m_1, \\\Sigma_{22\cdot 1} &= \sigma_\varepsilon^2 - \Sigma_{21}\Sigma_{11}^{-1}\Sigma_{12}.\end{aligned}$$

Having partitioned,

$$\begin{aligned} f(s_t | Y_{t-1}) &= \int J(s_t, z_{t-1}) f(z_t | s_{t-1}, Y_{t-1}) \cdot \\ &\quad f(z_{t-1}, k_{t-1} | Y_{t-1}) |_{k_{t-1}=\psi(s_t, z_{t-1})} dz_{t-1} \\ &\approx \int J(s_t, z_{t-1}) |f_{kk}| f^*(s_t) f^*(z_{t-1} | s_t) dz_{t-1} \\ &= f^*(s_t) \int J(s_t, z_{t-1}) |f_{kk}| f^*(z_{t-1} | s_t) dz_{t-1} \end{aligned}$$

# Dirac Distributions, cont.

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Implementation: Obtain drawings  $\{z_{t-1}^i\}$  from  $f^*(z_{t-1}|s_t)$ , approximate  $f(s_t|Y_{t-1})$  as

$$f(s_t|Y_{t-1}) \approx f^*(s_t) |f_{kk}| \frac{1}{R} \sum_{i=1}^R J(s_t, z_{t-1}^i)$$

# Constructing Effective Initial Samplers, Outer EIS

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Recall the objective for outer EIS:

$$f(y_t | Y_{t-1}) = \int f(y_t | s_t, Y_{t-1}) f(s_t | Y_{t-1}) ds_t.$$

Above, we constructed

$$f^*(s_t) \sim N_k(m_1, \Sigma_{11})$$

which serves to approximate  $f(s_t | Y_{t-1})$  (since we expect

$$|f_{kk}| \frac{1}{R} \sum_{i=1}^R J(s_t, z_{t-1})$$

to very nearly be one).

# Constructing Effective Initial Samplers, Outer EIS, cont.

Next, we seek a linear-Normal approximation  $f^*(y_t | s_t, Y_{t-1}) \sim N(\mu(s_t), V)$ . We obtain from our log-linear model approximation

$$\begin{aligned}y_{it} &= \bar{y}_{it} \exp \left( f_{ik} \tilde{k}_{t-1} + f_{iz} \tilde{z}_{t-1} \right) \\ &\approx \bar{y}_{it} \left( 1 + f_{ik} \tilde{k}_{t-1} + f_{iz} \tilde{z}_{t-1} \right), \\ \tilde{k}_{it-1} &= \ln \left( \frac{k_{t-1}}{\bar{k}} \right).\end{aligned}$$

Note that in terms of expected values, we may express  $\tilde{k}_t$  and  $\tilde{z}_t$  as

$$\tilde{k}_t = \frac{\tilde{k}_{t-1}}{\rho_k}, \quad \tilde{z}_t = \frac{\tilde{z}_{t-1}}{\rho_z}.$$

# Constructing Effective Initial Samplers, Outer EIS, cont.

Therefore,

$$\mu(s_t) = r + Ps_t,$$

with

$$\begin{aligned} r_i &= \bar{y}_{it}, \\ P_{i,.} &= \begin{bmatrix} \frac{\bar{y}_i f_{ik}}{\rho_k} & \frac{\bar{y}_i f_{iz}}{\rho_z} \end{bmatrix}. \end{aligned}$$

# Constructing Effective Initial Samplers, Outer EIS, cont.

So we have

$$\begin{aligned}f^*(s_t) &\sim N_k(m_1, \Sigma_{11}), \\f^*(y_t|s_t, Y_{t-1}) &\sim N(\mu(s_t), V).\end{aligned}$$

Letting  $Q = V^{-1}$ , and  $H = \Sigma_{11}^{-1}$ , their product is Normal with mean

$$(H + P'QP)^{-1} [Hm_1 + P'Q(y_t - r)],$$

and variance

$$(H + P'QP)^{-1}.$$

# Example: F-V/R-R (2005 JAE) RBC Model

EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

Experiment: Generate artificial realizations of  $\{y_t, i_t, n_t\}$ ,  $T = 100$  from the RBC model.

Using both the Particle Filter and the EIS filter, recover likelihood estimates at actual parameter values.

Note: Std. dev.s of measurement errors for  $\{y_t, i_t, n_t\}$  :  $[0.000158, 0.0011, 0.000866]$  ; in turn,  $\sigma_\varepsilon = 0.007$ .



## Example, cont.

- ▶ Particle Filter:  $N=60,000$ , CPU time = 40.6 seconds per likelihood evaluation.
- ▶ EIS Filter: Outer EIS draws = 20, one EIS iteration; Inner EIS draws = 10 (no iterations necessary). CPU time = 0.22 seconds per evaluation

Mean, Std. Dev. of likelihood values over 1,000 MC replications:

|                  |          |         |
|------------------|----------|---------|
| Particle Filter: | 1,285.51 | 33.48   |
| EIS Filter:      | 1,299.81 | 0.00177 |

# Example, cont.

EIS Filter

DND

Introduction

The EIS Filter


Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

## Log-likelihoods, Particle Filter



# Example, cont.

EIS Filter

DND

Introduction

The EIS Filter


Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example


## Log-likelihoods, Particle Filter



## Example, cont.

Period-by-period MC standard errors (log scale):

Standard  
Errors



## Example, cont.



Diagnosis of the challenges facing the Particle Filter

Blue dots: EIS representations of  $f(s_t|Y_{t-1})$ ,  $f(s_t|Y_t)$

Small ellipse: EIS representation of


$f(y_t|s_t, Y_{t-1}) f(s_t|Y_{t-1})$

Large ellipse:  $f(y_t|s_t, Y_{t-1})$ ,  $t = 3$



# Example, cont.

$t = 18$



EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS


Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

# Example, cont.

$t = 53$



EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS


Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example

# Example, cont.

$t = 66$



EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions


Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example



# Example, cont.

Likelihood countours:  $\alpha$



EIS Filter

DND

Introduction

The EIS Filter

Constructing  
Effective Initial  
Samplers, Inner  
EIS

Dirac Distributions

Constructing  
Effective Initial  
Samplers, Outer  
EIS

RBC Example