A Cosine-Projection Framework for Choosing Among Cox HR, D-RMST, and Average-Hazard Tests in Survival Trials

Xiang Meng, Lu Tian, Hajime Uno

October 2, 2026

Abstract

Three summary statistics are available for comparing two survival distributions, the hazard ratio (HR) from Cox model, the difference in restricted mean survival time (D-RMST), and the ratio of average hazards (RAH), yet they detect different patterns of treatment effect with different efficiency, and there is no guidance on which to use in a given trial. RAH, in particular, is a recent proposal, whose large-sample efficiency has not been studied systematically. In the paper, we show that all three statistics admit a first-order weighted comparison of cumulative hazard functions, represent each statistic by a direction in $L^2([0,\tau], \lambda_0(v)\,dv)$, and establish that its asymptotic efficiency relative to the oracle test is exactly the squared cosine of the angle between that direction and the true signal for the treatment effect; the asymptotic relative efficiency of any two tests is then a ratio of squared cosines. Because the censoring and accrual design enters only through the at-risk weight function $w(v)$, the cosine metric explains—and predicts—how design choices can reverse the efficiency ordering among tests, something raw weight-function plots cannot do, and it doubles as a design-stage diagnostic for test selection. We illustrate the framework using the CheckMate 057 immunotherapy trial and corroborate the theoretical results through simulations spanning proportional- and non-proportional-hazards settings.

1 Introduction

Three tests are available for comparing two survival distributions in randomized clinical trials, and they occupy very different positions in practice. The test based on Cox proportional hazards model (or the log-rank test) is the only one in widespread use: it is optimal under proportional hazards (PH) assumption and remains the default for regulatory submissions. The test based on restricted mean survival time (RMST) [Uno et al., 2014] is used increasingly often when the proportional hazards assumption is violated, offering a clinically interpretable estimand—the difference in restricted mean survival time (D-RMST) up to a horizon τ. Unlike D-RMST, the ratio of average hazards (RAH) [Uno and Horiguchi, 2023] is an estimand on the *rate* scale: it contrasts the two arms' average event rates per unit

of event-free person-time on a time window. As D-RMST, RAH also doesn't require PH assumption.

Being relatively new, the RAH-based test is also much less well understood than the other two tests. Although its null distribution is known [Uno and Horiguchi, 2023], its large-sample *efficiency*—that is, its power and the alternatives against which it is most sensitive, relative to the two established tests—has not been systematically studied. This is important because practitioners often face a practical question: *given a specific trial design and an anticipated treatment-effect pattern, which test is most appropriate?*

CheckMate 057 was a randomized, open-label Phase III trial comparing nivolumab with docetaxel in 582 patients with previously treated non-squamous NSCLC [Borghaei et al., 2015]. The primary endpoint was overall survival. Individual-level data were reconstructed from the published Kaplan–Meier (KM) curves using the algorithm of Guyot et al. [2012]. We considered two reconstructed datasets, both comprising $n_0 = 290$ docetaxel and $n_1 = 292$ nivolumab patients, that differed only in their censoring patterns. In the "*original*" dataset, the subject-level data were reconstructed from the published figure and the reported riskset information, yielding 419 observed events (226 in the docetaxel group and 193 in the nivolumab group). To illustrate the dependence of test results on the censoring-time distribution, we reconstructed a second dataset with the same marginal KM curves as those reported in the publication but different risk-set sizes over time. This results in only 267 observed events (134 in the docetaxel group and 133 in the nivolumab group). The time window of interest was set to $[0, \tau] = [0, 24]$ months.

Table 1 presents the results of the three aforementioned tests for both reconstructed datasets. The point estimates are broadly consistent between the two datasets (Cox HR: $_{0.75}0.740$ vs. $_{0.84}0.783$; D-RMST: $_{1.73}1.75$ vs. $_{1.75}1.73$ months; RAH: $_{0.74}0.739$ vs. $_{0.74}0.743$), all indicating a survival advantage for nivolumab. In contrast, the corresponding p-values differ substantially (Cox model-based test: $_{p=0.004}p = 0.002$ vs. $_{p=0.157}p = 0.052$; D-RMST-based test: $_{p=0.012}p = 0.012$ vs. $_{p=0.058}p = 0.058$; RAH-based test: $_{p=0.003}p = 0.003$ vs. $_{p=0.014}p = 0.014$). Thus, the p-values for the Cox model- and D-RMST-based tests cross the conventional 0.05 threshold between the two analyses, whereas the RAH-based test remains statistically significant in both datasets. This sensitivity to the censoring distribution does not indicate a failure of any particular test. Rather, it reflects a fundamental interplay among the underlying treatment-effect pattern, the choice of test, and the censoring distribution. Understanding and quantifying this interplay is the central objective of this paper.

The motivating example raises a question that is more general than "is RAH better than D-RMST?". Instead, the correct question is

> *Given assumed survival functions, a treatment-effect pattern, and a censoring/accrual design, which test—Cox, D-RMST, or RAH—is most statistically efficient?*

1.1 Related work

Reservations about the HR as a summary measure of treatment effect are long-standing. When the PH assumption is violated, the Cox maximum partial-likelihood estimator converges to a weighted average of the time-varying log hazard ratio, with weights depending on

Table 1: CheckMate 057 OS overall-survival results under two censoring reconstructions(. All three methods use an analysis horizon of $\tau = 24$ months); for Cox, follow-up is administratively censored at τ before model fitting. Values below 1 (for Cox , and RAH) , or above 0 (for D-RMST) favour , favor nivolumab.

Dataset	Method	Estimate	95% CI	p-value
Original	Cox	HR = 0.750 0.740	(0.618, 0.910) (0.609, 0.898)	0.004 0.002
Original	D-RMST	$\hat{\Delta} = 1.75$ mo 1.75	(0.39, 3.11) (0.39, 3.11)	0.012
Original	RAH	$\hat{\theta}_{\text{RAH}} = 0.739$ 0.739	(0.604, 0.902) (0.604, 0.902)	0.003
Sensitivity	Cox	HR = 0.840 0.783	(0.660, 1.069) (0.612, 1.003)	0.157 0.052
Sensitivity	D-RMST	$\hat{\Delta} = 1.73$ mo 1.73	(−0.06, 3.53) (-0.06, 3.53)	0.058
Sensitivity	RAH	$\hat{\theta}_{\text{RAH}} = 0.743$ 0.743	(0.586, 0.942) (0.586, 0.942)	0.014

the study-specific censoring distribution [Cox, 1972, Hernán, 2010, Horiguchi et al., 2019]. Consequently, two trials with the same underlying biological treatment effect can report materially different HRs simply because their accrual rates or follow-up durations differ. Moreover, because the HR is a relative measure without a direct absolute reference, it can be difficult to translate into clinically actionable terms [Uno et al., 2014]. These concerns have motivated estimands defined directly through the marginal survival function, such as the D-RMST [Royston and Parmar, 2013, Uno et al., 2014, Tian et al., 2014, Uno et al., 2022] and RAH [Uno and Horiguchi, 2023, Meng et al., 2026].

Efficiency comparisons among these tests have received much less attention. The closest antecedent to the present paper is Tian et al. [2018], who compare the logrank test with the D-RMST test through their asymptotic relative efficiency under both PH and non-PH settings. They find that the D-RMST test performs nearly as well as the logrank test when PH holds and can substantially outperform it when PH fails, whereas the logrank test retains an advantage when the two survival curves separate only late in follow-up. When the underlying alternative is unknown, combination (Max-combo) tests trade a small amount of efficiency for greater robustness by considering a family of tests [Lin et al., 2020].

1.2 Contributions and Paper Outline

This paper makes four three contributions. First, we place the Cox-, D-RMST-, and characterize the large-sample efficiency of the RAH-based tests test and place it on a common asymptotic-efficiency scale through Pitman efficacy, extending the comparison of Tian et al. [2018]to include the more recent RAH-based testPitman-efficacy scale with the Cox- and D-RMST-based tests, extending Tian et al. [2018]. Second, we introduce a *cosine/projection metric*, defined as the relative efficiency of a given testwith respect show that each test's efficiency relative to the oracle test . This metric facilitates the comparison of different tests and provides a geometric interpretation of their strengths and limitations under a given alternativeequals the squared cosine of the angle between its test direction and the signal direction, so that any pairwise ARE is a ratio of squared cosines. Third, we study how the censoring distribution and accrual pattern affect the relative efficiency of the three tests. Finally, we demonstrate how because censoring and accrual enter only through the at-risk weight $w(\cdot)$, we use the cosine metric can be used to inform test selection at the study-design stageby considering different plausible patterns of treatment effect, censoring and accrualto show how design choices can reverse the efficiency ordering and to select a test at the design stage. All results are supported

by extensive numerical studies, including simulations and an analysis of the reconstructed CheckMate 057 trial.

The paper is organized as follows. Section 2 defines the three test statistics, shows that each can be expressed as a form of weighted logrank test statistic, and presents their asymptotic efficiency under local alternative. Section 3 develops the cosine metric for facilitating the comparison of the three tests. Section 4 presents simulation studies under a range of settings, and Section 5 reports statistical analysis for a study design based on CheckMate 057 trial. Section 6 summarizes the main contributions and discusses directions for future research.

2 Three Statistics, a Weighted-Log-Rank Representation, and Their Efficiency

2.1 Setup and notation

Let T_k $(k = 0, 1)$ denote the event time in group k with survival function $S_k(t)$, cumulative hazard function $\Lambda_k(t)$, and hazard function $\lambda_k(t)$. Define

$$F_k(t) = 1 - S_k(t) \qquad \text{and} \qquad R_k(t) = \int_0^t S_k(u)\, du. \tag{1}$$

Thus, $F_k(t) = \Pr(T_k \le t)$ is the cumulative incidence rate (the probability that the event has occurred by time t) and $R_k(\tau)$ is the RMST on $(0, \tau]$. Let C_k be the censoring time, independent of T_k, with $G_k(t) = \Pr(T_k \wedge C_k \ge t)$ bounded away from zero on $(0, \tau]$. Let p_k be the limiting proportion of subjects in group k $(p_0 + p_1 = 1)$. Lastly, the at-risk weight function is defined as

$$w(t) = \frac{p_0\, p_1\, G_0(t)\, G_1(t)}{p_0\, G_0(t) + p_1\, G_1(t)}. \tag{2}$$

Two simplifying conditions recur throughout the paper. By *equal allocation* we mean a balanced design with $p_0 = p_1 = 1/2$. By *a common censoring distribution* we mean that the two groups share the same marginal censoring law, and we write $S_C(v) = \Pr(C \ge v)$ for the common censoring survival function; this is the case for administrative censoring. Under these two conditions, the at-risk weight (2) becomes $w(t) = 0.5 \times S_C(t)/\{S_0(t)^{-1} + S_1(t)^{-1}\}$. If the null hypothesis, $S_1(t) = S_0(t), t \in [0, \tau]$, is also true, then the weight function further reduces to $w(t) = S_0(t)S_C(t)/4$.

2.2 The three test statistics

We can test $H_0 : S_1(t) = S_0(t), t \in [0, \tau]$ against alternatives using three statistics. For each statistic, we state the corresponding estimand, point estimator, and asymptotic distribution of the test statistic. Throughout, $\hat{S}_k(\cdot)$ and $\hat{\Lambda}_k(\cdot)$ denote the Kaplan–Meier and Nelson–Aalen estimators of $S_k(\cdot)$ and $\Lambda_k(\cdot)$, respectively, $\hat{G}_k(\cdot)$ denote the empirical counterpart of $G_k(\cdot)$, and $\hat{R}_k(t)$ denote $\int_0^t \hat{S}_k(u)du$, a consistent estimator of $R_k(t)$.

HR. Under the PH assumption,

$$\theta = \log\left\{\frac{\lambda_1(t)}{\lambda_0(t)}\right\}$$

does not depend on t and can be consistently estimated by the maximum partial likelihood estimator $\hat{\theta}$. As $n \to \infty$,

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} N(0, \sigma_\theta^2),$$

where

$$\sigma_\theta^2 = \left\{\int_0^\tau \frac{e^\theta p_0 p_1 \, G_0(u) \, G_1(u)}{p_0 \, G_0(u) + e^\theta p_1 \, G_1(u)} \, d\Lambda_0(u)\right\}^{-1}. \tag{3}$$

Under $H_0 : \lambda_1(\cdot) = \lambda_0(\cdot)$,

$$\sigma_\theta^2 = \left\{\int_0^\tau w(u) \, d\Lambda_0(u)\right\}^{-1},$$

and the Wald statistic $Z_{\text{Cox}} = \sqrt{n}\,\hat{\theta}/\hat{\sigma}_\theta$ is asymptotically standard normal, where $\hat{\sigma}_\theta^2$ is a consistent estimator of σ_θ^2.

We use Z_{Cox} for all numerical studies in this paper, whereas its theoretical properties were examined based on an asymptotically equivalent score statistic, i.e., the log-rank statistic. The details are given in Section 2.3 and Appendix B.

D-RMST. The **D-RMST** test is based on the difference between two estimated RMSTs:

$$\hat{\theta}_{D\text{-RMST}} = \hat{R}_1(\tau) - \hat{R}_0(\tau) = \int_0^\tau \left[\hat{S}_1(t) - \hat{S}_0(t)\right] dt, \tag{4}$$

the area between the two Kaplan–Meier curves up to τ [Uno et al., 2014, 2022]. Let $\theta_{D\text{-RMST}} = R_1(\tau) - R_0(\tau)$ be the true RMST difference. It is a clinically interpretable estimand, and oftentimes preferred when the PH assumption may be violated. Then, $\hat{\theta}_{D\text{-RMST}}$ is a consistent estimator of $\theta_{D\text{-RMST}}$ and $\sqrt{n}(\hat{\theta}_{D\text{-RMST}} - \theta_{D\text{-RMST}}) \xrightarrow{d} N(0, \sigma^2_{D\text{-RMST}})$ with

$$\sigma^2_{D\text{-RMST}} = \sum_{k=0}^{1} \int_0^\tau \left\{\int_v^\tau S_k(u) \, du\right\}^2 \frac{d\Lambda_k(v)}{p_k \, G_k(v)},$$

as $n \to \infty$ [Tian et al., 2018]. Under the null, the Wald statistic $Z_{D\text{-RMST}} = \sqrt{n}\,\hat{\theta}_{D\text{-RMST}}/\hat{\sigma}_{D\text{-RMST}}$ is asymptotically standard normal, where $\hat{\sigma}^2_{D\text{-RMST}}$ is a consistent estimator of $\sigma^2_{D\text{-RMST}}$.

RAH. The AH for group k on $(0, \tau]$ is

$$\eta_k(\tau) = \frac{F_k(\tau)}{R_k(\tau)} = \int_0^\tau \lambda_k(t) \, \frac{S_k(t)}{\int_0^\tau S_k(u) \, du} \, dt,$$

a weighted average of hazard function over $[0, \tau]$ [Uno and Horiguchi, 2023]. $\eta_k(\tau)$ is a population-level person-time event rate. Two properties make AH attractive as a robust summary on the hazard scale. First, $\eta_k(\tau)$ is defined purely through the event-time distribution and is therefore not affected by the censoring distribution. On the other hand, the

commonly used person-time incidence rate $\Pr(T \le C)/\mathbb{E}(T \wedge C)$ depends on both survival and censoring distributions [Uno and Horiguchi, 2023]. Second, AH respects stochastic ordering: if $S_1(t) \ge S_0(t)$ for all $t \le \tau$, then $\eta_1(\tau) \le \eta_0(\tau)$. The ratio of AH (RAH) between two groups is

$$\theta_{\mathrm{RAH}} = \frac{\eta_1(\tau)}{\eta_0(\tau)} = \frac{F_1(\tau)/R_1(\tau)}{F_0(\tau)/R_0(\tau)}, \tag{5}$$

with $\theta_{\mathrm{RAH}} = 1$ under H_0. Its estimator $\hat{\theta}_{\mathrm{RAH}}$ can be obtained by replacing $F_k(\tau)$ and $R_k(\tau)$ with $1 - \hat{S}_k(\tau)$ and $\hat{R}_k(\tau)$ in (5), respectively. As $n \to \infty$, $\sqrt{n}(\log \hat{\theta}_{\mathrm{RAH}} - \log \theta_{\mathrm{RAH}}) \xrightarrow{d} N(0, \sigma^2_{\mathrm{RAH}})$, [Uno and Horiguchi, 2023], where

$$\sigma^2_{\mathrm{RAH}} = \sum_{k=0}^{1} \int_0^{\tau} \left\{ \frac{1}{F_k(\tau)} - \frac{R_k(u)}{R_k(\tau)} \right\}^2 \frac{d\Lambda_k(u)}{G_k(u)\, p_k}. \tag{6}$$

The test statistic $Z_{\mathrm{RAH}} = \sqrt{n} \log(\hat{\theta}_{\mathrm{RAH}})/\hat{\sigma}_{\mathrm{RAH}}$ is asymptotically $N(0,1)$ under H_0, where $\hat{\sigma}^2_{\mathrm{RAH}}$ is a consistent estimator of σ^2_{RAH}.

2.3 Weighted-log-rank representation

All three statistics, to the first order, can be written as a weighted contrast of Nelson–Aalen cumulative-hazard estimates from two groups:

$$U_W = \int_0^{\tau} W(t) d\hat{\Delta}_{\Lambda}(t), \tag{7}$$

where

$$\hat{\Delta}_{\Lambda}(v) = \hat{\Lambda}_1(v) - \hat{\Lambda}_0(v).$$

Under H_0, the two cumulative hazards coincide and U_W has mean zero. The weight $W(\cdot)$ decides which time periods contribute to the comparison, and choosing $W(\cdot)$ to match the anticipated alternative is the classical route to constructing powerful, distribution-free survival tests [Fleming and Harrington, 1991, Tian et al., 2018]. Rigorous arguments for expressing three aforementioned test statistics in this form can be found in Appendix B.

HR. The log-rank statistic, the score statistic under Cox model, can be written as

$$U_{\mathrm{Cox}} = \int_0^{\tau} \frac{p_1 p_0 \hat{G}_1(t) \hat{G}_0(t)}{p_1 \hat{G}_1(t) + p_0 \hat{G}_0(t)} d\hat{\Delta}_{\Lambda}(t). \tag{8}$$

Under the local alternatives considered in section 2.4, a one-step expansion of the score function from the partial likelihood function around $\theta = 0$ yields that $\sqrt{n}\,\hat{\theta} = \sigma^2_{\theta} \sqrt{n}\, U_{\mathrm{Cox}} + o_p(1)$. Thus, the Wald statistic, Z_{Cox} and the normalized $\sqrt{n} U_{\mathrm{Cox}}$ are asymptotically equivalent. Furthermore, the random weight in (8) converges to at-risk weight $w(\cdot)$ in (2) in probability, as $n \to \infty$,

D-RMST and RAH. Neither statistic is a log-rank type statistic by construction, but each is a smooth functional of $\hat{\Lambda}_1(\cdot)$ and $\hat{\Lambda}_0(\cdot)$. For D-RMST, $R(\Lambda) = \int_0^\tau \exp\{-\Lambda(t)\}\, dt$, and

$$\begin{aligned}
\hat{\theta}_{D\text{-RMST}} &= \int_0^\tau S_0(t) \left[\exp\left\{-\hat{\Delta}_\Lambda(t)\right\} - 1\right] dt + O\big(\|\hat{\Delta}_\Lambda\|^2\big) \\
&= -\int_0^\tau S_0(t)\, \hat{\Delta}_\Lambda(t)\, dt + O\big(\|\hat{\Delta}_\Lambda\|^2\big) \\
&= -\int_0^\tau \{R_0(\tau) - R_0(t)\}\, d\hat{\Delta}_\Lambda(t) + O\big(\|\hat{\Delta}_\Lambda\|^2\big) \\
&= -U_{D\text{-RMST}} + O_p\big(n^{-1}\big)
\end{aligned}$$

under local alternative. As the remainder is $O_p(n^{-1})$, $Z_{D\text{-RMST}}$ and the normalized $\sqrt{n}U_{D\text{-RMST}}$ are asymptotically equivalent. RAH can be handled the same way, with an extra delta-method step for the ratio $F_k(\tau)/R_k(\tau)$, and

$$\begin{aligned}
\log(\hat{\theta}_{\text{RAH}}) =& \int_0^\tau \left[\frac{S_0(\tau)}{F_0(\tau)} + \frac{R_0(\tau) - R_0(t)}{R_0(\tau)}\right] d\hat{\Delta}_\Lambda(t) + O\big(\|\hat{\Delta}_\Lambda\|^2\big) \\
=& U_{\text{RAH}} + O_p(n^{-1})
\end{aligned}$$

under local alternative. As a result, Z_{RAH} and the normalized $\sqrt{n}U_{\text{RAH}}$ are asymptotically equivalent.

All U_X are oriented so that $U_X < 0$ indicates benefit of treatment (hence $U_{D\text{-RMST}} = -\hat{\theta}_{D\text{-RMST}}$ to first order); since ξ_X depends on μ_X^2, the orientation does not affect efficiency.

In summary, under equal allocation and a common censoring assumption, the resulting weights for U_{Cox}, $U_{D\text{-RMST}}$ and U_{RAH} are

$$\begin{aligned}
W_{\text{Cox}}(t) &= S_0(t)\, S_C(t)/4, \\
W_{D\text{-RMST}}(t) &= R_0(\tau) - R_0(t), \text{ and} \\
W_{\text{RAH}}(t) &= \left[\frac{S_0(\tau)}{F_0(\tau)} + \frac{R_0(\tau) - R_0(t)}{R_0(\tau)}\right],
\end{aligned}$$

respectively. Since multiplying the weight function by any nonzero constant yields an equivalent test statistic, without loss of generality, we may use following equivalent weight functions

$$\begin{aligned}
W_{\text{Cox}}(t) =& S_0(t)\, S_C(t) \\
W_{D\text{-RMST}}(t) =& \frac{R_0(\tau) - R_0(t)}{R_0(\tau)} \\
\text{and } \ W_{\text{RAH}}(t) =& S_0(\tau) + \{1 - S_0(\tau)\}\, W_{D\text{-RMST}}(t)
\end{aligned}$$

for three test statistics. On the current scale, all three weights equal 1 at $t = 0$, their limits at $t = \tau$ are $S_0(\tau)S_C(\tau)$, 0, and $S_0(\tau)$. Also, weight for U_{RAH} is a simple linear transformation of that for $U_{D\text{-RMST}}$.

The Cox model-based log-rank weight tracks the amount of information in the risk set. It is the largest early in follow-up, when the risk set is large, and decreases over time as deaths

and censoring deplete the risk set. Under the PH assumption, this weighting is optimal. The D-RMST weight represents the RMST within the window $[t, \tau]$. It also places greater weight on earlier follow-up, which decreases monotonically to zero at τ, thereby also down-weighting survival differences in late follow-up. The weight for RAH is likewise monotone decreasing, but remains strictly positive at τ, and therefore places relatively greater emphasis on late-stage survival differences than the weight function for D-RMST. This common representation allows us to formally examine the test efficiency in the next section.

2.4 Asymptotic efficiency

To compare the three tests on a single scale we use the *Pitman efficacy*, the standard local-power measure for asymptotically normal tests [Pitman, 1949]. Following Tian et al. [2018], we embed the two-sample problem in a sequence of contiguous (local) alternatives indexed by a fixed log-hazard-ratio shape $\alpha(\cdot)$ on $(0, \tau]$,

$$H_{An}: \qquad \log\left\{\frac{\lambda_{1n}(t)}{\lambda_0(t)}\right\} = \frac{\alpha(t)}{\sqrt{n}}, \tag{9}$$

so that the two hazards coincide as $n \to \infty$, while the between-group difference is in the order of $O(1)$. Under (9), for test $X \in \{\text{Cox}, D\text{-RMST}, \text{RAH}\}$, the test statistic is asymptotically normal, i.e.,

$$\sqrt{n}\, U_X \xrightarrow{d} N\big(\mu_X,\ \sigma_X^2\big),$$

where

$$\mu_X = \int_0^\tau W_X(t)\, \alpha(t)\, \lambda_0(t)\, dt \qquad \text{and} \qquad \sigma_X^2 = \int_0^\tau \frac{W_X(t)^2}{w(t)}\, \lambda_0(t)\, dt.$$

The drift μ_X is obtained by substituting $\sqrt{n}\hat{\Delta}_\Lambda(t)$ by $\alpha(t)\lambda_0(t)$ in (7). The variance $\hat{\sigma}_X^2$ is the null variance of (7). The *asymptotic efficiency* (AE), or Pitman efficacy, of test X is the squared standardized drift

$$\xi_X = \frac{\mu_X^2}{\sigma_X^2}, \tag{10}$$

which governs the local power. The derivation of μ_X for RAH-based test can be found in Appendix A. Furthermore, the explicit expressions of ξ_{Cox}, $\xi_{D\text{-RMST}}$ and ξ_{RAH} are provided in Appendix A.2. The *asymptotic relative efficiency* (ARE) of test X versus test Y is the ratio of corresponding AEs:

$$\text{ARE}(X, Y) = \frac{\xi_X}{\xi_Y}, \tag{11}$$

which approximates the inverse ratio of the sample sizes the two tests require to reach the same local power against (9); $\text{ARE}(X, Y) > 1$ means that test X is more efficient and needs proportionally fewer subjects than Y [Tian et al., 2018, Pitman, 1949]. Because ARE may be a poor proxy for the sample size ratio in finite-sample setting, we also report the *empirical relative efficiency* (ERE), the ratio of squared standardized effect sizes estimated by simulation (Section 4),

$$\text{ERE}(X, Y) = \frac{E(U_X)^2/\text{Var}(U_X)}{E(U_Y)^2/\text{Var}(U_Y)}, \tag{12}$$

with the expectations and variances approximated by their Monte Carlo averages in simulation [Tian et al., 2018].

3 The geometric interpretation of AE

Section 2.3 reduced all three tests to the weighted logrank test (7) and showed that their AE depends on the test only through the weight function. This is the basis of the classical comparison of Tian et al. [2018], who contrast the Cox and D-RMST tests by inspecting their weight functions $W_{\text{Cox}}(\cdot)$ and $W_{D\text{-RMST}}(\cdot)$ directly. In this section, we replace the side-by-side weight comparison with a geometric quantity, a cosine, turning AE into an angle in a function space.

3.1 Signal direction and test direction

Suppose that all candidate statistics are evaluated on the same horizon τ, and consider the function space $L^2([0,\tau], \lambda_0(v)\,dv)$, where the inner product

$$\langle f, g\rangle = \int_0^\tau f(v)\, g(v)\, \lambda_0(v)\, dv, \tag{13}$$

and norm $\|f\|^2 = \langle f, f\rangle$. Next, we define the *signal direction* for local alternative $\alpha(\cdot)/\sqrt{n}$ and *test direction* for test $X \in \{\text{Cox}, D\text{-RMST}, \text{RAH}\}$, as

$$s_\alpha(t) = \alpha(t)\sqrt{w(t)} \qquad \text{and} \qquad h_X(t) = \frac{W_X(t)}{\sqrt{w(t)}}, \tag{14}$$

respectively.

The signal direction, $s_\alpha(t)$, describes where the detectable treatment effect is by combining the local log-HR representing the underlying between-group difference at t and $\sqrt{w(t)}$ representing the amount of information available to detect the difference at time t. Thus, $s_\alpha(t)$ is large where the true effect is big and sufficient information is available to detect it. It is a "direction" since only its shape over t, not its overall magnitude, matters. The test direction, $h_X(t) = W_X(t)/\sqrt{w(t)}$, is the analogous object built from the test's own weight. $W_X(\cdot)$ records where the test is prepared to look for the effect. $\sqrt{w(t)}$ in the denominator ensures that $h_X(t)$ represents test weight per unit of information available at time t. Next, we have

$$\langle s_\alpha, h_X\rangle = \int_0^\tau \alpha(t)\sqrt{w(t)}\,\frac{W_X(t)}{\sqrt{w(t)}}\,\lambda_0(t)\,dt = \int_0^\tau W_X(t)\,\alpha(t)\,\lambda_0(t)\,dt = \mu_X,$$

$$\|h_X\|^2 = \int_0^\tau \frac{W_X(v)^2}{w(v)}\,\lambda_0(v)\,dv = \sigma_X^2,$$

Consequently the AE (10)

$$\xi_X = \frac{\langle s_\alpha, h_X\rangle^2}{\|h_X\|^2}. \tag{15}$$

3.2 The cosine metric

Writing the inner product in (15) as

$$\langle s_\alpha, h_X \rangle = \|s_\alpha\| \, \|h_X\| \cos(\theta_X),$$

or equivalently,

$$\cos^2(\theta_X) = \frac{\langle s_\alpha, h_X \rangle^2}{\|s_\alpha\|^2 \|h_X\|^2} = \frac{\xi_X}{\|s_\alpha\|^2}, \tag{16}$$

where θ_X denotes the angle between the signal direction s_α and the test direction h_X. Intuitively, θ_X measures the mismatch between where the test places its weight and where the detectable treatment effect is concentrated: a small θ_X indicates good alignment and high local power. The quantity $\|s_\alpha\|^2$ is the AE of the oracle test using the optimal weight $\alpha(t)w(t)$. Therefore, $\cos^2(\theta_X)$ is the ARE of test X relative to the oracle test, O. In general, the ARE of test X vs test Y can be expressed as

$$\mathrm{ARE}(X, Y) = \frac{\cos^2(\theta_X)}{\cos^2(\theta_Y)} = \frac{\mathrm{ARE}(X, O)}{\mathrm{ARE}(Y, O)}. \tag{17}$$

As a preview, Figure 1 provides geometric representations for three data-generating processes considered in [Tian et al., 2018]. Specifically, the left column displays $s_\alpha(t)$, $h_{\mathrm{Cox}}(t)$, $h_{\mathrm{DRMST}}(t)$, and $h_{\mathrm{RAH}}(t)$, whereas the right column depicts the corresponding directions on a compass, with the signal direction fixed at 12 o'clock and each test's $\cos^2\theta_X$ shown at the tip of the corresponding arrow for test direction. More detailed results are presented in Section 4.

The cosine metric naturally applies at the trial design stage. Suppose that one has access to:

- assumed survival functions $S_0(t)$ and $S_1(t)$;
- assumed treatment-effect $\alpha(t)$ (e.g., PH, delayed benefit, et al.);
- planned censoring and accrual distribution (which affects $w(\cdot)$); and
- a set of test statistics.

Given these inputs, one can compute the signal and test directions and $\cos^2(\theta_X)$ for each candidate test and select the test with the highest $\cos^2(\theta_X)$, the one most aligned with the expected signal. Crucially, censoring and accrual are part of trial design and affect the value of $\cos^2(\theta_X)$. By varying $S_C(t)$, one can trace how the cosine metric changes and understand whether the selected test is robust to design assumptions. We demonstrate this in Section 5 for CheckMate 057, and specifically trace the preferred test across design choices in a sensitivity analysis.

4 Simulation Study

4.1 Design

We compare the Cox, D-RMST, and RAH-based tests via ARE and ERE using the same data-generating process of Tian et al. [2018]. The baseline event-time distribution is Weibull(κ, η_0) with scale parameter η_0 chosen so that $S_0(40)$ $S_0(40)$ equals a specified target survival rate at $\tau = 40$ $\tau = 40$ months and the default shape parameter $\kappa = 0.826$ yielding a decreasing hazard $\lambda_0(t) \propto t^{-0.174}$. Censoring time $C = \min(E, U)$, where $E \sim \text{Exp}(\lambda_c)$ and $U \sim \text{Uniform}(24, 43)$ represent loss of follow-up and random entry, respectively. Two rates $\lambda_c \in \{0.0006, 0.025\}$ for E yield light ($S_C(40) = 0.15$$S_C(40) = 0.154$) and heavy ($S_C(40) = 0.04$$S_C(40) = 0.058$) censoring. Sample size $n_0 = n_1 = 400$ per arm; $B = 2000$ replicates are used for each simulation. We consider the following alternatives:

PH scenario (Case 1): The event-time distribution in the treatment group is Weibull(κ, η_1), where $\eta_1 = \eta_0/0.70^{1/\kappa}$ yields a HR of 0.70. η_0 is adjusted so that $S_0(40) = 0.10, \ldots, 0.80$ and 0.9.

Non-PH scenarios: $\kappa = 0.826$ and $S_0(40) = 0.50$ for control subjects. The log-HR $\alpha(t) = \log[\lambda_1(t)/\lambda_0(t)]$ follows two patterns:

- **Case 2 (fading benefit):** $\alpha(t) = -c_0 \max(43 - t, 0)^s, s \in \{0.5, 1, 2\}$$\alpha(t) = -c_0 \max(43 - t, 0)^s, s \in \{0.5, 1, 2\}$. The treatment benefit is the biggest early and fades to null at month 43. 43.

- **Case 3 (delayed benefit):** $\alpha(t) = -c_0 t^s, s \in \{0.5, 1, 2\}$. The treatment benefit is absent at $t = 0$ and increases over follow-up.

In both cases $c_0 > 0$ is selected so that the mean log-HR over $[0, \tau]$ is approximately $\log(0.70)$. For Cases 2 and 3, event-times in treatment arm are generated by numerical inversion of the cumulative hazard, i.e., $T_1 \sim \Lambda_1^{-1}\{\text{Exp}(1)\}$.

Figures 2, 3, and 4 display the survival curves and HRs for there three cases. Each figure uses a four-panel layout: the top-left panel shows $\exp\{\alpha(t)\}$ (the HR), and the remaining panels show $S_0(t)$ (control, solid) against $S_1(t)$ (treatment, dashed) for different values of s.

Three simulation settings are also studied:

1. the PH setting with a HR=0.82 (Appendix C);

2. the PH setting with alternative baseline survival distributions (Weibull(κ, η_0) with $\kappa = 0.7, 1$ and 1.5), sample size (150 per arm), and trial horizon ($\tau = 10$) month (Appendix D);

3. the crossing-hazard setting with $\alpha(t) = \log(0.46) + (t/43)$ (the HR begins below 1 and rises to 1.25 at month 43) (Appendix E).

4.2 PH scenario (Case 1)

Table 2 summarizes the comparison for $S_0(40) \in \{0.10, \ldots, 0.90\}$, $\kappa = 0.826$, and HR $= 0.70$. Under light censoring, the ARE of the D-RMST-based test relative to the Cox model-based test decreases gradually from 0.99 to 0.90 Cox decreases gradually as $S_0(40)$ increases. Under heavy censoring, however, this ARE remains close to 1.0 1 across the full range of $S_0(40)$.

The ARE of the RAH-based test relative to the Cox model-based test is also below 1 throughout. This is expected as the Cox model-based test is optimal under the PH assumption. Under light censoring, it decreases from 0.94 to 0.83 The efficiency declines as $S_0(40)$ increases . This pattern is expected. and more strongly under heavy censoring. As $S_0(\tau)$ increases, the lower bound $W_{\mathrm{RAH}}(\tau) = S_0(\tau)$ becomes larger, so that $h_{\mathrm{RAH}}(\tau) = 2\sqrt{S_0(\tau)/S_C(\tau)}$ also increases. This inflates $\|h_{\mathrm{RAH}}\|^2$ and enlarges RAH's discrepancy from the optimal test direction. Under At $S_0(40) = 0.10$ under heavy censoring, this loss of efficiency is more pronounced: the AREdecreases from 0.81 at $S_0(40) = 0.10$ to 0.64 at $S_0(40) = 0.90$, because a larger $S_0(\tau)$ inflates $\|h_{\mathrm{RAH}}\|^2$ even more rapidly, when $S_C(\tau)$ is small. In summary, the efficiency loss from using the RAH-based test is modest under light censoring, with a maximum of approximately 17%, but can reach about 37% under heavy censoringARE(RAH/Cox) is 0.798, whereas ARE(RAH/D-RMST) is 0.812. The ordering

$$\text{Cox} \geq \text{RAH} \approx D\text{-RMST}$$

under PH with light censoring and small $S_0(\tau)$ is also evident geometrically in Figure 1 (top row): the signal direction is aligned with the Cox model-based test direction, whereas the RAH and D-RMST test directions are more off-axis.

The ARE of the RAH-based test relative to the D-RMST-based test is close to 1 under light censoring , but ranges from approximately 0.64 to 0.81 but substantially lower under heavy censoring, indicating a more substantial pronounced performance separation between two testing the two procedures.

Table 3 reports results for $S_0(40) = 0.50$ and $\kappa \in \{0.5, 0.826, 1, 1.5, 2\}$. The Under light censoring, the ARE of the RAH-based test relative to the Cox model-based test decreases from 0.92 0.922 when $\kappa = 0.5$, corresponding to a decreasing baseline hazard, to 0.73 0.737 when $\kappa = 2$, corresponding to an increasing baseline hazard. Intuitively, an increasing hazard shifts more events toward later follow-up, where RAH weighting differs more substantially from optimal weighting of Cox model-based test, thereby reducing its efficiency.

Both Tables 2 and 3 show systematically larger efficiency losses for the RAH-based test under heavy censoring than under light censoring. This pattern persists across alternative baseline survival distributions and trial horizons.

In summary, these results show that the ARE of the competing tests depends materially on the censoring distribution. The censoring and accrual patterns induced by the study design should not be viewed merely as nuisance features; rather, they are important design inputs for choosing testing procedures.

4.3 Non-PH: fading benefit (Case 2)

Table 4 presents the two monotone non-PH scenarios with $\kappa = 0.826$, $S_0(40) = 0.50$, and $s = 1$. Under Case 2, the D-RMST-based test slightly outperforms the RAH-based test, with the Cox model-based test performing intermediately, i.e.,

$$D\text{-RMST} \geq \text{Cox} \geq \text{RAH}.$$

Table 2: Theoretical ARE and empirical ERE under PH (HR= 0.70), Tian DGP. Weibull baseline ($\kappa = 0.826$), $\tau = 40$. Notation: R/C = RMST/Cox, A/C = RAH/Cox, A/R = RAH/RMST. $N = 400$ $n = 400$ per arm, $B = 2000$.

$S_0(40)$ Censoring Metric 0.10 0.30 0.50 0.70 0.90 ARE(R/C) 0.992 0.964 0.939 0.916 0.897 ARE(A/C) 0.940 0.896 0.868 0.847 0.831 ARE(A/R) 0.947 0.929 0.924 0.924 0.927 ERE(R/C) 1.015 0.967 0.966 0.931 0.930 ERE(A/C) 0.916 0.869 0.865 0.851 0.813 ERE(A/R) 0.902 0.899 0.895 0.914 0.874 ARE(R/C) 0.982 0.996 0.996 0.993 0.987 ARE(A/C) 0.797 0.719 0.679 0.653 0.634 ARE(A/R) 0.812 0.722 0.682 0.658 0.642 ERE(R/C) 0.989 1.024 1.014 1.014 1.035 ERE(A/C) 0.767 0.674 0.716 0.663 0.666 ERE(A/R) 0.775 0.658 0.707 0.654 0.643

Censoring	$S_0(40)$	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
Light	0.10	0.994	0.941	0.947	1.007	0.946	0.940
Light	0.30	0.967	0.898	0.929	0.978	0.860	0.879
Light	0.50	0.942	0.871	0.924	0.948	0.893	0.942
Light	0.70	0.920	0.851	0.924	0.912	0.866	0.950
Light	0.90	0.901	0.835	0.927	0.919	0.832	0.905
Heavy	0.10	0.982	0.798	0.812	0.997	0.725	0.727
Heavy	0.30	0.997	0.720	0.722	1.020	0.691	0.677
Heavy	0.50	0.998	0.680	0.682	1.024	0.671	0.656
Heavy	0.70	0.995	0.654	0.658	1.011	0.677	0.670
Heavy	0.90	0.990	0.635	0.642	1.019	0.658	0.646

Because the treatment effect is concentrated in the early follow-up period, the down-weighting of late differences by D-RMST causes little efficiency loss. In contrast, the RAH-based test continues to assign positive weight to late follow-up times, where the treatment effect has largely disappeared. This leads to the expected ordering, D-RMST $\geq$ Cox $\geq$ RAH. The corresponding geometric interpretation is shown in the middle row of Figure 1.

4.4 Non-PH: delayed benefit (Case 3)

Under Case 3, the ARE of the RAH-based test relative to the Cox model-based test exceeds 1 under both censoring regimes. The treatment effect is concentrated at late time, whereas the weight for D-RMST approaches zero near τ, effectively down-weighting the events in the most informative time window. The Cox model also assigns decreasing weight to later follow-up times, as its weight function $w(t)$ declines, but this down-weighting is less pronounced than that of D-RMST. In contrast, the RAH-based test maintains a bigger positive weight near τ, as $S_0(\tau) > S_0(\tau)S_C(\tau) > 0$. Therefore, it preserves a greater alignment with the delayed treatment effect, and yields the expected ordering,

$$\text{RAH} \geq \text{Cox} \geq \text{D-RMST}.$$

See Figure 1 (bottom row) for the corresponding geometric interpretation.

In all simulations, the EREs are close to the corresponding AREs, especially in terms > 1 vs < 1, supporting the validity of ARE under finite sample. Across the reported simulations, ARE and ERE generally agree in direction; disagreements occur only for comparisons close to 1 (within about 0.07), where Monte Carlo variation and finite-alternative effects can reverse the side of 1.

Table 3: ARE and ERE under PH (HR= 0.70), varying Weibull shape. $S_0(40) = 0.50$, $\tau = 40$, $N = 400$ $n = 400$ per arm, $B = 2000$.

κ Censoring ARE(R/C) ARE(A/C) ARE(A/R) ERE(R/C) ERE(A/C) ERE(A/R) 0.500 light 0.960 0.920 0.959 0.971 0.923 0.951 0.500 heavy 0.995 0.750 0.753 1.025 0.751 0.733 0.826 light 0.939 0.868 0.924 0.973 0.847 0.870 0.826 heavy 0.996 0.679 0.682 1.019 0.682 0.669 1.000 light 0.930 0.840 0.904 0.951 0.827 0.870 1.000 heavy 0.995 0.651 0.654 1.021 0.627 0.614 1.500 light 0.913 0.776 0.850 0.927 0.772 0.833 1.500 heavy 0.990 0.597 0.603 1.000 0.591 0.590 2.000 light 0.905 0.730 0.806 0.917 0.731 0.797 2.000 heavy 0.984 0.567 0.576 1.002 0.573 0.572

Censoring	κ	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
Light	0.500	0.962	0.922	0.959	0.970	0.914	0.942
Light	0.826	0.942	0.871	0.924	0.948	0.893	0.942
Light	1.000	0.934	0.844	0.904	0.953	0.838	0.879
Light	1.500	0.919	0.781	0.850	0.936	0.779	0.832
Light	2.000	0.914	0.737	0.806	0.917	0.738	0.805
Heavy	0.500	0.996	0.750	0.753	1.016	0.737	0.725
Heavy	0.826	0.998	0.680	0.682	1.024	0.671	0.656
Heavy	1.000	0.998	0.653	0.654	1.025	0.643	0.628
Heavy	1.500	0.994	0.599	0.603	1.015	0.577	0.569
Heavy	2.000	0.991	0.570	0.576	1.029	0.536	0.521

Table 4: ARE and ERE under two for monotone non-PH patterns, Tian DGP. Parameters: $\kappa = 0.826$, ; $\tau = 40$, $S_0(40) = 0.50$, $s = 1$$t_M = 43$, $N = 400$ $n = 400$ per arm, $B = 2000$.Rows: light (L) and heavy (H) censoring.

Case Cens ARE(R/C) ARE(A/C) ARE(A/R) ERE(R/C) ERE(A/C) ERE(A/R) Case 2 (fading) L 1.347 0.686 0.510 1.535 0.621 0.404 Case 2 (fading) H 1.021 0.384 0.376 1.102 0.318 0.289 Case 3 (delayed) L 0.615 1.169 1.901 0.634 1.171 1.849 Case 3 (delayed) H 0.952 1.334 1.402 0.959 1.435 1.496

Pattern	Cens.	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
fading	Light	1.108	0.741	0.669	1.115	0.711	0.637
fading	Heavy	1.011	0.499	0.494	1.040	0.472	0.454
delayed	Light	0.626	1.190	1.901	0.635	1.187	1.869
delayed	Heavy	0.962	1.349	1.402	0.974	1.374	1.411

5 CheckMate 057: Analysis and Sensitivity

In this section, we illustrate the practical performance of three two-sample tests using data from the CheckMate 057 trial [Borghaei et al., 2015]. To this end, we fit Weibull model to the trial data, compute the theoretical ARE for each test, and conduct simulations to compute the empirical power and required sample sizes under the fitted parametric model.

5.1 Trial background and survival model

The CheckMate 057 data were introduced in Section 1; . The Schoenfeld global test yields $p = 0.001$$p = 0.002$, providing strong evidence against the PH assumption. We therefore fit separate Weibull models to each arm and obtain:

$$\text{Docetaxel (arm 0):} \quad \kappa_0 = 1.264, \ \eta_0 = 13.076 \text{ mo},$$
$$\text{Nivolumab (arm 1):} \quad \kappa_1 = 0.987, \ \eta_1 = 17.320 \text{ mo}.$$

$$\text{Docetaxel (arm 0):} \quad \kappa_0 = 1.264, \ \eta_0 = 13.076 \text{ mo},$$
$$\text{Nivolumab (arm 1):} \quad \kappa_1 = 0.987, \ \eta_1 = 17.320 \text{ mo}.$$

Figure 5 overlays the Kaplan–Meier curves with the fitted Weibull curves, demonstrating their close agreement.

In the docetaxel arm, $\kappa_0 > 1$ inducing induces a monotone increasing hazard. In ; in the nivolumab arm, $\kappa_1 \approx 1$ indicating indicates an approximately constant hazard. The resulting HR is

$$\frac{h_1(t)}{h_0(t)} \approx \frac{1.206}{t^{0.277}},$$

which decreases from 1.21 at month 1 to 0.50 at month 24. This is a fitted log hazard ratio is time varying: $\alpha(1) = 0.187$, it crosses zero near month 1.97, and it reaches -0.693 at month 24. This pattern is characteristic of immunotherapy and is consistent with the Schoenfeld diagnosis (Figure 5), whose negative residual slope indicates a decreasing log hazard ratio over time.

Choice of the censoring model Patients were enrolled from November 2012 through December 2013(≈13-month accrual window), and the primary database lock occurred on March 18, 2015 [Borghaei et al., 2015], giving minimum and maximum . For the fitted-design simulations, we represent staggered entry and follow-up times of approximately 13 and 28 months respectively. We therefore model administrative censoring as $U \sim \text{Uniform}(13, 28)$ by administrative censoring $U \sim \text{Uniform}(6, 30)$ months. Additionally, a small fraction of patients withdrew consent in the study or were lost to follow-up [Borghaei et al., 2015]; we capture this with light exponential dropout, $E \sim \text{Exp}(\lambda_c)$ with $\lambda_c = 0.005$/month (≈6% annually). The combined censoring time is $C = \min(E, U)$. The D-RMST and RAH tests use $\tau = 24$ All three tests use $\tau = 24$ months, corresponding to the clinically natural two-year overall survival landmark; Cox follow-up is administratively censored at the same horizon.

Table 5 presents the ARE and ERE results. Both measures yield the same ordering of the three tests:

$$\text{RAH} \geq \text{Cox} \geq D\text{-RMST}.$$

Figure 6 presents the compass diagram for the CheckMate 057 data. The corresponding squared cosine similarities are

$$\cos^2(\theta_{\text{Cox}}) = 0.603, \quad \cos^2(\theta_{D\text{-RMST}}) = 0.462, \quad \cos^2(\theta_{\text{RAH}}) = 0.638.$$
$$\cos^2(\theta_{\text{Cox}}) = 0.552, \quad \cos^2(\theta_{\text{D-RMST}}) = 0.506, \quad \cos^2(\theta_{\text{RAH}}) = 0.662.$$

Because the ARE between two tests is given by the ratio of their squared cosine similarities, these values imply

$$\text{ARE(RAH/Cox)} \approx 1.058 \quad \text{and} \quad \text{ARE(RAH/}D\text{-RMST)} \approx 1.381.$$
$$\text{ARE(RAH/Cox)} = 1.199 \quad \text{and} \quad \text{ARE(RAH/D-RMST)} = 1.310.$$

Figure 6 also provides a geometric explanation for why the crossing-hazards alternative favors the RAH-based test. The signal direction places substantial weight on the late follow-up period, where the treatment benefit of nivolumab is greatest. In contrast, the Cox test

direction places relatively greater weight on earlier times and gradually decays over follow-up. The D-RMST test direction tapers more rapidly to zero as time approaches τ, thereby placing even less weight on the late follow-up period. The RAH test direction, by comparison, is bounded below by $S_0(\tau) \approx 0.12 S_0(\tau) \approx 0.12$, allowing it to retain meaningful weight at later times. Consequently, the RAH test direction is more closely aligned with the signal direction, resulting in greater efficiency under this crossing-hazards alternative.

Table 5: Theoretical ARE and empirical ERE ($n = 300$ per arm, $B = 500$) under the CheckMate 057 fitted non-PH Weibull DGP. D-RMST and RAH evaluated at ; all methods use $\tau = 24$ months; Cox at $t_H = 26$ months, $n = 300$ per arm, $B = 500$.

Comparison	ARE (theoretical)	ERE ($n = 300$, empirical)
D-RMST$_{vs}$ /Cox	0.906 0.915	0.883 0.935
RAH$_{vs}$ /Cox	1.178 1.199	1.232 1.332
RAH$_{vs}$ /D-RMST	1.300 1.310	1.395 1.425

Beyond the efficiency metrics themselves, a key practical question is how differences in ARE translate into differences in power and required sample size. Figure 7 shows the empirical power as a function of sample size per arm, with the RAH-based test consistently achieving the highest power, followed by the Cox model-based test and then the D-RMST-based test. Table 6 summarizes these differences in two ways: power at a fixed sample size of power at $n_0 = n_1 = 300$ per arm and the minimum sample size required to achieve first simulated sample size on the prespecified 50-patient grid to reach 80% power. The RAH-based test achieves 80% power at approximately 250 RAH reaches the target at 300 patients per arm, compared with approximately 350 400 for Cox and 400 patients per arm for the Cox model- and D-RMST-based tests, respectively400 for D-RMST. Thus, relative to the RAH-based test, the Cox model- and D-RMST-based tests require approximately 40% and 60Cox and D-RMST require 33% and 33% more patients than RAH, respectively, to achieve the same 80% power. Conversely, at a fixed sample size of $n_0 = n_1 = 300$ on this grid. At 300 patients per arm, the empirical power is approximately 85% for the RAH-based test, 76% for the Coxmodel-based test, and only 67% for the D-RMST-based test84% for RAH, 72% for Cox, and 68% for D-RMST.

Table 6: Empirical power at $N = 300$ $n = 300$ per arm and first simulated sample size required for reaching 80% power under the CheckMate 057 Weibull DGP (50-patient grid, $B = 2000$replications), $\tau = 24$ monthswith simulated, uniform censoring $[6, 30]$ monthsARE-, dropout rate 0.005)and ERE-implied ratios relative to RAH.

Method	Power ($N = 300$/arm)	N for 80% power n_{80}	Sim. ratio	ARE ratio	ERE ratio
Cox	0.756 0.717	350 400	1.33	1.20	1.33
D-RMST	0.688 0.680	400	1.33	1.31	1.43
RAH	0.848 0.843	250 300	1.00	1.00	1.00

We now assess the robustness of the ordering RAH $\geq$ Cox $\geq$ D-RMST to the analysis horizon τ, dropout rate λ_c, and assumed alternative. The results are reported in Table 7.

1. **Horizon τ:** ARE(D-RMST/Cox) increases substantially with τ, from 0.531 0.651 at $\tau = 18$ to 0.766 0.915 at $\tau = 24$. The relatively low efficiency of D-RMST at shorter

horizons reflects the truncation of the follow-up period, which ignores a substantial portion of the late treatment benefitof nivolumab. In contrast, ARE(RAH/Cox) remains relatively stable across horizons (0.964–1.110), consistent with the robustness provided by the positive weight floor of RAH. is comparatively stable (1.198, 1.211, and 1.199 at 18, 21, and 24 months), while ARE(RAH/D-RMST) exceeds 1.381 across all cases, ranging from 1.30 at $\tau = 24$ to 1.82 at $\tau = 18$decreases from 1.841 to 1.310.

2. **Dropout rate:** Increasing λ_c from 0.001 to 0.020 shifts ARE(D-RMST/Cox) upward from 0.87 to 1.03from 0.884 to 1.039, indicating that heavier dropout reduces the relative efficiency of the Coxmodel-based test. This is consistent with the Cox weight function being increasingly attenuated at later times as the censoring survival function $S_C(t)$ decreases. In contrast, Cox. ARE(RAH/Cox) remains relatively stable (1.21–1.32). Because the RAH weight floor is not directly affected by censoring, RAH maintains its alignment with the late-concentrated treatment signal as dropout increases . increases modestly from 1.180 to 1.265, whereas ARE(RAH/D-RMST) declines slightly, from 1.32 to 1.21, under heavier censoring. decreases from 1.335 to 1.217. This pattern reflects the growing attenuation of the Cox weight at later times as $S_C(t)$ decreases.

3. **PH vs. non-PH:** Under a PH alternative (HR= 0.75, with the same Weibull shape as for docetaxel), the Cox model-based test Cox is the most efficient, with ARE(D-RMST/Cox)= 1.00 = 0.995 and ARE(RAH/Cox)= 0.87. Both RAH and D-RMST are slightly less efficient than Cox under the PH alternative, highlighting that the efficiency advantage of RAHin the = 0.870. This highlights that RAH's advantage in the fitted non-PH setting arises from the crossing-hazards structure. This finding is consistent with the geometric interpretation: RAH's weight floor improves its greater alignment with the treatment-effect signal when the signal is concentrated in the late follow-up period, but late treatment benefit; the positive weight floor provides no such advantage when the treatment effect follows a PHpatternunder PH.

In summary, the ERE and ARE results are generally consistent. Under the assumed non-PH alternative, the RAH-based test is more efficient than both Cox model-based and D-RMST-based tests. Under the PH alternative, however, the RAH-based test is slightly less efficient than the Cox model-based test.

6 Discussion

This paper has two main contributions: first, to characterize makes three contributions: it characterizes the efficiency of the recently proposed RAH-based test, whose large-sample power properties have not been systematically studied; and second, to develop a geometric interpretation for comparing the efficiency of the Cox model- provides a cosine metric that places the Cox-, D-RMST-, and RAH-based tests on a common geometric scale, and shows how censoring and accrual shape which test is preferred at the design stage. The main message is that no single test uniformly dominates; rather, the most efficient test depends jointly on the underlying pattern of treatment effect and the censoring distribution. The Cox model-based test is most efficient under the PH assumption. The D-RMST-based test is particularly well suited for detecting fading treatment effects, as it places greater weight on early survival differences. In contrast, the RAH-based test is particularly attractive for detecting delayed treatment benefits, a pattern that has been repeatedly observed in studies of immunotherapy in oncology. In addition, the censoring

Table 7: Sensitivity of pairwise ARE (theoretical) and ERE ($n = 300$ per arm, $B = 500$) to analysis horizon, dropout rate, and hazard structure. The ; the common baseline scenario (is $\tau = 24$, $\lambda_c = 0.005$, fitted non-PH) matches Table 5 on ARE, ERE differs by Monte-Carlo error$n = 300$ per arm and $B = 500$.

ARE (theoretical) ERE (empirical) (lr)3-5(lr)6-8Sensitivity Setting $\frac{\text{D−RMST}}{\text{Cox}}$ $\frac{\text{RAH}}{\text{Cox}}$ $\frac{\text{RAH}}{\text{D−RMST}}$ $\frac{\text{D−RMST}}{\text{Cox}}$ $\frac{\text{RAH}}{\text{Cox}}$ $\frac{\text{RAH}}{\text{D−RMST}}$ $\tau = 18$ mo 0.531 0.964 1.817 0.451 0.943 2.091 $\tau = 21$ mo 0.731 1.110 1.519 0.676 1.145 1.694 $\tau = 24$ mo 0.906 1.178 1.300 0.886 1.221 1.379 $\lambda_c = 0.001$ 0.874 1.158 1.324 0.834 1.213 1.455 $\lambda_c = 0.005$ 0.906 1.178 1.300 0.882 1.210 1.371 $\lambda_c = 0.020$ 1.032 1.246 1.207 1.058 1.461 1.381 Non-PH (fitted) 0.906 1.178 1.300 0.874 1.295 1.483 PH (HR = 0.75) 0.999 0.870 0.871 0.996 0.862 0.865

Panel	Setting	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
Horizon	$\tau = 18$	0.651	1.198	1.841	0.607	1.292	2.127
Horizon	$\tau = 21$	0.789	1.211	1.533	0.768	1.299	1.691
Horizon	$\tau = 24$	0.915	1.199	1.310	0.935	1.332	1.425
Dropout	$\lambda_c = 0.001$	0.884	1.180	1.335	0.895	1.223	1.367
Dropout	$\lambda_c = 0.005$	0.915	1.199	1.310	0.935	1.332	1.425
Dropout	$\lambda_c = 0.020$	1.039	1.265	1.217	1.071	1.369	1.278
Hazard	Non-PH (fitted)	0.915	1.199	1.310	0.935	1.332	1.425
Hazard	PH (HR= 0.75)	0.995	0.870	0.875	0.990	0.904	0.913

and accrual designs can substantially affect the relative performance of these tests. These effects can be readily visualized through the proposed cosine metrics, which quantify the alignment between the underlying treatment-effect signal and the directions targeted by each test. In practice, the cosine metric is best used to compare candidate tests across a range of plausible scenarios rather than to commit to one in advance in the study design.

Several extensions are natural. The cosine metric could be used as an objective function for optimizing accrual and follow-up duration, rather than merely as a diagnostic for comparing prespecified tests. In addition, because W_{RAH} is not a Fleming–Harrington weight [Fleming and Harrington, 1991], incorporating it alongside W_{Cox} and $W_{D\text{-RMST}}$ in a Max-combo test would broaden the class of treatment-effect alternatives that can be effectively detected.

A Asymptotic Distribution of the RAH Statistic

A.1 Local-alternative drift

Throughout we work under *Pitman contiguous alternatives*:

$$H_{An}: \quad \log \frac{\lambda_{1n}(t)}{\lambda_0(t)} = \frac{\alpha(t)}{\sqrt{n}}, \tag{18}$$

where $\alpha(\cdot)$ is a fixed function on $(0, \tau]$. Under (18), to first order in $n^{-1/2}$: $\sqrt{n}[S_{1n}(t) - S_0(t)] \to -S_0(t) \int_0^t \lambda_0(u)\alpha(u)\, du$.

Proposition 1. *Under H_{An}, $\sqrt{n}\log(\hat{\theta}_{\text{RAH}})$ is asymptotically normal with mean*

$$\mu_{\text{RAH}} = \int_0^\tau \left[\frac{1}{F_0(\tau)} - \frac{R_0(t)}{R_0(\tau)}\right] \alpha(t)\lambda_0(t)\, dt. \tag{19}$$

Proof. $\sqrt{n}\log(\hat{\theta}_{\mathrm{RAH}}) = U_n + V_n$, where $U_n = \sqrt{n}[\log \hat{F}_1(\tau) - \log \hat{F}_0(\tau)]$ and $V_n = -\sqrt{n}[\log \hat{R}_1(\tau) - \log \hat{R}_0(\tau)]$. Next, we have

$$\begin{aligned} U_n &= F_0(\tau)^{-1}\sqrt{n}[\hat{F}_1 - \hat{F}_0](\tau) + o_p(1) \\ &= F_0(\tau)^{-1}\sqrt{n}[F_1 - F_0](\tau) - F_0(\tau)^{-1}\sqrt{n}[(\hat{S}_1 - S_1) - (\hat{S}_0 - S_0)](\tau) + o_p(1) \\ &= \int_0^\tau \frac{S_0(\tau)}{F_0(\tau)}\alpha(t)\lambda_0(t)\,dt + \eta_{1n} + o_p(1), \end{aligned}$$

and

$$\begin{aligned} V_n &= -R_0(\tau)^{-1}\int_0^\tau \sqrt{n}[\hat{S}_1(t) - \hat{S}_0(t)]\,dt + o_p(1) \\ &= -R_0(\tau)^{-1}\int_0^\tau \sqrt{n}[S_1(t) - S_0(t)]\,dt + R_0(\tau)^{-1}\int_0^\tau \sqrt{n}\left[(\hat{S}_1 - S_1) - (\hat{S}_0 - S_0)\right](t)\,dt + o_p(1) \\ &= R_0(\tau)^{-1}\int_0^\tau S_0(t)A(t)\,dt + \eta_{2n} + o_p(1), \end{aligned}$$

$$\begin{aligned} \mathsf{V_n} &= -\mathsf{R_0}(\tau)^{-1}\int_0^\tau \sqrt{\mathsf{n}}[\hat{\mathsf{S}}_1(\mathsf{t}) - \hat{\mathsf{S}}_0(\mathsf{t})]\,\mathsf{dt} + \mathsf{o_p}(1) \\ &= -\mathsf{R_0}(\tau)^{-1}\int_0^\tau \sqrt{\mathsf{n}}[\mathsf{S}_1(\mathsf{t}) - \mathsf{S}_0(\mathsf{t})]\,\mathsf{dt} - \mathsf{R_0}(\tau)^{-1}\int_0^\tau \sqrt{\mathsf{n}}\left[(\hat{\mathsf{S}}_1 - \mathsf{S}_1) - (\hat{\mathsf{S}}_0 - \mathsf{S}_0)\right](\mathsf{t})\,\mathsf{dt} + \mathsf{o_p}(1) \\ &= \mathsf{R_0}(\tau)^{-1}\int_0^\tau \mathsf{S_0(t)A(t)}\,\mathsf{dt} + \eta_{2n} + \mathsf{o_p}(1), \end{aligned}$$

where $A(t) = \int_0^t \lambda_0(u)\alpha(u)\,du$ and $(\eta_{1n}, \eta_{2n})'$ converging in distribution to a mean zero bivariate Gaussian distribution as $n \to \infty$. Furthermore,

$$R_0(\tau)^{-1}\int_0^\tau S_0(t)A(t)\,dt = A(\tau) - \int_0^\tau \lambda_0(t)\alpha(t)\frac{R_0(t)}{R_0(\tau)}\,dt = \int_0^\tau \left[1 - \frac{R_0(t)}{R_0(\tau)}\right]\alpha(t)\lambda_0(t)dt.$$

Substituting and simplifying gives (19). □

A.2 Pitman efficacy of RAH-based test

The Pitman efficacy formula follows from $\xi_{\mathrm{RAH}} = \mu^2_{\mathrm{RAH}}/\sigma^2_{\mathrm{RAH}}$:

$$\begin{aligned} \xi_{\mathrm{RAH}} &= \frac{\left[\int_0^\tau \left\{\frac{1}{F_0(\tau)} - \frac{R_0(t)}{R_0(\tau)}\right\}\alpha(t)\lambda_0(t)\,dt\right]^2}{\int_0^\tau \left\{\frac{1}{F_0(\tau)} - \frac{R_0(t)}{R_0(\tau)}\right\}^2 \frac{\lambda_0(t)}{w(t)}\,du}. \\ \mathsf{\xi_{RAH}} &= \frac{\left[\int_0^\tau \left\{\frac{1}{\mathsf{F_0}(\tau)} - \frac{\mathsf{R_0(t)}}{\mathsf{R_0}(\tau)}\right\}\alpha(\mathsf{t})\lambda_0(\mathsf{t})\,\mathsf{dt}\right]^2}{\int_0^\tau \left\{\frac{1}{\mathsf{F_0}(\tau)} - \frac{\mathsf{R_0(t)}}{\mathsf{R_0}(\tau)}\right\}^2 \frac{\lambda_0(\mathsf{t})}{\mathsf{w(t)}}\,\mathsf{dt}}. \end{aligned} \tag{20}$$

The efficacies of Cox and D-RMST follow from Tian et al. [2018]:

$$\xi_{\text{Cox}} = \frac{\left[\int_0^\tau w(t)\,\alpha(t)\lambda_0(t)\,dt\right]^2}{\int_0^\tau w(t)\,\lambda_0(t)\,dt}, \tag{21}$$

$$\xi_{D\text{-RMST}} = \frac{\left[\int_0^\tau \{R_0(\tau) - R_0(t)\}\,\alpha(t)\lambda_0(t)\,dt\right]^2}{\int_0^\tau \{R_0(\tau) - R_0(t)\}^2\,\dfrac{\lambda_0(t)}{w(t)}\,dt}. \tag{22}$$

B Derivation of the Weight Function Representations

We prove that each of the three statistics of Section 2.2 is asymptotically equivalent to a version of weighted logrank statistic (7),

$$\int_0^\tau W(t)\,d[\hat{\Lambda}_1(t) - \hat{\Lambda}_0(t)].$$

All derivations use: (i) the KM linear representation

$$\sqrt{n}(\hat{S}_k - S_k) = -S_k\sqrt{n}(\hat{\Lambda}_k - \Lambda_k) + o_p(1);$$

(ii) the functional delta method; and (iii) the local-alternative approximation

$$S_{1n}(t) = S_0(t) - \frac{S_0(t)A(t)}{\sqrt{n}} + o(n^{-1/2}).$$

For simplicity, we also assume equal allocation and common censoring $S_C(\cdot)$.

- **Cox model-based statistic.** Let $\ell_n(\theta)$ be the log partial likelihood over $[0,\tau]$, $U_n(\theta) = \partial\ell_n(\theta)/\partial\theta$ its score, and $\mathcal{I}_n(\theta) = -\partial^2\ell_n(\theta)/\partial\theta^2$ the observed information. The score at $\theta = 0$ is exactly the log-rank statistic, $n^{-1}U_n(0) = U_{\text{Cox}}$ in (8), and

  $$n^{-1}\mathcal{I}_n(0) \longrightarrow \int_0^\tau w(t)\lambda_0(t)dt = \sigma_\theta^{-2}$$

  in probability, uniformly in a neighbourhood of $\theta = 0$. Since $\hat{\theta} = O_p(n^{-1/2})$ under the local alternatives (18) and $U_n(\hat{\theta}) = 0$, a one-term Taylor expansion of $U_n(\theta)$ about $\theta = 0$ gives

  $$0 = n^{-1}U_n(\hat{\theta}) = U_{\text{Cox}} - n^{-1}\mathcal{I}_n(0)\,\hat{\theta} + o_p(n^{-1/2}) \Rightarrow \sqrt{n}\,\hat{\theta} = \sigma_\theta^2\sqrt{n}\,U_{\text{Cox}} + o_p(1).$$

  Hence $\hat{\theta}$ and U_{LR} have the same limit$\sqrt{n}\,\hat{\theta} = \sigma_\theta^2\sqrt{n}\,U_{\text{Cox}} + o_p(1)$, and Z_{Cox} and the normalized log-rank statistic are asymptotically equivalent. The logrank statistic (8) is already of the weighted-log-rank form, with random weight

  $$\hat{w}_{\text{Cox}}(t) = \frac{Y_1(t)Y_0(t)}{Y_1(t) + Y_0(t)},$$

where $Y_k(t)$ is the size of the risk set of group k at t. By law of large numbers, $n^{-1}\hat{w}_{\text{Cox}}(t)$ converges to $w(t)$, which is $S_0(t)S_C(t)/4$ under equal allocation and common censoring. Therefore, we can set

$$W_{\text{Cox}}(t) = 4w(t) = S_0(t)S_C(t),$$

in (7).

- **D-RMST.** By functional delta-method, we have

$$\begin{aligned}
\sqrt{n}\hat{\theta}_{D\text{-RMST}} &= -\sqrt{n}\int_0^\tau S_0(t)\,\hat{\Delta}_\Lambda(t)\,dt + O(\sqrt{n}\|\hat{\Delta}_\Lambda\|^2) \\
&= -\sqrt{n}\int_0^\tau \int_0^t S_0(t)d\Delta_\Lambda(u)\,dt + O(\sqrt{n}\|\sqrt{\Delta}_\Lambda\|^2) \\
&= -\sqrt{n}\int_0^\tau \left\{\int_u^\tau S_0(t)dt\right\} d\hat{\Delta}_\Lambda(u) + O_p(n^{-1/2}) \\
&= -\sqrt{n}\int_0^\tau \{R_0(\tau) - R_0(t)\}\; d\Delta_\Lambda(t) + o_p(1).
\end{aligned}$$

$$\begin{aligned}
\sqrt{\mathsf{n}}\hat{\theta}_{\text{D-RMST}} &= -\sqrt{\mathsf{n}}\int_0^\tau \mathsf{S}_0(\mathsf{t})\,\hat{\Delta}_\Lambda(\mathsf{t})\,\mathsf{dt} + \mathsf{O}\big(\sqrt{\mathsf{n}}\|\hat{\Delta}_\Lambda\|^2\big) \\
&= -\sqrt{\mathsf{n}}\int_0^\tau \mathsf{S}_0(\mathsf{t})\int_0^{\mathsf{t}} \mathsf{d}\hat{\Delta}_\Lambda(\mathsf{u})\,\mathsf{dt} + \mathsf{O}_\mathsf{p}\big(\mathsf{n}^{-1/2}\big) \\
&= -\sqrt{\mathsf{n}}\int_0^\tau \left\{\int_\mathsf{u}^\tau \mathsf{S}_0(\mathsf{t})\,\mathsf{dt}\right\} \mathsf{d}\hat{\Delta}_\Lambda(\mathsf{u}) + \mathsf{O}_\mathsf{p}\big(\mathsf{n}^{-1/2}\big) \\
&= -\sqrt{\mathsf{n}}\int_0^\tau \{\mathsf{R}_0(\tau) - \mathsf{R}_0(\mathsf{u})\}\,\mathsf{d}\hat{\Delta}_\Lambda(\mathsf{u}) + \mathsf{o}_\mathsf{p}(1).
\end{aligned}$$

  Therefore, we may set the weight as

$$W_{D\text{-RMST}}(t) = \frac{R_0(\tau) - R_0(t)}{R_0(\tau)},$$

  as $Z_{D\text{-RMST}}$ is invariant to multiplying any nonzero constant to the weight function.

- **RAH.** Using the same method, we have

$$W_{\text{RAH}}(t) = -\left[\frac{S_0(\tau)}{F_0(\tau)} + \frac{R_0(\tau) - R_0(t)}{R_0(\tau)}\right].$$

  argument together with the delta method for $\log F_k(\tau)$ and $\log R_k(\tau)$,

$$\log\hat{\theta}_{\text{RAH}} = \int_0^\tau \left[\frac{\mathsf{S}_0(\tau)}{\mathsf{F}_0(\tau)} + \frac{\mathsf{R}_0(\tau) - \mathsf{R}_0(\mathsf{t})}{\mathsf{R}_0(\tau)}\right] \mathsf{d}\hat{\Delta}_\Lambda(\mathsf{t}) + \mathsf{O}_\mathsf{p}(\mathsf{n}^{-1}).$$

  Multiplying $W_{\text{RAH}}(\cdot)$ by $-F_0(\tau)$ yields an equivalent weight

$$W_{\text{RAH}}(t) = S_0(\tau) + F_0(\tau)\,\frac{R_0(\tau) - R_0(t)}{R_0(\tau)} = S_0(\tau) + [1 - S_0(\tau)]\,W_{D\text{-RMST}}(v).$$

  by $F_0(\tau) > 0$ gives the equivalent weight

$$\mathsf{W}_{\text{RAH}}(\mathsf{t}) = \mathsf{S}_0(\tau) + \mathsf{F}_0(\tau)\,\frac{\mathsf{R}_0(\tau) - \mathsf{R}_0(\mathsf{t})}{\mathsf{R}_0(\tau)} = \mathsf{S}_0(\tau) + \big[1 - \mathsf{S}_0(\tau)\big]\,\mathsf{W}_{\text{D-RMST}}(\mathsf{t}).$$

C Simulation Study for PH Setting at HR = 0.82

We repeat the simulation for PH scenarios with HR=0.82. The results are reported in Tables 8 and 9. Both ARE and ERE are similar to to those for HR= 0.70:

Table 8: ARE and ERE under PH (HR= 0.82), $\kappa = 0.826$, $\tau = 40$, $n = 400$ per arm, $B = 2000$.

$S_0(40)$ Censoring ARE(R/C) ARE(A/C) ARE(A/R) ERE(R/C) ERE(A/C) ERE(A/R) 0.10 light 0.992 0.940 0.947 0.990 0.944 0.954 0.10 heavy 0.982 0.797 0.812 1.000 0.836 0.836 0.20 light 0.978 0.915 0.935 0.975 0.882 0.905 0.20 heavy 0.992 0.750 0.756 0.993 0.735 0.741 0.30 light 0.964 0.896 0.929 0.970 0.908 0.936 0.30 heavy 0.996 0.719 0.722 1.006 0.661 0.657 0.40 light 0.951 0.880 0.926 0.945 0.883 0.935 0.40 heavy 0.997 0.697 0.699 1.005 0.699 0.696 0.50 light 0.939 0.868 0.924 0.952 0.850 0.893 0.50 heavy 0.996 0.679 0.682 0.998 0.692 0.693 0.60 light 0.927 0.857 0.924 0.915 0.856 0.935 0.60 heavy 0.995 0.665 0.668 0.998 0.671 0.672 0.70 light 0.916 0.847 0.924 0.928 0.847 0.914 0.70 heavy 0.993 0.653 0.658 0.996 0.643 0.646 0.80 light 0.906 0.838 0.925 0.892 0.848 0.950 0.80 heavy 0.990 0.643 0.649 1.009 0.636 0.631 0.90 light 0.897 0.831 0.927 0.915 0.829 0.906 0.90 heavy 0.987 0.634 0.642 1.020 0.563 0.552

Censoring	$S_0(40)$	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
Light	0.10	0.994	0.941	0.947	1.001	0.939	0.938
Light	0.20	0.980	0.917	0.935	0.987	0.898	0.910
Light	0.30	0.967	0.898	0.929	0.968	0.911	0.942
Light	0.40	0.954	0.883	0.926	0.957	0.872	0.911
Light	0.50	0.942	0.871	0.924	0.929	0.885	0.953
Light	0.60	0.931	0.860	0.924	0.958	0.852	0.889
Light	0.70	0.920	0.851	0.924	0.930	0.839	0.903
Light	0.80	0.911	0.842	0.925	0.952	0.831	0.872
Light	0.90	0.901	0.835	0.927	0.913	0.834	0.914
Heavy	0.10	0.982	0.798	0.812	0.989	0.727	0.735
Heavy	0.20	0.993	0.751	0.756	1.001	0.718	0.717
Heavy	0.30	0.997	0.720	0.722	1.004	0.704	0.701
Heavy	0.40	0.998	0.698	0.699	1.004	0.689	0.686
Heavy	0.50	0.998	0.680	0.682	1.007	0.672	0.667
Heavy	0.60	0.997	0.666	0.668	1.001	0.676	0.675
Heavy	0.70	0.995	0.654	0.658	0.996	0.685	0.688
Heavy	0.80	0.992	0.644	0.649	0.993	0.661	0.665
Heavy	0.90	0.990	0.635	0.642	1.054	0.698	0.662

D Simulation Study for PH Setting with Different Sample Size, Horizon and Survival Distributions

We also repeat simulation study the PH setting under different design: $\tau = 10$, $n_0 = n_1 = 150$, and three baseline distributions (an exponential with $S_0(10) \approx 0.37$; a Weibull baseline with a decreasing hazard ($\kappa = 0.7$) and $S_0(10) = 0.5$; and a Weibull baseline with an increasing hazard ($\kappa = 1.5$) and $S_0(10) = 0.5$.) We also consider light censoring with $C \sim \min\{U(8, 12), \text{Exp}(0.02)\}$ and heavy censoring with $C \sim \min\{U(8, 12), \text{Exp}(0.20)\}$.

The results are reported in Table 10. It is interesting to observe that RAH-based test is RAH is more efficient than D-RMST-based test D-RMST under light censoring(ARE ranges from 1.067 to 1.119), while

Table 9: ARE, empirical power, and ERE under PH (HR= 0.82)with different baseline distributions in the control , $S_0(40) = 0.50$, $\tau = 40$, $n = 400$ per arm, $B = 2000$.

Censoring	κ	ARE(D/C)	ARE(R/C)	ARE(R/D)	Power C	Power D	Power R	ERE(D/C)	ERE(R/C)	ERE(R/D)
Light	0.500	0.962	0.922	0.959	0.458	0.452	0.433	0.980	0.918	0.937
Light	0.826	0.942	0.871	0.924	0.435	0.418	0.395	0.929	0.885	0.953
Light	1.000	0.934	0.844	0.904	0.448	0.406	0.393	0.941	0.838	0.891
Light	1.500	0.919	0.781	0.850	0.415	0.382	0.341	0.931	0.802	0.862
Light	2.000	0.914	0.737	0.806	0.383	0.349	0.307	0.906	0.765	0.844
Heavy	0.500	0.996	0.750	0.753	0.376	0.379	0.307	1.003	0.744	0.741
Heavy	0.826	0.998	0.680	0.682	0.336	0.337	0.260	1.007	0.672	0.667
Heavy	1.000	0.998	0.653	0.654	0.298	0.301	0.237	0.996	0.632	0.635
Heavy	1.500	0.994	0.599	0.603	0.277	0.284	0.224	0.991	0.632	0.638
Heavy	2.000	0.991	0.570	0.576	0.247	0.253	0.183	0.995	0.584	0.587

D-RMST-based test , whereas D-RMST is substantially more efficient than RAH-based test RAH under heavy censoring(ARE ranges from 0.66 to 0.69).

Table 10: ARE and ERE under PH (HR= 0.70), $\tau = 10$, $n_0 = n_1 = 150$$n = 150$ per arm, and light and heavy censoring$B = 500$.

Baseline Cens ARE(R/C) ARE(A/C) ARE(A/R) ERE(R/C) ERE(A/C) ERE(A/R) Exp(λ_0=0.10) L 0.896 0.958 1.068 0.891 0.960 1.079 Exp(λ_0=0.10) H 0.968 0.662 0.684 0.966 0.523 0.542 Weibull(κ=0.7) L 0.906 0.966 1.067 0.932 0.959 1.029 Weibull(κ=0.7) H 0.966 0.666 0.690 0.982 0.628 0.640 Weibull(κ=1.5) L 0.823 0.921 1.119 0.737 0.950 1.289 Weibull(κ=1.5) H 0.975 0.643 0.659 0.985 0.592 0.601

Censoring	Baseline	ARE(D/C)	ARE(R/C)	ARE(R/D)	ERE(D/C)	ERE(R/C)	ERE(R/D)
Light	Weibull(kappa=0.7)	0.920	0.981	1.067	0.985	0.974	0.989
Light	Exponential(lambda0=0.10)	0.913	0.976	1.068	0.941	0.989	1.051
Light	Weibull(kappa=1.5)	0.853	0.954	1.119	0.895	0.906	1.013
Heavy	Weibull(kappa=0.7)	0.970	0.669	0.690	1.025	0.722	0.704
Heavy	Exponential(lambda0=0.10)	0.973	0.665	0.684	1.033	0.670	0.648
Heavy	Weibull(kappa=1.5)	0.987	0.651	0.659	0.976	0.611	0.626

E Additional Simulations: Crossing Hazards

In this simulation, $\alpha(t) = \log(0.46) + (t/t_M)$ with $t_M = 43$ months, $\kappa = 0.826$, and $S_0(40) = 0.50$: the HR starts at 0.46, crosses one, and rises above 1 late in follow-up. Thus, an early treatment benefit is reversed to harm at late times. Table 11 reports the ARE and ERE under light and heavy censoring. Both RAH D-RMST is more efficient than Cox, whereas RAH is less efficient than both Cox and D-RMST-based tests exceed Cox model-based test (ARE and ERE above 1). The ordering and magnitudes are similar across censoringlevelsD-RMST, under both censoring regimes. The efficiency separation is larger under heavy censoring.

Table 11: ARE and ERE under the crossing-hazard scenario. $\alpha(u) = \log(0.46) + (u/t_M)$. $\kappa = 0.826$crossing hazards; $t_M = 43$, $\tau = 40$, $S_0(40) = 0.50$, and $n_0 = n_1 = 400$ $n = 400$ per arm, $B = 2000$.

CensCens.	ARE(RD/C)	ARE(AR/C)	ARE(A/RR/D)	ERE(RD/C)	ERE(AR/C)	ERE(A/RR/D)
L Light	1.040 1.193	1.072 0.683	1.031 0.573	1.027 1.272	1.075 0.599	1.047 0.471
H Heavy	1.041 1.016	1.053 0.429	1.012 0.422	1.021 1.060	1.070 0.374	1.048 0.353

References

Hossein Borghaei, Luis Paz-Ares, Leora Horn, David R. Spigel, Martin Steins, Neal E. Ready, Laura Q. Chow, Everett E. Vokes, Enriqueta Felip, Esther Holgado, et al. Nivolumab versus docetaxel in advanced nonsquamous non–small-cell lung cancer. *New England Journal of Medicine*, 373(17):1627–1639, 2015. doi: 10.1056/NEJMoa1507643.

D. R. Cox. Regression models and life-tables. *Journal of the Royal Statistical Society, Series B*, 34(2):187–220, 1972.

Thomas R. Fleming and David P. Harrington. *Counting Processes and Survival Analysis*. Wiley, New York, 1991.

Patricia Guyot, A. E. Ades, Mario J. N. M. Ouwens, and Nicky J. Welton. Enhanced secondary analysis of survival data: Reconstructing the data from published kaplan-meier survival curves. *BMC Medical Research Methodology*, 12:9, 2012. doi: 10.1186/1471-2288-12-9.

Miguel A. Hernán. The hazards of hazard ratios. *Epidemiology*, 21(1):13–15, 2010. doi: 10.1097/EDE.0b013e3181c1ea43.

Miki Horiguchi, Michael J. Hassett, and Hajime Uno. How do the accrual pattern and follow-up duration affect the hazard ratio estimate when the proportional hazards assumption is violated? *The Oncologist*, 24(7):867–871, 2019. doi: 10.1634/theoncologist.2018-0141.

Ray S. Lin, Ji Lin, Satrajit Roychoudhury, Keaven M. Anderson, Tianle Hu, Bo Huang, Larry J. Leon, Jason J.Z. Liao, Rong Liu, Xiaodong Luo, Pralay Mukhopadhyay, Rui Qin, Kay Tatsuoka, Xuejing Wang, Yang Wang, Jian Zhu, Tai-Tsang Chen, and Rinaldo Iacona. Alternative analysis methods for time to event endpoints under nonproportional hazards: A comparative analysis. *Statistics in Biopharmaceutical Research*, 12(2):187–198, 2020. doi: 10.1080/19466315.2019.1697804.

Xiang Meng, Lu Tian, Kenneth Kehl, and Hajime Uno. Efficient and debiased learning of average hazard under non-proportional hazards. *arXiv preprint arXiv:2602.13475*, 2026.

E. J. G. Pitman. Notes on non-parametric statistical inference. Technical Report 27, North Carolina State University, Department of Statistics, 1949. URL `https://www.lib.ncsu.edu/resolver/1840.4/2430`. Repository date listed as 1949?; notes based on lectures at Columbia University.

Patrick Royston and Mahesh K. B. Parmar. Restricted mean survival time: An alternative to the hazard ratio for the design and analysis of randomized trials with a time-to-event outcome. *BMC Medical Research Methodology*, 13:152, 2013. doi: 10.1186/1471-2288-13-152.

Lu Tian, Lihui Zhao, and L. J. Wei. Predicting the restricted mean event time with the subject's baseline covariates in survival analysis. *Biostatistics*, 15(2):222–233, 2014. doi: 10.1093/biostatistics/kxt050.

Lu Tian, Haoda Fu, Stephen J. Ruberg, Hajime Uno, and Lee-Jen Wei. Efficiency of two sample tests via the restricted mean survival time for analyzing event time observations. *Biometrics*, 74(2):694–702, 2018. doi: 10.1111/biom.12770.

Hajime Uno and Miki Horiguchi. Ratio and difference of average hazard with survival weight: New measures to quantify survival benefit of new therapy. *Statistics in Medicine*, 42(7): 936–952, 2023. doi: 10.1002/sim.9651.

Hajime Uno, Brian Claggett, Lu Tian, Eisuke Inoue, Paul Gallo, Toshio Miyata, Deborah Schrag, Masahiro Takeuchi, Yoshiaki Uyama, Lihui Zhao, Hicham Skali, Scott Solomon, Susanna Jacobus, Michael Hughes, Milton Packer, and Lee-Jen Wei. Moving beyond the hazard ratio in quantifying the between-group difference in survival analysis. *Journal of Clinical Oncology*, 32(22):2380–2385, 2014. doi: 10.1200/JCO.2014.55.2208.

Hajime Uno, Lu Tian, Miki Horiguchi, Angel Cronin, Chakib Battioui, and James Bell. *survRM2: Comparing Restricted Mean Survival Time*, 2022. URL `https://CRAN.R-project.org/package=survRM2`. R package version 1.0-4.




Figure 1: Unit-vector functions (left column) and compass diagrams (right column) for three scenarios (Weibull baseline: $\kappa = 0.826$, $S_0(40) = 0.50$, light censoring). The D-RMST and RAH tests are evaluated at $\tau = 40$ months; Signal direction is placed at 12 o'clock in the compass diagrams; each test arrow is annotated with its $\cos^2(\theta_X)$ value. Under PH (top), Cox is aligned, while D-RMST and RAH lose some efficiencies. Under delayed benefit (bottom), the signal rotates toward the RAH direction, raising $\cos^2(\theta_{\text{RAH}})$ above $\cos^2(\theta_{\text{Cox}})$.




Figure 2: **PH scenario** (HR $= 0.70$) and $S_0(\tau) \in \{0.10, 0.30, 0.70\}$. The HR is flat at 0.70; higher event rates produce greater absolute separation between two survival curves.




Figure 3: **Case 2: fading benefit.** $\alpha(u) = -c_0 \max(t_M - u, 0)^s$, mean log-HR $= \log(0.70)$. The HR is below 1 early and converges to 1 by t_M; larger s concentrates the benefit closer to $u = 0$.




Figure 4: **Case 3: delayed benefit.** $\alpha(u) = -c_0\, u^s$, mean log-HR $= \log(0.70)$. The HR starts at 1 and decreases; larger s delays the onset of the treatment effect, with the benefit concentrated at later time points for $s = 2$.




Figure 5: KM curves Survival model fit (solidleft) and fitted Weibull models proportional-hazards diagnostic (dashedright) for docetaxel (black) and nivolumab (red)the CheckMate 057 trial.Parameters estimated by maximum likelihood.

Scaled Schoenfeld residuals for the treatment indicator. The negative slope (Schoenfeld global test $p = 0.001$) indicates a time-varying hazard ratio: the nivolumab advantage grows over follow-up. Survival model fit (left) and proportional-hazards diagnostic (right) for the CheckMate 057 trial.




Figure 6: Unit-vector functions (left) and compass diagram (right) for Weibull distributions fitted to CheckMate 057 trial. D-RMST and RAH All three tests use $\tau = 24$ $\tau = 24$ months; the function panel displays them only on $[0, \tau]$. The signal direction is placed at 12 o'clock and reflects the crossing-hazards pattern. Each test direction is annotated with $\cos^2(\theta_X)$. RAH is closest to the signal direction, followed by Cox and D-RMST.




Figure 7: Empirical power for Cox model-(bluered, solid), D-RMST- D-RMST (orangeblue, dashed), and RAH- RAH (green, dotted) for under the fitted CheckMate 057 non-PH alternativeobserved in CheckMate 057. . Horizontal line: 80% power.