

jaxparrow: a Python package solving the cyclogeostrophic balance using a variational formulation

Vadim BERTRAND^{1*}, Victor E V Z DE ALMEIDA^{1*}, Julien LE
SOMMER^{1*}, and Emmanuel COSME^{1*}

¹ Université Grenoble Alpes, France * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

Summary

Sea Surface Height (SSH) variations measured by satellite altimeters are widely used to estimate Sea Surface Currents (SSC) in oceanographic operational or research applications. The geostrophic balance approximation, which relates the pressure gradient, the current velocity, and the Coriolis force, is commonly employed to estimate SSC from SSH. It is known that under some configurations, the velocity advection term, neglected in the geostrophic formulation, should be included in the balance, leading to the cyclogeostrophic balance approximation. In general, solving the cyclogeostrophic balance can not be done analytically and numerical methods are needed. However, (1) existing iterative approaches are known to diverge, and ad-hoc methods are used to avoid local discontinuities; (2) publicly available, well maintained implementations are missing.

To overcome these limitations, we propose the Python package `jaxparrow`. `jaxparrow` formulates the cyclogeostrophic balance as a variational problem and solve it using a collection of well known optimizers. Its implementation heavily relies on JAX, the Python library bringing together automatic differentiation and just-in-time compilation, and the growing ecosystem around it. `jaxparrow` can be used as a package for an easy integration to existing oceanographic pipelines, or as a standalone executable working directly with NetCDF files.

Statement of need

Sea Surface Currents (SSC) can be easily approximated from satellite altimetry observations of the Sea Surface Height (SSH) using the geostrophic balance. Geostrophy describes the balance between the pressure gradient force (indirectly observed via SSH), and the Coriolis force. Geostrophic currents satisfy this equilibrium:

$$f(\vec{k} \times \vec{u}_g) = -g\nabla\eta, \quad (1)$$

where f is the Coriolis parameter, \vec{k} the vertical unit vector, \vec{u}_g the geostrophic velocity, g the gravity, and η the SSH.

The geostrophic equation represents a drastic approximation of the Navier-Stokes equation adapted to ocean dynamics. However, as discussed by Bakun (2006), Charney (1955), and Maximenko & Niiler (2006), geostrophy alone is not always sufficient to accurately estimate SSC. In particular, Penven et al. (2014) showed that, in the highly energetic Mozambique Channel, the geostrophic approximation can produce errors of the order of 30% in velocity estimates, and that the advection term $\vec{u} \cdot \nabla \vec{u}$ was needed in the balance to reduce these

errors. Considering a horizontal, stationary, and inviscid flow, the momentum equation linking SSC velocities \vec{u} with SSH —through geostrophic velocities \vec{u}_g from Equation 1— can be expressed as:

$$\vec{u}_c - \frac{\vec{k}}{f} \times (\vec{u}_c \cdot \nabla \vec{u}_c) = \vec{u}_g, \quad (2)$$

where \vec{u}_c is the cyclogeostrophic velocity.

Ocean data and services providers, such as Copernicus Marine Environment Monitoring Service (Taburet et al., 2019), use geostrophic balance to estimate SSC from SSH. Cao et al. (2023) demonstrates that applying cyclogeostrophic corrections to the global ocean over a 25-years period results in significantly different estimates of SSC. Ocean products could therefore greatly benefit from a robust and open estimation method of cyclogeostrophic currents, which, to our knowledge, is not presently available.

Numerical resolution of the cyclogeostrophic inverse problem

Because of the advective term $\vec{u}_c \cdot \nabla \vec{u}_c$, Equation 2 is nonlinear, and solving it analytically is conceivable only in idealized scenarios, making numerical approaches essential. The current state-of-the-art method to solve the cyclogeostrophic equation is the iterative formulation introduced by Arnason et al. (1962) and Endlich (1961), which consists of reaching balance using the following iterative scheme:

$$\vec{u}_c^{(n+1)} = \vec{u}_g + \frac{\vec{k}}{f} \times (\vec{u}_c^{(n)} \cdot \nabla \vec{u}_c^{(n)}), \quad (3)$$

with $\vec{u}_c^{(0)} = \vec{u}_g$. This approach is known to diverge since Arnason et al. (1962), and in practice (Ioannou et al., 2019; Penven et al., 2014) the residual $res = |\vec{u}_c^{(n+1)} - \vec{u}_c^{(n)}|$ is used to control point by point the iteration process. The iterative procedure is usually stopped when the residual locally falls below 0.01 m/s or starts to increase.

To avoid the local divergence issue of the iterative process, and its ad-hoc control, we propose to formulate the cyclogeostrophy as the variational problem:

$$J(\vec{u}_c) = \left\| \vec{u}_c - \frac{\vec{k}}{f} \times (\vec{u}_c \cdot \nabla \vec{u}_c) - \vec{u}_g \right\|^2, \quad (4)$$

where $\|\cdot\|$ is the discrete L^2 norm. jaxparrow implements this approach, leveraging JAX (Bradbury et al., 2021). Thanks to JAX automatic differentiation capabilities, ∇J is numerically available, and the cyclogeostrophic currents are estimated by minimizing Equation 4 using a gradient-based optimizer, with $\vec{u}_c^{(0)} = \vec{u}_g$ as initial guess.

Application to the Alboran sea

The Alboran sea is an energetic area of the Mediterranean sea. We demonstrate below the need to consider cyclogeostrophy in this region, and the benefit of the variational formulation implemented in jaxparrow. The data and results presented here can be found in the Alboran sea notebook hosted on GitHub.

We use SSH and SSC from the eNATL60 configuration (Brodeau et al., 2020; Uchida et al., 2022) of the state-of-the-art NEMO ocean circulation model (Madec et al., 2022) as reference data. Figure 1 shows SSH, SSC, and normalized relative vorticities in this region.




Figure 1: Reference data: on the left and middle panels, SSH and SSC velocity (colored by the magnitude, with arrows giving the direction) simulated by NEMO; on the right, the corresponding normalized vorticity.

71 Using SSH, jaxparrow can first estimate the geostrophic SSC with Equation 1. As geostrophy
72 is a major mechanism governing ocean dynamics, vorticities derived from those velocities
73 present an overall similarity with the ones obtain from NEMO data. However, we can clearly
74 identify irregular areas (around $(-4, 35.5)$, $(-3, 36)$, and $(-2.5, 35.5)$ in (longitude, latitude)
75 coordinates, see Figure 2) where the geostrophic balance fails to accurately reconstruct SSC.




Figure 2: The qualitative comparison between reference (left panel) and geostrophic (right panel) normalized vorticities reveals several regions with highly erroneous estimations.

76 Starting from geostrophic currents, jaxparrow solves the variational formulation of the cyclo-
77 geostrophy (Equation 4), using in this example the classical gradient descent (Kantorovich &
78 Akilov, 2016). As a result, almost all the problematic areas are now much more accurately
79 reconstructed, leaving mainly costal or domain boundary regions with large differences from our
80 reference vorticity (see Figure 3, left and middle panels). By evaluating the cyclogeostrophic
81 disequilibrium (the functional J in Equation 4) along the optimization process, we observe that
82 jaxparrow iteratively converges towards cyclogeostrophic balance (right panel of Figure 3).




Figure 3: In contrast to the geostrophic approximation, variational cyclogeostrophy (middle panel) provide accurate reconstruction of the reference (left panel) normalized vorticities. The right panel demonstrates the fast convergence towards cyclogeostrophic balance.

83 For comparison, jaxparrow can also estimate the cyclogeostrophic currents using the iterative
84 scheme (Equation 3). In this example, the evolution of J at each iteration reveals that this
85 approach is not able to fill the cyclogeostrophic balance (see Figure 4, right panel). We even
86 notice that this estimation is qualitatively worse than the geostrophy (left and middle panels
87 of Figure 4).




Figure 4: As exhibited in the right panel, the iterative approach diverges from the cyclogeostrophic balance; and we can notice from the two other panels that the resulting normalized vorticity is qualitatively worse than the geostrophic one.

Those qualitative observations are supported by more quantitative analysis. We computed the 1000 first percentiles of the vorticity distributions, and we observe, via a Q-Q plot (Wilk & Gnanadesikan, 1968), that the percentiles of the variational distribution are the closest to the ones of the reference distribution (Figure 5).




Figure 5: The percentiles of the normalized vorticity distributions demonstrate that our variational estimation of the cyclogeostrophy (in orange) corrects the geostrophy approximation (in blue), while the iterative scheme (in green) tends to diverge from the reference (in black).

Availability

Beside the novel variational formulation, jaxparrow also offers the first to our knowledge open implementation of the cyclogeostrophy inversion. The code is available on [GitHub](#), with the specific tag joss for the version matching this publication; and the documentation, with pip-installation instructions, usage examples, and toy notebooks, is hosted on [Read the Docs](#).

References

- Arnason, G., Haltiner, G., & Frawley, M. (1962). Higher-order geostrophic wind approximations. *Monthly Weather Review*, 90(5), 175–185. [https://doi.org/10.1175/1520-0493\(1962\)090%3C0175:HGWA%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1962)090%3C0175:HGWA%3E2.0.CO;2)
- Bakun, A. (2006). Fronts and eddies as key structures in the habitat of marine fish larvae: Opportunity, adaptive response and competitive advantage. *Scientia Marina*, 70(S2), 105–122. <https://doi.org/10.3989/scimar.2006.70s2105>

- 104 Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Leary, C., Maclaurin, D., Necula, G.,
105 Paszke, A., VanderPlas, J., Wanderman-Milne, S., & others. (2021). JAX: Autograd and
106 XLA. *Astrophysics Source Code Library*, ascl-2111. <https://github.com/google/jax>
- 107 Brodeau, L., Sommer, J. L., & Albert, A. (2020). *ocean-next/eNATL60: Material describing*
108 *the set-up and the assessment of NEMO-eNATL60 simulations* (Version v1). Zenodo.
109 <https://doi.org/10.5281/zenodo.4032732>
- 110 Cao, Y., Dong, C., Stegner, A., Bethel, B. J., Li, C., Dong, J., Lü, H., & Yang, J. (2023).
111 Global sea surface cyclogeostrophic currents derived from satellite altimetry data. *Journal*
112 *of Geophysical Research: Oceans*, 128(1), e2022JC019357. [https://doi.org/10.1029/](https://doi.org/10.1029/2022JC019357)
113 [2022JC019357](https://doi.org/10.1029/2022JC019357)
- 114 Charney, J. G. (1955). The gulf stream as an inertial boundary layer. *Proceedings of the*
115 *National Academy of Sciences*, 41(10), 731–740. <https://doi.org/10.1073/pnas.41.10.731>
- 116 Endlich, R. M. (1961). Computation and uses of gradient winds. *Monthly Weather Review*,
117 89(6), 187–191. [https://doi.org/10.1175/1520-0493\(1961\)089%3C0187:CAUOGW%3E2.](https://doi.org/10.1175/1520-0493(1961)089%3C0187:CAUOGW%3E2.0.CO;2)
118 [0.CO;2](https://doi.org/10.1175/1520-0493(1961)089%3C0187:CAUOGW%3E2.0.CO;2)
- 119 Ioannou, A., Stegner, A., Tuel, A., LeVu, B., Dumas, F., & Speich, S. (2019). Cyclostrophic
120 corrections of AVISO/DUACS surface velocities and its application to mesoscale eddies in
121 the mediterranean sea. *Journal of Geophysical Research: Oceans*, 124(12), 8913–8932.
122 <https://doi.org/10.1029/2019JC015031>
- 123 Kantorovich, L. V., & Akilov, G. P. (2016). *Functional analysis*. Elsevier.
- 124 Madec, G., Bourdallé-Badie, R., Chanut, J., Clementi, E., Coward, A., Ethé, C., Iovino, D.,
125 Lea, D., Lévy, C., Lovato, T., Martin, N., Masson, S., Mocavero, S., Rousset, C., Storkey,
126 D., Müller, S., Nurser, G., Bell, M., Samson, G., ... Moulin, A. (2022). *NEMO ocean*
127 *engine* (Version v4.2). Zenodo. <https://doi.org/10.5281/zenodo.6334656>
- 128 Maximenko, N., & Niiler, P. (2006). Mean surface circulation of the global ocean
129 inferred from satellite altimeter and drifter data. *Proceeding of the Symposium*
130 *on 15 Years of Progress in Radar Altimetry*, Eur. Space Agency Spec. Publ.,
131 *ESA SP*, 614. [https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=](https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=3a37ef4ad98374b642a360e1445474bb2bdcafed)
132 [3a37ef4ad98374b642a360e1445474bb2bdcafed](https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=3a37ef4ad98374b642a360e1445474bb2bdcafed)
- 133 Penven, P., Halo, I., Pous, S., & Marié, L. (2014). Cyclogeostrophic balance in the mozambique
134 channel. *Journal of Geophysical Research: Oceans*, 119(2), 1054–1067. [https://doi.org/](https://doi.org/10.1002/2013JC009528)
135 [10.1002/2013JC009528](https://doi.org/10.1002/2013JC009528)
- 136 Taburet, G., Sanchez-Roman, A., Ballarotta, M., Pujol, M.-I., Legeais, J.-F., Fournier, F.,
137 Faugere, Y., & Dibarboure, G. (2019). DUACS DT2018: 25 years of reprocessed sea
138 level altimetry products. *Ocean Science*, 15(5), 1207–1224. [https://doi.org/10.5194/](https://doi.org/10.5194/os-15-1207-2019)
139 [os-15-1207-2019](https://doi.org/10.5194/os-15-1207-2019)
- 140 Uchida, T., Le Sommer, J., Stern, C., Abernathey, R. P., Holdgraf, C., Albert, A., Brodeau,
141 L., Chassignet, E. P., Xu, X., Gula, J., & others. (2022). Cloud-based framework for
142 inter-comparing submesoscale-permitting realistic ocean models. *Geoscientific Model*
143 *Development*, 15(14), 5829–5856. <https://doi.org/10.5194/gmd-15-5829-2022>
- 144 Wilk, M. B., & Gnanadesikan, R. (1968). Probability plotting methods for the analysis for the
145 analysis of data. *Biometrika*, 55(1), 1–17. <https://doi.org/10.1093/biomet/55.1.1>