




Figure 3. Left Ventricular Diastolic Parameters and Cardiopulmonary Fitness. Primary restricted cubic spline models adjusted for age, sex, and BMI, with each diastolic index analyzed in all patients in whom it was measured; panel annotations give the analyzed N. Solid lines represent spline estimates with shaded 95% confidence intervals; dashed vertical lines denote ASE abnormality thresholds; Panel annotations report the analyzed N, overall association P value, and nonlinearity q value. The overall P value tests whether the full spline adds information beyond age, sex, and BMI; the nonlinearity q value tests whether curvature improves fit beyond a straight-line relationship after adjustment across the six primary comparisons. **(A)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of E/e' . **(B)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of LAVi. **(C)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of TRV_{max}. **(D)** $\dot{V}E/\dot{V}CO_2$ slope as a function of E/e' . **(E)** $\dot{V}E/\dot{V}CO_2$ slope as a function of LAVi. **(F)** $\dot{V}E/\dot{V}CO_2$ slope as a function of TRV_{max}.