




Figure 3. Left Ventricular Diastolic Parameters and Cardiopulmonary Fitness. Primary restricted cubic spline models adjusted for age, sex, and BMI in the fixed E/e' + LAVi cohort; TRVmax was analyzed where measured and panel annotations give the analyzed N. Solid lines represent spline estimates with shaded 95% confidence intervals; dashed vertical lines denote ASE abnormality thresholds; Panel annotations report the analyzed N, overall association P value, and nonlinearity q value. The overall P value tests whether the full spline adds information beyond age, sex, and BMI; the nonlinearity q value tests whether curvature improves fit beyond a straight-line relationship after adjustment across the six primary comparisons. **(A)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of E/e' . **(B)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of LAVi. **(C)** Peak $\dot{V}O_2$ (FRIEND 2.0 % predicted) as a function of TRV_{max}. **(D)** $\dot{V}E/\dot{V}CO_2$ slope as a function of E/e' . **(E)** $\dot{V}E/\dot{V}CO_2$ slope as a function of LAVi. **(F)** $\dot{V}E/\dot{V}CO_2$ slope as a function of TRV_{max}.