

L2: Basics of Linear Algebra

Topics to be covered today:

- Vectors
- Inner product
- Linear independence
- Basis, dimension of a subspace
- Matrices
- Rank of a matrix

Def: A vector is an ordered finite list of numbers.

$$\text{Eg: } a = \begin{pmatrix} -1.1 \\ 2.3 \\ 4 \\ 7 \end{pmatrix}$$

↳ 4-dimensional vector



We say $a \in \mathbb{R}^4$ or that a is a vector of length/dimension 4.

a_i - i^{th} entry/component of a .

$$\text{Eg: } a_3 = 4$$

It is convenient to ~~visu~~ visualize a vector as:



- a point in $\mathbb{R}^n$
- an arrow from the origin to this point in $\mathbb{R}^n$




Vector addition

Vectors of the same size can be added (elementwise).

The visual interpretation of vector addition is useful.




These visualizations are equivalent.



Scalar multiplication:

$$\underset{\substack{\downarrow \\ \text{scalar}}}{\alpha} \underset{\substack{\downarrow \\ \text{vector}}}{a} = \begin{pmatrix} \alpha a_1 \\ \vdots \\ \alpha a_n \end{pmatrix}$$



Scalar multiplication also has a useful interpretation

Inner / Dot product

Say $a, b \in \mathbb{R}^n$

$$\text{Then } \langle a, b \rangle = \sum_{i=1}^n a_i b_i$$

This file is meant for personal use by me (Pravish Patil) for educational purposes only. Sharing or publishing the content in part or full is liable for legal action.

Also convenient to view inner product as a matrix multiplication:

View a & b as $n \times 1$ column matrices.

Then $\langle a, b \rangle = a^T b$

$$a = \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} \quad b = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}$$

$$a^T b = (a_1, \dots, a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \dots + a_n b_n$$

The norm of a vector a is defined as:


$$\|a\| = \sqrt{\langle a, a \rangle}$$

↓
magnitude of vector a

★ Important property of norms


Norm satisfies triangle inequality:

$$\|a+b\| \leq \|a\| + \|b\|$$



The distance between vectors a & b is given by:

$$\text{dist}(a, b) = \|a - b\|$$



★ Another important property of inner product is norm

Cauchy - Schwartz inequality :

$$|\langle a, b \rangle| \leq \|a\| \cdot \|b\|$$



$$-\|a\| \cdot \|b\| \leq \langle a, b \rangle \leq \|a\| \cdot \|b\|$$

- You get equality on the right,
i.e. $\langle a, b \rangle = \|a\| \cdot \|b\|$

iff $a = \alpha b$, where $\alpha > 0$.

(Vectors are pointed in the same "direction")


- You get equality on the left,
i.e. $\langle a, b \rangle = -\|a\| \cdot \|b\|$

iff $a = -\alpha b$, where $\alpha > 0$.

(Vectors are pointed in opposite "directions")

We say a is orthogonal / perpendicular to b if $\langle a, b \rangle = 0$.

Projection



The projection of b onto a is $\frac{\langle a, b \rangle}{\|a\|^2} a$


Magnitude of this projection is : $\frac{|\langle a, b \rangle|}{\|a\|}$.

Def A collection of vectors $\{a_1, \dots, a_k\}$ is linearly dependent if

$$\beta_1 a_1 + \beta_2 a_2 + \dots + \beta_k a_k = 0$$

holds for some $\beta_1, \dots, \beta_k$ that are NOT all zero.

{ This means at least one vector in the collection can be expressed as a linear combination of the others! }



Eg: The vectors $\{a, b, c\}$ ^{in $\mathbb{R}^2$} are ~~not~~ linearly dependent - We can write

$$c = \beta_1 a + \beta_2 b$$

$$\Rightarrow \beta_1 a + \beta_2 b + (-1)c = 0.$$

Def A collection $\{a_1, \dots, a_n\}$ is linearly independent if it is NOT linearly dependent.

{ This means no vector in this collection can be expressed as a linear combination of the others. }

Eg The collection $\{a, b\}$ in the above example is linearly independent.

Eg The collection $\{e_1, \dots, e_n\}$ of vectors in $\mathbb{R}^n$, where

unit $\leftarrow$ vectors $e_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \text{ in } i\text{th position} \\ \vdots \\ 0 \end{pmatrix}$

is linearly independent. - Exercise

Fact: If $\{a_1, \dots, a_k\}$ is an independent collection of vectors in $\mathbb{R}^n$, then $k \leq n$.

{ You cannot have a collection of 4 independent vectors in $\mathbb{R}^3$. }

Basis & dimension

Def A collection of n independent vectors in $\mathbb{R}^n$ forms a basis of $\mathbb{R}^n$.

Fact: If $\{a_1, \dots, a_n\}$ is a basis of $\mathbb{R}^n$, any vector $b \in \mathbb{R}^n$ can be expressed as a unique linear combination of the basis vectors,

$$\text{i.e., } b = \sum_{i=1}^n \beta_i a_i.$$

Eg $\{e_1, \dots, e_n\}$ just defined is a basis of $\mathbb{R}^n$.

In fact, this is an orthogonal basis.

{ any pair of basis vectors }
is orthogonal.

In fact, this is an orthonormal basis

{ orthogonal basis with
Unit norm basis vectors }

More generally, we can also ~~design~~ define basis & dimension of subspaces of $\mathbb{R}^n$.

Consider $\{a_1, \dots, a_k\}$.

Let L denote the set of all vectors that may be expressed as linear combinations of this collection.
→ subspace spanned by $\{a_1, \dots, a_k\}$

- If $\{a_1, \dots, a_k\}$ is linearly independent, it forms a basis of the subspace L .

We say L is k -dimensional.

- If not, there is a subset $\{a_{i_1}, \dots, a_{i_m}\}$ of the collection s.t. every vector in L is a unique linear combination of the vectors in this subset.

We say L is m -dimensional in this case.

Eg $\left\{ \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}, \begin{pmatrix} -3 \\ 4 \\ -1 \end{pmatrix} \right\}$
Span a 2-dimensional subspace of $\mathbb{R}^3$.

Exercise: Check that any two vectors of this collection are linearly independent, but

$$\begin{pmatrix} -3 \\ 4 \\ 1 \end{pmatrix} = -\frac{1}{5} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} - \frac{7}{5} \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}.$$

Matrices

A matrix is a rectangular array of numbers.

Eg

$$A = \begin{bmatrix} 0 & 1 & -2.3 & 0.1 \\ 1.3 & 4 & -0.1 & 0 \\ 4.1 & -1 & 0 & 1.7 \end{bmatrix}$$

We say A is $m \times n$ if it has m rows & n columns.

Eg: Above matrix is 3×4 .

$A_{i,j}$

We use $A_{i,j}$ to denote the entry in i^{th} row & j^{th} col.

$A_{3,2}$

Eg $A_{3,2} = -1.$

$$\left\{ \begin{array}{l} m = n \Rightarrow \text{square matrix} \\ m > n \Rightarrow \text{tall matrix} \\ m < n \Rightarrow \text{wide matrix} \end{array} \right\}$$

For $m \times n$ matrix,

- There are m row vectors, each in $\mathbb{R}^n$

Subspace spanned by rows - row space

Dimension of row space - row rank

Note: $\text{Row rank}(A) \leq \min(m, n)$
of independent rows

- There are n column vectors, each in $\mathbb{R}^m$

Subspace spanned by columns - column space

Dimension of column space - column rank

Note: $\text{Col. rank of } A \leq \min(m, n)$
of independent columns

Facts

- $\text{Col. rank}(A) = \text{Row rank}(A) = \text{rank}(A)$
- ~~For~~ For square $n \times n$ matrix A ,

$\text{rank}(A) = n \iff A$ is invertible
 A is "full rank"

$\exists n \times n$ matrix A^{-1} s.t.
 $AA^{-1} = A^{-1}A = I$
identity matrix

Q: Can a tall matrix have full row rank?

Q: Can a tall matrix have full col. rank?