

# Propositional Logic

Mădălina Eraşcu

West University of Timișoara and Institute e-Austria Timișoara  
bvd. V. Parvan 4, Timișoara, Romania

`madalina.erascu@e-uvt.ro`



*Based on the book: Chin-Liang Chang and Richard Char-Tung Lee. Symbolic Logic and Mechanical Theorem Proving, Chapter 2*

# Outline

Syntax

Semantics

Normal Forms in the Propositional Logic

Logical Consequence

# Outline

Syntax

Semantics

Normal Forms in the Propositional Logic

Logical Consequence

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\iff$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \iff H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Leftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Leftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\iff$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \iff H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\iff$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \iff H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?



# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Syntax

## Definition

A *proposition* is a declarative sentence that is either true ( $\mathbb{T}$ ) or false ( $\mathbb{F}$ ), but not both.

Can you give some examples?

We use symbols like  $P$ ,  $Q$ ,  $R$ , etc. for denoting propositions. They are called **atomic formulas** or **atoms**.

Complex propositions are built using logical connectives:  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\Rightarrow$  (implication),  $\Longleftrightarrow$  (equivalence).

## Definition (Syntax)

*Well-formed formulas (formulas)* in propositional logic are defined recursively as follows:

1. An *atom* is a formula.
2. If  $G$  is a formula, then  $\neg G$  is a formula.
3. If  $G$  and  $H$  are formulas, then  $G \wedge H$ ,  $G \vee H$ ,  $G \Rightarrow H$ , and  $G \Longleftrightarrow H$  are formulas.
4. All formulas are generated by applying the rules above.

What is the meaning of 4. ?

Can you give some examples/counterexamples of formulas?

# Outline

Syntax

**Semantics**

Normal Forms in the Propositional Logic

Logical Consequence

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$\mathcal{B}_{\neg}$		$\mathcal{B}_{\wedge}$	T	F	$\mathcal{B}_{\vee}$	T	F	$\mathcal{B}_{\Rightarrow}$	T	F	$\mathcal{B}_{\iff}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = \mathcal{B}_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?



# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$\mathcal{B}_{\neg}$		$\mathcal{B}_{\wedge}$	T	F	$\mathcal{B}_{\vee}$	T	F	$\mathcal{B}_{\Rightarrow}$	T	F	$\mathcal{B}_{\Leftrightarrow}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = \mathcal{B}_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$B_{\neg}$		$B_{\wedge}$	T	F	$B_{\vee}$	T	F	$B_{\Rightarrow}$	T	F	$B_{\Leftrightarrow}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = B_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$B_{\neg}$		$B_{\wedge}$	T	F	$B_{\vee}$	T	F	$B_{\Rightarrow}$	T	F	$B_{\Leftrightarrow}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = B_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$B_{\neg}$		$B_{\wedge}$	T	F	$B_{\vee}$	T	F	$B_{\Rightarrow}$	T	F	$B_{\Leftrightarrow}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = B_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$\mathcal{B}_{\neg}$		$\mathcal{B}_{\wedge}$	T	F	$\mathcal{B}_{\vee}$	T	F	$\mathcal{B}_{\Rightarrow}$	T	F	$\mathcal{B}_{\iff}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = \mathcal{B}_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$\mathcal{B}_{\neg}$		$\mathcal{B}_{\wedge}$	T	F	$\mathcal{B}_{\vee}$	T	F	$\mathcal{B}_{\Rightarrow}$	T	F	$\mathcal{B}_{\iff}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = \mathcal{B}_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Semantics

## Definition (Semantics)

The semantics of a formula  $G$ , is a function  $f_G : \mathcal{I} \rightarrow \{\mathbb{T}, \mathbb{F}\}$  with  $\mathcal{I} = \{I \mid \text{Vars}(G) \rightarrow \{\mathbb{T}, \mathbb{F}\}\}$ .

We introduce the notation  $\langle G \rangle_I$  instead of  $f_G(I)$  meaning *the truth evaluation of the formula  $G$  in the interpretation  $I$* .

## Definition (Interpretation)

Given a propositional formula  $G$ , let  $A_1, \dots, A_n$  be the atoms occurring in the formula  $G$ . Then an **interpretation** of  $G$  is an assignment of truth values to  $A_1, \dots, A_n$  in which every  $A_i$  is assigned either  $\mathbb{T}$  or  $\mathbb{F}$ , but not both.

## Example

Evaluate the truth value of  $(A \wedge (A \Rightarrow B)) \Rightarrow B$ .

To evaluate it we need to know an interpretation  $I$  as well as the semantics of the logical connectives.

$\mathcal{B}_{\neg}$		$\mathcal{B}_{\wedge}$	T	F	$\mathcal{B}_{\vee}$	T	F	$\mathcal{B}_{\Rightarrow}$	T	F	$\mathcal{B}_{\Longleftrightarrow}$	T	F
T	F	T	T	F	T	T	T	T	T	F	T	T	F
F	T	F	F	F	F	T	F	F	T	T	F	F	T

Then we have

$$\langle (A \wedge (A \Rightarrow B)) \Rightarrow B \rangle_I = \mathcal{B}_{\Rightarrow}(\langle A \wedge (A \Rightarrow B) \rangle_I, \langle B \rangle_I) = \dots$$

For a formula with  $n$  distinct atoms, how many distinct interpretation do exist for it?

# Validity/Invalidity, Inconsistency (unsatisfiability) / Consistency (satisfiability)

## Definition (Validity/Invalidity)

A formula  $F$  is said to be **valid** iff it is true under all its interpretations (For any  $I \in \mathcal{I} : \langle F \rangle_I = \mathbb{T}$ ). A formula is said to be **invalid** iff it is not valid.

## Definition (Inconsistent (unsatisfiable) / Consistent (satisfiable))

A formula is said to be **inconsistent (unsatisfiable)** iff it is false under all its interpretations (For any  $I \in \mathcal{I} : \langle F \rangle_I = \mathbb{F}$ ). A formula is said to be **consistent (satisfiable)** iff it is not inconsistent.



# Validity/Invalidity, Inconsistency (unsatisfiability) / Consistency (satisfiability)

## Definition (Validity/Invalidity)

A formula  $F$  is said to be **valid** iff it is true under all its interpretations (For any  $I \in \mathcal{I} : \langle F \rangle_I = \mathbb{T}$ ). A formula is said to be **invalid** iff it is not valid.

## Definition (Inconsistent (unsatisfiable) / Consistent (satisfiable))

A formula is said to be **inconsistent** (**unsatisfiable**) iff it is false under all its interpretations (For any  $I \in \mathcal{I} : \langle F \rangle_I = \mathbb{F}$ ). A formula is said to be **consistent** (**satisfiable**) iff it is not inconsistent.

# Outline

Syntax

Semantics

**Normal Forms in the Propositional Logic**

Logical Consequence

# Equivalent Formulas

## Definition (Equivalent Formulas)

Let  $F, G$  be two formulas. Then  $F = G$  iff for any  $I \in \mathcal{I} : \langle F \rangle_I = \langle G \rangle_I$ .

Proving the equivalence of two formulas:

1. By examining the truth tables of them
2. By rewriting
3. By bringing the two formulas in the normal form

# Equivalent Formulas

## Definition (Equivalent Formulas)

Let  $F, G$  be two formulas. Then  $F = G$  iff for any  $I \in \mathcal{I} : \langle F \rangle_I = \langle G \rangle_I$ .

Proving the equivalence of two formulas:

1. By examining the truth tables of them
2. By rewriting
3. By bringing the two formulas in the normal form

# Equivalent Formulas

## Definition (Equivalent Formulas)

Let  $F, G$  be two formulas. Then  $F = G$  iff for any  $I \in \mathcal{I} : \langle F \rangle_I = \langle G \rangle_I$ .

Proving the equivalence of two formulas:

1. By examining the truth tables of them
2. By rewriting
3. By bringing the two formulas in the normal form

# Equivalent Formulas

## Definition (Equivalent Formulas)

Let  $F, G$  be two formulas. Then  $F = G$  iff for any  $I \in \mathcal{I} : \langle F \rangle_I = \langle G \rangle_I$ .

Proving the equivalence of two formulas:

1. By examining the truth tables of them
2. By rewriting
3. By bringing the two formulas in the normal form

# Equivalent Formulas

## Definition (Equivalent Formulas)

Let  $F, G$  be two formulas. Then  $F = G$  iff for any  $I \in \mathcal{I} : \langle F \rangle_I = \langle G \rangle_I$ .

Proving the equivalence of two formulas:

1. By examining the truth tables of them
2. By rewriting
3. By bringing the two formulas in the normal form

# Equivalent transformations

Let  $\square$  be the formula which is always false,  $\blacksquare$  the formula which is always true (tautology).

We have the followings:

$$F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$F \Rightarrow G = \neg F \vee G$$

$$F \vee G = G \vee F$$

$$F \vee (G \vee H) = (F \vee G) \vee H$$

$$F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$F \vee \square = F$$

$$F \vee \blacksquare = \blacksquare$$

$$F \vee \neg F = \blacksquare$$

$$\neg(\neg F) = F$$

$$\neg(F \vee G) = \neg F \wedge \neg G$$

$$F \wedge G = G \wedge F$$

$$F \wedge (G \wedge H) = (F \wedge G) \wedge H$$

$$F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

$$F \wedge \blacksquare = F$$

$$F \wedge \square = \square$$

$$F \wedge \neg F = \square$$

$$\neg(F \wedge G) = \neg F \vee \neg G$$

(comutativity)

(associativity)

(distributivity)

(de Morgan)

## Definition (Literal)

A literal is an atom or the negation of an atom.



# Equivalent transformations

Let  $\square$  be the formula which is always false,  $\blacksquare$  the formula which is always true (tautology).

We have the followings:

$$F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$F \Rightarrow G = \neg F \vee G$$

$$F \vee G = G \vee F$$

$$F \vee (G \vee H) = (F \vee G) \vee H$$

$$F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$F \vee \square = F$$

$$F \vee \blacksquare = \blacksquare$$

$$F \vee \neg F = \blacksquare$$

$$\neg(\neg F) = F$$

$$\neg(F \vee G) = \neg F \wedge \neg G$$

$$F \wedge G = G \wedge F$$

$$F \wedge (G \wedge H) = (F \wedge G) \wedge H$$

$$F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

$$F \wedge \blacksquare = F$$

$$F \wedge \square = \square$$

$$F \wedge \neg F = \square$$

$$\neg(F \wedge G) = \neg F \vee \neg G$$

(comutativity)

(associativity)

(distributivity)

(de Morgan)

## Definition (Literal)

A literal is an atom or the negation of an atom.

# Equivalent transformations

Let  $\square$  be the formula which is always false,  $\blacksquare$  the formula which is always true (tautology).

We have the followings:

$$F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$F \Rightarrow G = \neg F \vee G$$

$$F \vee G = G \vee F$$

$$F \vee (G \vee H) = (F \vee G) \vee H$$

$$F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$F \vee \square = F$$

$$F \vee \blacksquare = \blacksquare$$

$$F \vee \neg F = \blacksquare$$

$$\neg(\neg F) = F$$

$$\neg(F \vee G) = \neg F \wedge \neg G$$

$$F \wedge G = G \wedge F$$

$$F \wedge (G \wedge H) = (F \wedge G) \wedge H$$

$$F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

$$F \wedge \blacksquare = F$$

$$F \wedge \square = \square$$

$$F \wedge \neg F = \square$$

$$\neg(F \wedge G) = \neg F \vee \neg G$$

(comutativity)

(associativity)

(distributivity)

(de Morgan)

## Definition (Literal)

A **literal** is an atom or the negation of an atom.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

► **Step 1.** Use the laws

$$\text{► } F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$\text{► } F \Rightarrow G = \neg F \vee G$$

to eliminate  $\iff$  and  $\Rightarrow$ .

► **Step 2.** Repeatedly use the laws

$$\text{► } \neg(\neg F) = F$$

and de Morgan's laws

$$\text{► } \neg(F \vee G) = \neg F \wedge \neg G$$

$$\text{► } \neg(F \wedge G) = \neg F \vee \neg G$$

to bring the negation signs immediately before atoms.

► **Step 3.** Repeatedly use the distributive laws

$$\text{► } F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$\text{► } F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

and the other laws to obtain a normal form.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

► **Step 1.** Use the laws

$$\text{► } F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$\text{► } F \Rightarrow G = \neg F \vee G$$

to eliminate  $\iff$  and  $\Rightarrow$ .

► **Step 2.** Repeatedly use the laws

$$\text{► } \neg(\neg F) = F$$

and de Morgan's laws

$$\text{► } \neg(F \vee G) = \neg F \wedge \neg G$$

$$\text{► } \neg(F \wedge G) = \neg F \vee \neg G$$

to bring the negation signs immediately before atoms.

► **Step 3.** Repeatedly use the distributive laws

$$\text{► } F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$\text{► } F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

and the other laws to obtain a normal form.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

► **Step 1.** Use the laws

$$\text{► } F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$\text{► } F \Rightarrow G = \neg F \vee G$$

to eliminate  $\iff$  and  $\Rightarrow$ .

► **Step 2.** Repeatedly use the laws

$$\text{► } \neg(\neg F) = F$$

and de Morgan's laws

$$\text{► } \neg(F \vee G) = \neg F \wedge \neg G$$

$$\text{► } \neg(F \wedge G) = \neg F \vee \neg G$$

to bring the negation signs immediately before atoms.

► **Step 3.** Repeatedly use the distributive laws

$$\text{► } F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$\text{► } F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

and the other laws to obtain a normal form.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

► **Step 1.** Use the laws

$$\text{► } F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$$

$$\text{► } F \Rightarrow G = \neg F \vee G$$

to eliminate  $\iff$  and  $\Rightarrow$ .

► **Step 2.** Repeatedly use the laws

$$\text{► } \neg(\neg F) = F$$

and de Morgan's laws

$$\text{► } \neg(F \vee G) = \neg F \wedge \neg G$$

$$\text{► } \neg(F \wedge G) = \neg F \vee \neg G$$

to bring the negation signs immediately before atoms.

► **Step 3.** Repeatedly use the distributive laws

$$\text{► } F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$$

$$\text{► } F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$$

and the other laws to obtain a normal form.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

- ▶ **Step 1.** Use the laws

- ▶  $F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$

- ▶  $F \Rightarrow G = \neg F \vee G$

to eliminate  $\iff$  and  $\Rightarrow$ .

- ▶ **Step 2.** Repeatedly use the laws

- ▶  $\neg(\neg F) = F$

and de Morgan's laws

- ▶  $\neg(F \vee G) = \neg F \wedge \neg G$

- ▶  $\neg(F \wedge G) = \neg F \vee \neg G$

to bring the negation signs immediately before atoms.

- ▶ **Step 3.** Repeatedly use the distributive laws

- ▶  $F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$

- ▶  $F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$

and the other laws to obtain a normal form.

# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

- ▶ **Step 1.** Use the laws

- ▶  $F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$

- ▶  $F \Rightarrow G = \neg F \vee G$

to eliminate  $\iff$  and  $\Rightarrow$ .

- ▶ **Step 2.** Repeatedly use the laws

- ▶  $\neg(\neg F) = F$

and de Morgan's laws

- ▶  $\neg(F \vee G) = \neg F \wedge \neg G$

- ▶  $\neg(F \wedge G) = \neg F \vee \neg G$

to bring the negation signs immediately before atoms.

- ▶ **Step 3.** Repeatedly use the distributive laws

- ▶  $F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$

- ▶  $F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$

and the other laws to obtain a normal form.



# Normal Forms

## Definition (Negation Normal Form)

A formula  $F$  is in **negation normal form (NNF)** iff  $F$  contains only the connectives  $\neg$ ,  $\wedge$ , and  $\vee$  and that negations appear only in literals.

## Definition (Conjunctive Normal Form)

A formula  $F$  is in **conjunctive normal form (CNF)** iff  $F$  is in the form  $F_1 \wedge \dots \wedge F_n$ ,  $n \geq 1$ , where each  $F_i$  is a disjunction of literals.

## Definition (Disjunctive Normal Form)

A formula  $F$  is in **disjunctive normal form (DNF)** iff  $F$  is in the form  $F_1 \vee \dots \vee F_n$ ,  $n \geq 1$ , where each  $F_i$  is a conjunction of literals.

A formula can be brought into a normal form by following the next steps:

- ▶ **Step 1.** Use the laws

- ▶  $F \iff G = (F \Rightarrow G) \wedge (G \Rightarrow F)$

- ▶  $F \Rightarrow G = \neg F \vee G$

to eliminate  $\iff$  and  $\Rightarrow$ .

- ▶ **Step 2.** Repeatedly use the laws

- ▶  $\neg(\neg F) = F$

and de Morgan's laws

- ▶  $\neg(F \vee G) = \neg F \wedge \neg G$

- ▶  $\neg(F \wedge G) = \neg F \vee \neg G$

to bring the negation signs immediately before atoms.

- ▶ **Step 3.** Repeatedly use the distributive laws

- ▶  $F \vee (G \wedge H) = (F \vee G) \wedge (F \vee H)$

- ▶  $F \wedge (G \vee H) = (F \wedge G) \vee (F \wedge H)$

and the other laws to obtain a normal form.

# Outline

Syntax

Semantics

Normal Forms in the Propositional Logic

**Logical Consequence**

# Logical Consequences

## Definition

Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff for all interpretation  $I$  in which  $F_1 \wedge F_2 \wedge \dots \wedge F_n$  is true,  $G$  is also true.  $F_1, F_2, \dots, F_n$  are called **axioms/postulates/premises**.

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is valid.*

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $F_1 \wedge F_2 \wedge \dots \wedge F_n \wedge \neg G$  is inconsistent.*

If  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$ , the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is called a **theorem**, and  $G$  is also called the **conclusion of the theorem**.

# Logical Consequences

## Definition

Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff for all interpretation  $I$  in which  $F_1 \wedge F_2 \wedge \dots \wedge F_n$  is true,  $G$  is also true.  $F_1, F_2, \dots, F_n$  are called **axioms/postulates/premises**.

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is valid.*

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $F_1 \wedge F_2 \wedge \dots \wedge F_n \wedge \neg G$  is inconsistent.*

If  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$ , the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is called a **theorem**, and  $G$  is also called the **conclusion of the theorem**.

# Logical Consequences

## Definition

Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff for all interpretation  $I$  in which  $F_1 \wedge F_2 \wedge \dots \wedge F_n$  is true,  $G$  is also true.  $F_1, F_2, \dots, F_n$  are called **axioms/postulates/premises**.

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is valid.*

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $F_1 \wedge F_2 \wedge \dots \wedge F_n \wedge \neg G$  is inconsistent.*

If  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$ , the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is called a **theorem**, and  $G$  is also called the **conclusion of the theorem**.

# Logical Consequences

## Definition

Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff for all interpretation  $I$  in which  $F_1 \wedge F_2 \wedge \dots \wedge F_n$  is true,  $G$  is also true.  $F_1, F_2, \dots, F_n$  are called **axioms/postulates/premises**.

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is valid.*

## Theorem

*Given formulas  $F_1, F_2, \dots, F_n$  and a formula  $G$ ,  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$  iff the formula  $F_1 \wedge F_2 \wedge \dots \wedge F_n \wedge \neg G$  is inconsistent.*

If  $G$  is a logical consequence of  $F_1, F_2, \dots, F_n$ , the formula  $(F_1 \wedge F_2 \wedge \dots \wedge F_n) \Rightarrow G$  is called a **theorem**, and  $G$  is also called the **conclusion of the theorem**.