

Yr 3: Chemical Engineering Thermodynamics - Exam Crib Sheet

Genevieve Clifford

Semester 3, Week 4

1 1st Law of Thermodynamics

The total energy of an isolated system is constant; energy can be transformed from one form to another, but can neither be created nor destroyed.

Internal energy change = heat in - work done by the system on surroundings

$$\begin{aligned}dU &= dq - dw \\dU &= Tds - pdV \\H &= U + pV \\dH &= dU + Vdp + pdV \\dH &= Tds + Vdp\end{aligned}$$

Some partial differentiation yields:

$$\left(\frac{\partial^2 S}{\partial P \partial T}\right) = \frac{1}{T} \left(\frac{\partial H}{\partial P \partial T}\right) \quad (2.8)$$

$$T \left(\frac{\partial^2 S}{\partial T \partial P}\right) + \left(\frac{\partial S}{\partial P}\right)_T = \left(\frac{\partial^2 H}{\partial T \partial P}\right) - \left(\frac{\partial V}{\partial T}\right) \quad (2.9)$$

By making use of Schwarz's Theorem, we yield:

$$\left(\frac{\partial^2 S}{\partial P \partial T}\right) \equiv \left(\frac{\partial^2 S}{\partial T \partial P}\right) \quad (2.10)$$

$$\left(\frac{\partial^2 H}{\partial P \partial T}\right) \equiv \left(\frac{\partial^2 H}{\partial T \partial P}\right) \quad (2.11)$$

$$\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial V}{\partial T}\right)_P \quad (2.12)$$

$$(2.13)$$

2 2nd Law of Thermodynamics

The total entropy of an isolated system can never decrease over time, and is constant if and only if all processes are reversible. In general, the total entropy of an isolated system tends to increase.

$$\Delta s = \int \frac{dq}{T} \quad (2.1)$$

It is now useful to derive the specific heats:

$$C_p = \left(\frac{\partial H}{\partial T}\right)_P \quad (2.14)$$

$$C_v = \left(\frac{\partial U}{\partial T}\right)_V \quad (2.15)$$

2.1 Analytic solution to 2nd Law

$$dH = Tds + Vdp \quad (2.2)$$

We can include the generic statement that dH and dS are equal to the sum of their partial derivatives:

$$dH = \left(\frac{\partial H}{\partial T}\right)_P dT + \left(\frac{\partial H}{\partial P}\right)_T dP \quad (2.3)$$

$$dS = \left(\frac{\partial S}{\partial T}\right)_P dT + \left(\frac{\partial S}{\partial P}\right)_T dP \quad (2.4)$$

By setting Equation 2.2 equal to Equation 2.3, it is possible to extract:

$$dS = \frac{1}{T} \left(\frac{\partial H}{\partial T}\right)_P dT + \frac{1}{T} \left[\left(\frac{\partial H}{\partial P}\right)_T - \frac{V}{T} \right] dP \quad (2.5)$$

By directly comparing Equations 2.4 and 2.5, it is possible to derive the following equations:

$$\left(\frac{\partial S}{\partial T}\right)_P = \frac{1}{T} \left(\frac{\partial H}{\partial T}\right)_P \quad (2.6)$$

$$T \left(\frac{\partial S}{\partial P}\right)_T = \left(\frac{\partial H}{\partial P}\right)_T - V \quad (2.7)$$

Through careful combination of previous derived equations it is now possible to derive:

$$dS = \frac{C_p dT}{T} - \left(\frac{\partial V}{\partial T}\right)_P dP \quad (2.16)$$

Assuming C_p is constant and that the ideal gas law applies, we gain the following useful formula:

$$\Delta S = nC_p \ln \left(\frac{T_2}{T_1}\right) - nR \ln \left(\frac{P_2}{P_1}\right) \quad (2.17)$$

2.2 Derivation of a formula for enthalpy change in terms of Equation of State

By use of Equation 2.7 and 2.12, it is possible to derive an equation that delivers the enthalpy change, fit for use with Equations of State (i.e. Ideal Gas Law, van der Waals, Peng-Robinson, Soave-Redlich-Kwong etc.)

$$\left(\frac{\partial H}{\partial P}\right)_T = T \left(\frac{\partial S}{\partial P}\right)_T + V \quad (2.18)$$

$$\left(\frac{\partial H}{\partial P}\right)_T = V - T \left(\frac{\partial V}{\partial T}\right)_P \quad (2.19)$$

Combining Equation 2.3 with 2.19 and 2.14, it is possible to derive the full general equation:

$$dH = C_p dT + \left[V - T \left(\frac{\partial V}{\partial T} \right)_P \right] dP \quad (2.20)$$

If we make the assumption that the Ideal Gas Law applies, i.e. $Pv = RT$, the equation becomes:

$$dH = C_p dT \quad (2.21)$$

2.3 Derivation for equations linking the heat capacities

It is necessary to define dU , then derive an equation to gain dS in terms of dU :

$$dU = TdS - PdV \quad (2.22)$$

$$dU = \left(\frac{\partial U}{\partial T} \right)_V dT + \left(\frac{\partial U}{\partial V} \right)_T dV \quad (2.23)$$

$$dS = \frac{1}{T} \left(\frac{\partial U}{\partial T} \right)_V dT + \frac{1}{T} \left[\left(\frac{\partial U}{\partial V} \right)_T + \frac{P}{T} \right] dV \quad (2.24)$$

Setting Equation 2.4 equal to 2.24, the following is yielded:

$$T \left(\frac{\partial S}{\partial T} \right)_V = \left(\frac{\partial U}{\partial T} \right)_V = C_V \quad (2.25)$$

Along with the relation gained in Equation 2.6, the following equation can be drawn:

$$C_p - C_V = T \left[\left(\frac{\partial S}{\partial T} \right)_P - \left(\frac{\partial S}{\partial T} \right)_V \right] \quad (2.26)$$

Dividing Equation 2.4 by ∂T yields:

$$\left(\frac{\partial S}{\partial T} \right)_V = \left(\frac{\partial S}{\partial T} \right)_P + \left(\frac{\partial S}{\partial P} \right)_T \left(\frac{\partial P}{\partial T} \right)_V \quad (2.27)$$

This is merged with previously derived quantities to give:

$$C_p - C_V = T \left(\frac{\partial V}{\partial T} \right)_P \left(\frac{\partial P}{\partial T} \right)_V \quad (2.28)$$

When used in conjunction with the Ideal Gas Law, the equation simplifies to:

$$C_p - C_V = R \quad (2.29)$$

3 3rd Law of Thermodynamics

The entropy of a system approaches a constant value as its temperature approaches absolute zero (0 K). Commonly given as "The entropy of a perfect crystal is zero when the temperature of the crystal is equal to absolute zero (0 K)".

4 Hess' Law

The enthalpy change accompanying a chemical change is independent of the route by which the chemical change occurs.

$$\Delta H^\circ = \sum_i \Delta H_{fi, \text{products}}^\circ - \sum_i \Delta H_{fi, \text{reactants}}^\circ \quad (4.1)$$

$$\Delta H_T^\circ = \Delta H_{298 \text{ K}}^\circ + \int_{298 \text{ K}}^T \Delta C_p dT \quad (4.2)$$

4.1 Formation Equations

$$\Delta_r H^\ominus = \sum v H^\ominus(\text{products}) - \sum v H^\ominus(\text{reactants}) \quad (4.3)$$

$$\Delta_r S^\ominus = \sum v S^\ominus(\text{products}) - \sum v S^\ominus(\text{reactants}) \quad (4.4)$$

$$\Delta_r G^\ominus = \sum v \Delta_f^\ominus G(\text{products}) - \sum v \Delta_f^\ominus G(\text{reactants}) \quad (4.5)$$

5 Free Energy and Equilibrium

Helmholtz Free Energy, with $dA = 0$ for equilibrium at constant temperature and volume:

$$A = U - TS \quad (5.1)$$

Gibbs Free Energy, with $dG = 0$ for equilibrium at a specified constant temperature and pressure:

$$G = H - TS \quad (5.2)$$

5.1 Derivation of the Clausius-Clapeyron equation

Assume there is a substance which is formed of two phases, α and β , these phases are in equilibrium. The two phases are at a temperature T and the system has a pressure, P . The value of dG for both phases is 0.

$$dG^\alpha = dG^\beta \quad (5.3)$$

$$dG^\beta = v^\beta dP - S^\beta dT = dG^\alpha = v^\alpha dP - S^\alpha dT \quad (5.4)$$

$$\frac{dP}{dT} = \frac{S^\beta - S^\alpha}{V^\beta - V^\alpha} \quad (5.5)$$

$$S^\beta - S^\alpha = \frac{dQ}{T} = \frac{L}{T} \quad (5.6)$$

$$\frac{dP}{dT} = \frac{L}{T \Delta V} \quad (5.7)$$

$$\Delta V = V^G - V^L \text{ and } V^G = RT/P \gg V^L \quad (\text{Ideal Gas})$$

$$\frac{dP}{dT} = \frac{LP}{RT^2} \quad (5.8)$$

$$\frac{d \ln P}{dT} = \frac{L}{RT^2} \quad (5.9)$$

6 Chemical Potential

6.1 Derivation of equation linking equilibrium constant and Gibbs Free Energy

Assume the following reaction takes place and all species are Ideal and Perfect Gases:



$$\mu_i = \mu_i^\circ + \int_{P_o}^{P_i} \bar{V}_{mi} dP \quad (6.1)$$

$$\bar{V}_{mi} = V_{mi} = \frac{R_m T}{P} \quad P_i = y_i P \quad (6.2)$$

$$\mu_i = G_{mi}^\circ + \int_{P_o}^{P_i} \left(\frac{R_m T}{P} \right) dP \quad (6.3)$$

$$\mu_i = \mu_i^\circ + RT \ln \left(\frac{P_i}{P_o} \right) \quad (6.4)$$

$$R_m T \ln K^* + \Delta G^\circ = \Delta G \quad (6.5)$$

Where:

$$K^* = \left[\frac{\left(\frac{P_C}{P_o} \right)^c \left(\frac{P_D}{P_o} \right)^d}{\left(\frac{P_A}{P_o} \right)^a \left(\frac{P_B}{P_o} \right)^b} \right] \quad (6.6)$$

$$\Delta G^\circ = cG_{mC}^\circ + dG_{mD}^\circ - aG_{mA}^\circ + bG_{mB}^\circ \quad (6.7)$$

$$\Delta G = c\mu_C + d\mu_D - a\mu_A + b\mu_B \quad (6.8)$$

By definition, $\Delta G = 0$ at equilibrium, so:

$$\ln K = -\frac{\Delta G_T^\circ}{R_m T} \quad (6.9)$$

6.2 Derivation of the Gibbs-Helmholtz Equation

Taking and rearranging the work from the previous section (Equations 5.2 and 6.9), the following formulae are to be used as a starting point:

$$\Delta G_T^\circ = \Delta H_T^\circ - T \Delta S_T^\circ \quad (6.10)$$

$$\Delta G^\circ = -RT \ln K \quad (6.11)$$

$$\frac{\Delta G_T^\circ}{T} = \frac{\Delta H_T^\circ}{T} - \Delta S_T^\circ \quad (6.12)$$

The equation is then implicitly differentiated with partial respect to T. The enthalpy term is differentiated using the product rule.

$$\frac{\partial \frac{\Delta G_T^\circ}{T}}{\partial T} = \frac{1}{T} \frac{\partial H_T^\circ}{\partial T} - \frac{H_T^\circ}{T^2} + \frac{\partial S_T^\circ}{\partial T} \quad (6.13)$$

$$\frac{\partial H}{\partial T} \equiv C_p \therefore \frac{1}{T} \frac{\partial H_T^\circ}{\partial T} = \frac{C_p}{T} \quad (6.14)$$

$$ds \equiv \frac{dq}{dT} = \frac{C_p dT}{T} \therefore \frac{\partial S_T^\circ}{\partial T} = \frac{C_p}{T} \quad (6.15)$$

$$\frac{\partial \frac{\Delta G_T^\circ}{T}}{\partial T} = \frac{C_p}{T} - \frac{\Delta H_T^\circ}{T^2} - \frac{C_p}{T} \quad (6.16)$$

$$\frac{\partial \frac{\Delta G_T^\circ}{T}}{\partial T} = -\frac{\Delta H_T^\circ}{T^2} \quad (6.17)$$

6.3 Derivation of the van 't Hoff Equation

Using Equation 6.17, it is possible to derive the van 't Hoff Equation:

$$\Delta G_T^\circ = -RT \ln K \quad (6.18)$$

$$\frac{\Delta G_T^\circ}{T} = -R \ln K \quad (6.19)$$

$$-\frac{\partial (R \ln K)}{\partial T} = -\frac{\Delta H_T^\circ}{T^2} \quad (6.20)$$

$$\frac{\partial \ln K}{\partial T} = \frac{\Delta H_T^\circ}{RT^2} \quad (6.21)$$

7 Useful Maths

7.1 Schwarz's Theorem

$$\frac{\partial}{\partial x_j} \left(\frac{\partial \phi}{\partial x_i} \right) \equiv \frac{\partial}{\partial x_i} \left(\frac{\partial \phi}{\partial x_j} \right) \quad (7.1)$$

7.2 Differentiation Rules

7.2.1 Product Rule

$$y = u(x) \cdot v(x) \quad (7.2)$$

$$\frac{dy}{dx} = v(x) \frac{du(x)}{dx} + u(x) \frac{dv(x)}{dx} \quad (7.3)$$

7.2.2 Quotient Rule

$$y = \frac{u(x)}{v(x)} \quad (7.4)$$

$$\frac{dy}{dx} = \frac{\left[v(x) \frac{du(x)}{dx} - u(x) \frac{dv(x)}{dx} \right]}{v(x)^2} \quad (7.5)$$

7.2.3 Chain Rule

$$\frac{dz}{dx} = \frac{dz}{dy} \cdot \frac{dy}{dx} \quad (7.6)$$

7.3 Integration Rules

7.3.1 Integrating by Parts

$$\int u(x) \frac{dv(x)}{dx} dx = u(x)v(x) - \int v(x) \frac{du(x)}{dx} \quad (7.7)$$

7.3.2 Integrating Factor

$$\frac{dy}{dx} + P(x)y = Q(x) \quad (7.8)$$

$$y = \frac{\int IQ(x)dx}{I} \quad (7.9)$$

$$I = e^{\int P(x)dx} \quad (7.10)$$