

Quantitative skills for genomic data analysis (2): statistics

BBMS 3009: Genome Science (First Semester, 2021)

Dr. Yuanhua Huang / 黃淵華

School of Biomedical Sciences &

Department of Statistics and Actuarial Science



香港大學

THE UNIVERSITY OF HONG KONG

Today's learning objectives (Statistics)

1. Main idea of statistical hypothesis testing
2. Regression based test
3. Types of errors and evaluation metrics
4. More testing methods: t-test, permutation test, etc

Resource: Chapter 6 in book *Modern Statistics for Modern Biology*

<https://www.huber.embl.de/msmb/Chap-Testing.html>

R script: Moodle / Lecture handouts / Dr. YH Huang → R-statistics.Rmd



1. Example of hypothesis testing

- Hypothetic example:
 - Does advertising in newspaper have impact on the sales of cars
- Decision to make
 - Yes or No
- Factors to consider when using collected data:
 - Uncertain measurement
 - Sample size is not big enough to represent the whole population



Decision making and hypothesis testing

- Common scientific questions from observed data
 - Difference between groups
 - Tendency, e.g., along with drug doses (effects)
- How to make the decision with considering
 - Uncertainty in observations, e.g., patient responses
 - Sample size is not big enough to represent the population, e.g., in clinical trial
- Statistical hypothesis testing
 - An approach for decision making **under uncertainty**
 - Estimate the **probability** to be wrong
 - Maximize expected utility (subjective value)



Main idea of hypothesis testing

A statistical hypothesis, sometimes called confirmatory data analysis, is a hypothesis that is **testable** in the light of observed data that is modeled via **a set of random variables**.

Main idea

- It is difficult to prove that a fact is “right”.
- But it is easy to prove that an opposite fact is “wrong”.
- Then you only have to find one counter example.



Main idea of hypothesis testing

- It is difficult to prove that a fact is “right”, but it is easy to prove that an opposite fact is “wrong”.

Research hypothesis (alternative hypothesis)

- H_1 : the newspaper advertising **has impact** on sales

$$H_1: y = \beta_0 + \beta_1 \times \text{Newspaper}; \beta_1 \neq 0$$

Null hypothesis (default hypothesis, you don't need to prove it, just assume it)

- H_0 : the newspaper advertising **has no impact** on sales

$$H_0: y = \beta_0 + \beta_1 \times \text{Newspaper}; \beta_1 = 0$$




Main idea of hypothesis testing

- With null and alternative hypotheses set up, we then **try to show** that, in light of our collected data, the **null hypothesis is false**.
- We do this by calculating the probability of obtaining **test results** at least as **extreme** as the results **actually observed**, under the **assumption that the null hypothesis is correct**
 - If this probability is very small, it suggests that the null hypothesis is false.
 - If this probability is large, it suggests that there is not enough evidence to reject the null hypothesis.
- This probability is called the **p value** of the test



2. Regression-based testing

- Data collection
 - 200 samples with both newspaper advertising costs and sales values



```
library(datarium)
head(marketing)

# plotting
ggplot(marketing,
  aes(x=newspaper, y=sales)) +
  geom_point() +
  geom_smooth(method=lm)
```

Dataset: <https://search.r-project.org/CRAN/refmans/datarium/html/marketing.html>




香港大學

THE UNIVERSITY OF HONG KONG

Regression-based testing

- Fitting a regression model with maximum likelihood
 - $y = \beta_0 + \beta_1 \times \text{Newspaper};$



Maximum likelihood estimate:
mean and standard error

Intercept: $\beta_0 = 14.82 \pm 0.746$

Newspaper: $\beta_1 = 0.0547 \pm 0.0166$

T-statistic for β_1 :


t value = $0.0547 / 0.0166 = 3.3$




香港大學

THE UNIVERSITY OF HONG KONG

Regression-based testing



Under the null, the distribution of t-statistic;
Degree of freedom = $n_{\text{sample}} - n_{\text{coefficient}} = 198$



P value = $\text{prob}(x > t \text{ value}) = 0.00115$



香港大學

THE UNIVERSITY OF HONG KONG


P value and level of significance

- **P value**: under the null hypothesis, the probability to see the test results as extreme as observed test result
 - If p value is very small, it suggests that the null hypothesis is false.
 - But how low does this probability has to be before we can conclude that the null hypothesis is false?
- Convention: choose a **level of significance** before the experiment that dictates how low the p value should be before we reject the null hypothesis.
- It is common to choose a significance level of 0.01 or 0.05.
- Here $p = 0.00115$, so we **reject** the Null hypothesis at the significance level of 0.01.



More on regression-based testing (1)

- Likelihood ratio test
 - Null model likelihood L_0 : $y = \beta_0$
 - Alternative model likelihood L_1 : $y = \beta_0 + \beta_1 \times \text{Newspaper}$
- Likelihood with maximum likelihood estimate
 - Null hypothesis: $L_0 = -650.15$
 - Alternative hypothesis: $L_1 = -644.8$
- Likelihood ratio statistic
 - Observed results: $\lambda = -2(L_0 - L_1) = 10.7$
 - Distribution under the Null: $\lambda \sim \chi^2 (df = 1)$
 - P value: $P(x > \lambda) = 0.00107$



More on regression-based testing (2)


➤ Condition on other covariate, e.g., advertising on Facebook

- $H_1: y = \beta_0 + \beta_1 \times \text{Newspaper} + \beta_2 \times \text{Facebook} ; \beta_1 \neq 0$
- $H_0: y = \beta_0 + \beta_1 \times \text{Newspaper} + \beta_2 \times \text{Facebook} ; \beta_1 = 0$

	youtube <dbl>	facebook <dbl>	newspaper <dbl>	sales <dbl>
1	276.12	45.36	83.04	26.52
2	53.40	47.16	54.12	12.48
3	20.64	55.08	83.16	11.16
4	181.80	49.56	70.20	22.20
5	216.96	12.96	70.08	15.48
6	10.44	58.68	90.00	8.64

➤ Fitting the model with collected data

- $\beta_0 = 11.02 \pm 0.753$
 - $\beta_1 = 0.0066 \pm 0.0149$; t value = $0.0066/0.0149 = 0.446$
 - $\beta_2 = 0.199 \pm 0.022$
-
- P value = 0.656; fail to reject the null hypothesis at significance level of 0.01.



3. Types of errors

- Now, let's think about genome and we are testing if a gene expression is different between treatment and control for a cancer patient
- We will have a hypothesis testing for each gene, so around 10,000 tests in total.
- What errors on our decision?
 - **False positive (type I error):** Genes are **genuine not different**, but we thought they are (reject the null hypothesis)
 - **False negative (type II error):** Genes are **genuine different**, but we missed it (we didn't reject the null hypothesis)



Evaluation metrics

➤ True positive rate (**Power**, **Sensitivity**, Hit rate, **Recall**): $TPR = \frac{TP}{TP+FN}$

➤ True Negative Rate (**Specificity**): $TNR = \frac{TN}{TN+FP}$

➤ **Precision** (Positive Predictive Value; 1- **false discovery rate**):


$$Precision = \frac{TP}{TP + FP} = 1 - FDR$$

		Predicted condition	
		Positive (PP)	Negative (PN)
Actual condition	Positive (P)	True positive (TP) , hit	False negative (FN) , type II error, miss, underestimation
	Negative (N)	False positive (FP) , type I error, false alarm, overestimation	True negative (TN) , correct rejection



4. More hypothesis testing methods

- Is the Sepal width different between versicolor and virginica?
- Difference between groups
 - T test
 - Permutation test
 - Wilcoxon rank-sum test / Mann–Whitney U test
- These won't be examined but can be very useful for your future studies or self-learning



Resources

- Chapter 6 in book *Modern Statistics for Modern Biology*
 - <https://www.huber.embl.de/msmb/Chap-Testing.html>
- R script: Moodle / Lecture handouts / Dr. YH Huang → R-statistics.Rmd

