

Associations between optical, physical and chemical properties of aerosols measured at ground-based networks

Meredith Franklin¹, Meytar Sorek-Hamer², Olga Kalashnikova³, Michael Garay³, and David Diner³

¹University of Southern California, Los Angeles, CA, United States

²NASA Ames Research Center, Moffett Field, CA, USA

³Jet Propulsion Laboratory, Pasadena, CA, USA

Correspondence: Meredith Franklin (meredith.franklin@usc.edu)

Abstract. The abstract goes here. It can also be on *multiple lines*.

Copyright statement. The author's copyright for this publication is transferred to institution/company.

1 Introduction

<<<<<<< HEAD A large body of literature has shown that satellite-derived aerosol optical depth (AOD), typically retrieved
5 at 550 nm wavelength, reliably correlates with mass-volume concentrations of fine mode particulate matter with aerodynamic
diameter less than 2.5 μm ($\text{PM}_{2.5}$). Studies that have used satellite observations to generate $\text{PM}_{2.5}$ have been instrumental for
air pollution - health effects research. The associations between AOD and different chemical components of $\text{PM}_{2.5}$ are lesser
known. A handful of studies using the Multiangle Imaging SpectroRadiometer (MISR), an instrument onboard the NASA
Terra satellite that provides observations of optical properties by particle type (size, shape, absorption), have provided evidence
10 that different optical properties relate to different physical and chemical properties of particulate matter. Results have been
somewhat inconsistent, showing differences depending on geographic area of analysis, optical components used, and statistical
tools applied.

A large body of literature has shown that satellite-derived aerosol optical depth (AOD), typically retrieved at 550 nm wave-
length, reliably correlates with mass-volume concentrations of fine mode particulate matter with aerodynamic diameter less
15 than 2.5 μm ($\text{PM}_{2.5}$). Studies that have used satellite observations to generate $\text{PM}_{2.5}$ have been instrumental for air pollution
- health effects research. The associations between AOD and different chemical components of $\text{PM}_{2.5}$ are lesser known. A
handful of studies using the Multiangle Imaging SpectroRadiometer (MISR), an instrument onboard the NASA Terra satellite
that provides observations of optical properties by particle type (size, shape, absorption), have provided evidence that differ-
ent optical properties relate to different physical and chemical properties of particulate matter. Results have been somewhat
20 inconsistent, showing differences depending on geographic area of analysis, optical components used, and statistical tools
applied.

The purpose of this analysis is to make a detailed examination of the statistical relationships between ground-level PM_{2.5} and PM_{2.5} chemical components (nitrate, sulfate, elemental carbon, organic carbon, dust) and optical measures of aerosols (e.g. aerosol optical depth, angstrom exponent).

AERONET ????

5 – Specific sections of the manuscript:

- running with title and author
- competing interests
- copyright statement (optional)
- availability (strongly recommended if any used), one of code, data, or codedata
- 10 – author contribution
- disclaimer
- acknowledgements

See the defaults and examples from the skeleton and the official Copernicus documentation for details.

Important: Always double-check with the official manuscript preparation guidelines at https://publications.copernicus.org/for_authors/manuscript_preparation.html, especially the sections “Technical instructions for LaTeX” and “Manuscript composition”. Please contact Daniel Nüst, daniel.nuest@uni-muenster.de, with any problems.

2 Methods

The study encompasses the San Joaquin Valley region of central California (Figure 1). We leverage datasets available at four sites - Bakersfield, Fresno, Modesto, and Visalia. At these sites there are co-located instruments from EPA’s chemical speciation
20 network (CSN), EPA’s air quality system (AQS) and NASA’s AERONET network.

2.1 Data

The CSN monitors are on a 1-in-3 or 1-in-6 day sampling schedule, providing PM_{2.5} mass and component PM_{2.5} concentrations of metals (e.g. Aluminium Al, Silicon Si, Calcium Ca, Titanium Ti, Iron Fe) obtained from X-ray fluorescence (XRF), ions (nitrate NO₃⁻ and sulfate SO₄²⁻) from ion chromatography, and carbons (organic OC and elemental EC) from thermal/optical
25 analysis. To quantify dust we use the following equation?: $\text{dust} = 2.2\text{Al} + 2.49\text{Si} + 1.63\text{Ca} + 1.94\text{Ti} + 2.42\text{Fe}$

The AQS monitors provide daily concentrations of PM_{2.5} mass by the EPA’s Federal Reference Method, which is the highest quality gravimetric measurement method used for regulatory purposes.

AERONET sites are sunphotometers providing a “ground-up” measurement of aerosol optical properties at multiple wavelengths and have been used extensively to validate “top-down” satellite observations of related properties. Wavelength-specific

AOD and angstrom exponents are the primary sunphotometer variables. Using quadratic log-log interpolation we calculated AOD 550 nm from AOD 440, 500, 675, 870 nm in log-log space. AOD at 550 nm is the most common wavelength retrieved from satellite instruments. A retrieval-based AERONET product, called the inversion product, provides an additional suite of aerosol properties that help distinguish size (fine, coarse effective radius), shape (asymmetry), and absorption. We excluded
5 sunphotometer and inversion variables that had a significant proportion of missing data (~90% missing). A list of the variables included in the analysis are shown in the Appendix. In a separate test we examine data from the SPARTAN site in Rehovot, Israel. The SPARTAN network provides data for PM_{2.5} mass and speciation concentrations on an integrated 1 in 9 day sampling schedule, and is colocated with an AERONET site (We don't have the speciation data for this site so we could only look at PM_{2.5} for now).

10 The CSN monitors are on a 1 in 3 or 1 in 6 day sampling schedule, providing PM_{2.5} mass and component PM_{2.5} concentrations of metals (e.g. Aluminium Al, Silicon Si, Calcium Ca, Titanium Ti, Iron Fe) obtained from X-ray fluorescence (XRF), ions (nitrate NO₃⁻ and sulfate SO₄²⁻) from ion chromatography, and carbons (organic OC and elemental EC) from thermal/optical analysis. To quantify dust we use the following equation: dust = 2.2Al + 2.49Si + 1.63Ca + 1.94Ti + 2.42Fe

The AQS monitors provide daily concentrations of PM_{2.5} mass by the EPA's Federal Reference Method, which is the highest
15 quality gravimetric measurement method used for regulatory purposes.

AERONET sites are sunphotometers providing a "ground-up" measurement of aerosol optical properties at multiple wavelengths and have been used extensively to validate "top-down" satellite observations of related properties. Wavelength-specific AOD and angstrom exponents are the primary sunphotometer variables. Using quadratic log-log interpolation we calculated AOD 550 nm from AOD 440, 500, 675, 870 nm in log-log space. AOD at 550 nm is the most common wavelength retrieved
20 from satellite instruments. A retrieval-based AERONET product, called the inversion product, provides an additional suite of aerosol properties that help distinguish size (fine, coarse effective radius), shape (asymmetry), and absorption. We excluded sunphotometer and inversion variables that had a significant proportion of missing data (~90% missing). A list of the variables included in the analysis are shown in the Appendix. In a separate test we examine data from the SPARTAN site in Rehovot, Israel. The SPARTAN network provides data for PM_{2.5} mass and speciation concentrations on an integrated 1 in 9 day sampling schedule, and is colocated with an AERONET site (We don't have the speciation data for this site so we could only look
25 at PM_{2.5} for now).

2.2 Statistical methods

Prior to model building we examined a cluster-based correlation heat map (Figure 2), which provides the Pearson correlations between all pairs of AERONET variables grouped by a decision tree. To avoid collinearity in the regression models, we kept the
30 most relevant of a group of variables that had a correlation coefficient > |0.9|. We then examined and picked a subset of variables connected at the mid-tier level of the tree to construct interactions. We fit simple linear regression models separately for PM_{2.5} mass, sulfate, nitrate, EC, OC, and dust with AOD 550 nm as the sole predictor variable. Multiple linear regression models were again fit separately for PM_{2.5} mass, sulfate, nitrate, EC, OC, and dust, but with the combined AERONET sunphotometer and inversion product as predictor variables and model selection was conducted using the "all possible subset method". This method

constructs models based on all combinations from 1 to k variable models. We select the best model from the combinations based on highest R², lowest RMSE, and Mallows' Cp statistic that is close to k+1. Model selection for the Fresno and Bakersfield sites were examined separately and in combination in a "total CA" analysis, which combined data from Fresno, Bakersfield, Modesto, Visalia, and a special DRAGON campaign in late 2012-early 2013 over the region (8 co-located sites with PM_{2.5} mass).

All models were cross validated (CV) with 10-fold CV, and we report the CV R² and RMSE. Models were fit in R using the leaps() library.

3 Results

3.1 Statistical methods

Prior to model building we examined a cluster-based correlation heat map (Figure 2), which provides the Pearson correlations between all pairs of AERONET variables grouped by a decision tree. To avoid collinearity in the regression models, we kept the most relevant of a group of variables that had a correlation coefficient > 0.9. We then examined and picked a subset of variables connected at the mid-tier level of the tree to construct interactions. We fit simple linear regression models separately for PM_{2.5} mass, sulfate, nitrate, EC, OC, and dust with AOD 550 nm as the sole predictor variable. Multiple linear regression models were again fit separately for PM_{2.5} mass, sulfate, nitrate, EC, OC, and dust, but with the combined AERONET sunphotometer and inversion product as predictor variables and model selection was conducted using the "all possible subset method". This method constructs models based on all combinations from 1 to k variable models. We select the best model from the combinations based on highest R², lowest RMSE, and Mallows' Cp statistic that is close to k+1. Model selection for the Fresno and Bakersfield sites were examined separately and in combination in a "total CA" analysis, which combined data from Fresno, Bakersfield, Modesto, Visalia, and a special DRAGON campaign in late 2012-early 2013 over the region (8 co-located sites with PM_{2.5} mass).

All models were cross validated (CV) with 10-fold CV, and we report the CV R² and RMSE. Models were fit in R using the leaps() library.

4 Results

5 Content section with R code chunks

You should always use `echo = FALSE` on R Markdown code blocks as they add formatting and styling not desired by Copernicus. The hidden workflow results in 42.

You can add verbatim code snippets without extra styles by using `` `` `` without additional instructions.

```
sum <- 1 + 41
```

6 Content section with list

If you want to insert a list, you must

- leave
- empty lines
- between each list item

because the `\tightlist` format used by R Markdown is not supported in the Copernicus template. Example:

- leave
- empty lines
- between each list item

7 Examples from the official template

7.1 FIGURES

When figures and tables are placed at the end of the MS (article in one-column style), please add




Figure 1. one column figure

between bibliography and first table and/or figure as well as between each table and/or figure.

7.1.1 ONE-COLUMN FIGURES

Include a 12cm width figure of Nikolaus Copernicus from Wikipedia with caption using R Markdown.

7.1.2 TWO-COLUMN FIGURES

- 5 You can also include a larger figure.

7.2 TABLES

You can add `\LaTeXtable` in an R Markdown document to meet the template requirements.




Figure 2. two column figure

Table 1. TEXT

a	b	c
1	2	3

Table Footnotes

Table 2. TEXT

a	b	c
1	2	3

Table footnotes

7.2.1 ONE-COLUMN TABLE

7.2.2 TWO-COLUMN TABLE

7.3 MATHEMATICAL EXPRESSIONS

All papers typeset by Copernicus Publications follow the math typesetting regulations given by the IUPAC Green Book (IU-

5 PAC: Quantities, Units and Symbols in Physical Chemistry, 2nd Edn., Blackwell Science, available at: <http://old.iupac.org/publications/book1993>).

Physical quantities/variables are typeset in italic font (t for time, T for Temperature)

Indices which are not defined are typeset in italic font (x, y, z, a, b, c)

Items/objects which are defined are typeset in roman font (Car A, Car B)

10 Descriptions/specifications which are defined by itself are typeset in roman font (abs, rel, ref, tot, net, ice)

Abbreviations from 2 letters are typeset in roman font (RH, LAI)

Vectors are identified in bold italic font using \boldsymbol{x}

Matrices are identified in bold roman font

Multiplication signs are typeset using the LaTeX commands `\times` (for vector products, grids, and exponential notations)

15 or `\cdot`

The character `*` should not be applied as multiplication sign

7.4 EQUATIONS

7.4.1 Single-row equation

Unnumbered equations (i.e. using `$$` and getting inline preview in RStudio) are not supported by Copernicus.

$$1 \times 1 \cdot 1 = 42 \tag{1}$$

5 $A = \pi r^2$ (2)

$$x = \frac{2b \pm \sqrt{b^2 - 4ac}}{2c}. \tag{3}$$

7.4.2 Multiline equation

$$3 + 5 = 8 \tag{4}$$

$$3 + 5 = 8 \tag{5}$$

10 $3 + 5 = 8$ (6)

7.5 MATRICES

$$\begin{matrix} x & y & z \end{matrix}$$

$$\begin{matrix} x & y & z \end{matrix}$$

$$\begin{matrix} x & y & z \end{matrix}$$

7.6 ALGORITHM

If you want to use algorithms, you can either enable the required packages in the header (the default, see `algorithms: true`), or make sure yourself that the `LaTeX` packages `algorithms` and `algorithmicx` are installed so that `algorithm.sty` respectively `algorithmic.sty` can be loaded by the Copernicus template. Copernicus staff will remove all undesirable packages from your LaTeX source code, so please stick to using the header option, which only adds the two acceptable packages.

7.7 CHEMICAL FORMULAS AND REACTIONS

20 For formulas embedded in the text, please use `\chem{ }`, e.g. $A \rightarrow B$.


The reaction environment creates labels including the letter R, i.e. (R1), (R2), etc.

```
i ← 10
if i ≥ 5 then
  i ← i − 1
else
  if i ≤ 3 then
    i ← i + 2
  end if
end if
```

- \rightarrow should be used for normal (one-way) chemical reactions
- \rightleftharpoons should be used for equilibria
- \leftrightarrow should be used for resonance structures



5



7.8 PHYSICAL UNITS

10 Please use \unit{} (allows to save the math/\$ environment) and apply the exponential notation, for example 3.14km h^{−1} (using LaTeX mode: \ (3.14\, \unit{...} \)) or 0.872ms^{−1} (using only \unit{0.872\,m\,s^{−1}}).

8 Conclusions

The conclusion goes here. You can modify the section name with \conclusions[modified heading if necessary].

Code and data availability. use this to add a statement when having data sets and software code available

15 *Sample availability.* use this section when having geoscientific samples available

Appendix A: Figures and tables in appendices

Regarding figures and tables in appendices, the following two options are possible depending on your general handling of figures and tables in the manuscript environment:

A1 Option 1

- 5 If you sorted all figures and tables into the sections of the text, please also sort the appendix figures and appendix tables into the respective appendix sections. They will be correctly named automatically.

A2 Option 2

If you put all figures after the reference list, please insert appendix tables and figures after the normal tables and figures.

- To rename them correctly to A1, A2, etc., please add the following commands in front of them: `\appendixfigures`
10 needs to be added in front of appendix figures `\appendixtables` needs to be added in front of appendix tables

Please add `\clearpage` between each table and/or figure. Further guidelines on figures and tables can be found below.

Author contributions. M. Franklin conducted analyses and wrote the manuscript. M. Sorek-Hamer conducted analyses and reviewed the manuscript. O. Kalashnikova and D. Diner conceptualized the study. D. Diner edited the manuscript.

Competing interests. The authors declare no competing interests.

- 15 *Disclaimer.* We like Copernicus.

Acknowledgements. This work was supported by NASA Grant 80NSSC19K0225

References