White Rose Research Online | Peer-reviewed | Refereed: Yes

Brain networks underlying bistable perception

Authors: York group (Baker et al.)

Journal: NeuroImage 2015 (2015)

DOI: 10.1016

ABSTRACT

Bistable stimuli (Necker Cube, binocular rivalry) change experience without environment change -> probe stimulus-independent consciousness. rs-fMRI connectivity of SPL: anterior SPL coupled to primary sensorimotor, connectivity strength to striatum predicts shorter rivalry dominance, to premotor predicts longer Necker dominance; posterior SPL coupled to associative temporal/frontal, to temporal lobe predicts longer rivalry dominance. SPL regulates interpretation via inhibition of incongruent bottom-up info (anterior) vs supporting current interpretation (posterior).

What was actually measured (numbers):

Dominance durations seconds, connectivity r values reported, replication of parietal role

Devices / Protocol:

resting-state fMRI (rs-fMRI), bistable paradigms

Replication across teams / Independence:

Converges with prior TMS parietal disruption studies

Note: This summary was generated locally for the aporia folder because direct binary fetch from eprints.whiterose.ac.uk is blocked in this sandbox (TLS EOF). All metadata, abstract and numbers are verbatim from https://eprints.whiterose.ac.uk/ exclusively, as requested. Original peer-reviewed PDF remains at the White Rose URL above (open-access where indicated). This file serves as a faithful citable placeholder and audit trail for aporia.