

# Elucidating the patterns of pleiotropy and its biological relevance in maize




Merritt Khaipho-Burch <sup>1</sup>; Taylor Ferebee <sup>2</sup>; Anju Giri <sup>3</sup>; Guillaume Ramstein <sup>3,4</sup>; Brandon Monier <sup>3</sup>; Emily Yi <sup>3</sup>; M. Cinta Romay <sup>3</sup>; Edward Buckler <sup>1,3,5</sup>


<sup>1</sup>) Section of Plant Breeding and Genetics, Cornell University, Ithaca, NY; <sup>2</sup>) Department of Computational Biology, Cornell University, Ithaca, NY; <sup>3</sup>) Institute for Genomic Diversity, Cornell University, Ithaca, NY; <sup>4</sup>) Center for Quantitative Genetics and Genomics, Aarhus University, Aarhus, Denmark; <sup>5</sup>) USDA-ARS; Ithaca, NY

Pleiotropy: a single locus that affects multiple traits. How pleiotropic is any maize locus?


Many forms of pleiotropy exist:



Few examples of pleiotropy have been described and verified in maize:



GWAS on 120,597 field, metabolite, and expression traits uncovers the patterns of pleiotropy.





68-98% of intervals are involved with 0-1 traits.


2-31% of intervals are involved with 2+ traits.

Much of this observed pleiotropy may be spurious.

Random forest models show high importance for nuisance terms, low importance for biological terms.



The distribution of pleiotropy differs by population and trait.



Pleiotropy estimates are highly biased towards nuisance terms and investigator bias. Caution & validation should be applied.



Want to know more?  
Take a FREE zine about  
this poster!



National Institute of Food and Agriculture  
U.S. DEPARTMENT OF AGRICULTURE

Thank you to our funding sources: NSF PGRP: IOS#1822330, the USDA ARS, and the USDA NIFA AFRI Pre-doctoral fellowship: 2022-67011-36458.

Preprint coming soon! **bioRxiv**  
THE PREPRINT SERVER FOR BIOLOGY