

BMA

Hoff Chapter 9, Liang et al 2007, Hoeting et al (1999), Clyde
& George (2004) Statistical Science

October 29, 2015

Prior Distributions

- ▶ “Spike and Slab” - Lempers (1971) Mitchell & Beauchamp (1988)

Prior Distributions

- ▶ “Spike and Slab” - Lempers (1971) Mitchell & Beauchamp (1988)
- ▶ “Spike and Bell” Leamer (1978) in BMA

Prior Distributions

- ▶ “Spike and Slab” - Lempers (1971) Mitchell & Beauchamp (1988)
- ▶ “Spike and Bell” Leamer (1978) in BMA
- ▶ mixture of 2 normals - concentrated and dispersed - SSVS
Gibbs Sampler - George & McCulloch (1993)

Prior Distributions

- ▶ “Spike and Slab” - Lempers (1971) Mitchell & Beauchamp (1988)
- ▶ “Spike and Bell” Leamer (1978) in BMA
- ▶ mixture of 2 normals - concentrated and dispersed - SSVS Gibbs Sampler - George & McCulloch (1993)
- ▶ Back to “Spike and Bell” Hoeting, Raftery & Madigan MC^3 (1997) and George (1997) collapsed MCMC integrate out β_γ
- ▶ Conjugate Normal-Gammas lead to closed form expressions for marginal likelihoods, Zellner's g-prior is the most popular.

Prior Distributions

- ▶ “Spike and Slab” - Lempers (1971) Mitchell & Beauchamp (1988)
- ▶ “Spike and Bell” Leamer (1978) in BMA
- ▶ mixture of 2 normals - concentrated and dispersed - SSVS Gibbs Sampler - George & McCulloch (1993)
- ▶ Back to “Spike and Bell” Hoeting, Raftery & Madigan MC^3 (1997) and George (1997) collapsed MCMC integrate out β_γ
- ▶ Conjugate Normal-Gammas lead to closed form expressions for marginal likelihoods, Zellner's g-prior is the most popular.

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

- ▶ $p(\alpha) \propto 1$

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

- ▶ $p(\alpha) \propto 1$
- ▶ $p(\sigma^2) \propto 1/\sigma^2$

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

- ▶ $p(\alpha) \propto 1$
- ▶ $p(\sigma^2) \propto 1/\sigma^2$
- ▶ $\boldsymbol{\beta}_\gamma \mid \alpha, \sigma^2, \gamma \sim N(0, g\sigma^2(\mathbf{X}_\gamma^c \mathbf{X}_\gamma^c)^{-1})$

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

- ▶ $p(\alpha) \propto 1$
- ▶ $p(\sigma^2) \propto 1/\sigma^2$
- ▶ $\boldsymbol{\beta}_\gamma \mid \alpha, \sigma^2, \gamma \sim N(0, g\sigma^2(\mathbf{X}_\gamma^c' \mathbf{X}_\gamma^c)^{-1})$

which leads to marginal likelihood of \mathcal{M}_γ that is proportional to

$$p(\mathbf{Y} \mid \mathcal{M}_\gamma) = C(1 + g)^{\frac{n-p-1}{2}} (1 + g(1 - R_\gamma^2))^{-\frac{(n-1)}{2}}$$

where R^2 is the usual R^2 for model \mathcal{M}_γ .

Zellner's g-prior

Centered model:

$$\mathbf{Y} = \mathbf{1}_n \alpha + \mathbf{X}^c \boldsymbol{\beta} + \epsilon$$

where \mathbf{X}^c is the centered design matrix where all variables have had their mean subtracted

- ▶ $p(\alpha) \propto 1$
- ▶ $p(\sigma^2) \propto 1/\sigma^2$
- ▶ $\boldsymbol{\beta}_\gamma \mid \alpha, \sigma^2, \gamma \sim N(0, g\sigma^2(\mathbf{X}_\gamma^c' \mathbf{X}_\gamma^c)^{-1})$

which leads to marginal likelihood of \mathcal{M}_γ that is proportional to

$$p(\mathbf{Y} \mid \mathcal{M}_\gamma) = C(1 + g)^{\frac{n-p-1}{2}} (1 + g(1 - R_\gamma^2))^{-\frac{(n-1)}{2}}$$

where R^2 is the usual R^2 for model \mathcal{M}_γ .

Trade-off of model complexity versus goodness of fit

Lastly, assign distribution to space of models

Priors on Model Space

$$p(\mathcal{M}_\gamma) \Leftrightarrow p(\gamma)$$

- ▶ $p(\gamma_j = 1) = .5 \Rightarrow P(\mathcal{M}_\gamma) = .5^p$ Uniform on space of models

Priors on Model Space

$$p(\mathcal{M}_\gamma) \Leftrightarrow p(\gamma)$$

- ▶ $p(\gamma_j = 1) = .5 \Rightarrow P(\mathcal{M}_\gamma) = .5^p$ Uniform on space of models
 $p_\gamma \sim \text{Bin}(p, .5)$
- ▶ $\gamma_j \mid \pi \stackrel{\text{iid}}{\sim} \text{Ber}(\pi)$ and $\pi \sim \text{Beta}(a, b)$ then $p_\gamma \sim \text{BB}(p, a, b)$

$$p(p_\gamma \mid p, a, b) = \frac{\Gamma(p+1)\Gamma(p_\gamma+a)\Gamma(p-p_\gamma+b)\Gamma(a+b)}{\Gamma(p_\gamma+1)\Gamma(p-p_\gamma+1)\Gamma(p+a+b)\Gamma(a)\Gamma(b)}$$

- ▶ $p_\gamma \sim \text{BB}(p, 1, 1) \sim \text{Unif}(0, p)$
- ▶ For sparsity. want b to be a function of p typically.

USair Data

```
library(BAS)
poll.bma = bas.lm(log(SO2) ~ temp + log(mgffirms) +
                  log(popn) + wind +
                  precip+ raindays,
                  data=pollution,
                  prior="g-prior",
                  alpha=41, # n
                  n.models=2^6,
                  modelprior = uniform(),
                  initprobs="Uniform")
```





```
> poll.bma
```

Marginal Posterior Inclusion Probabilities:

Intercept	temp	log(mgffirms)	log(popn)	wind	precip
1.0000	0.9755	0.7190	0.2757	0.7654	0.5994


Plots

`plot(poll.bma, ask=F)`




Posterior Distribution with Uniform Prior on Model Space

image(poll.bma)



Posterior Distribution with BB(1,p) Prior on Model Space

image(poll-bb.bma)



Jeffreys Scale of Evidence

- ▶ Bayes Factor = ratio of marginal likelihoods

Jeffreys Scale of Evidence

- ▶ Bayes Factor = ratio of marginal likelihoods
- ▶ Posterior odds = Bayes Factor \times Prior odds

Jeffreys Scale of Evidence

- ▶ Bayes Factor = ratio of marginal likelihoods
- ▶ Posterior odds = Bayes Factor \times Prior odds

$$B = BF[\mathcal{M}_0 : \mathcal{M}_\gamma] \text{ and } 1/B = BF[\mathcal{M}_\gamma : \mathcal{M}_0]$$

Jeffreys Scale of Evidence

- ▶ Bayes Factor = ratio of marginal likelihoods
- ▶ Posterior odds = Bayes Factor \times Prior odds

$$B = BF[\mathcal{M}_0 : \mathcal{M}_\gamma] \text{ and } 1/B = BF[\mathcal{M}_\gamma : \mathcal{M}_0]$$

Bayes Factor	Interpretation
$B \geq 1$	H_0 supported
$1 > B \geq 10^{-\frac{1}{2}}$	minimal evidence against H_0
$10^{-\frac{1}{2}} > B \geq 10^{-1}$	substantial evidence against H_0
$10^{-1} > B \geq 10^{-2}$	strong evidence against H_0
$10^{-2} > B$	decisive evidence against H_0

Jeffreys Scale of Evidence

- ▶ Bayes Factor = ratio of marginal likelihoods
- ▶ Posterior odds = Bayes Factor \times Prior odds

$$B = BF[\mathcal{M}_0 : \mathcal{M}_\gamma] \text{ and } 1/B = BF[\mathcal{M}_\gamma : \mathcal{M}_0]$$


Bayes Factor	Interpretation
$B \geq 1$	H_0 supported
$1 > B \geq 10^{-\frac{1}{2}}$	minimal evidence against H_0
$10^{-\frac{1}{2}} > B \geq 10^{-1}$	substantial evidence against H_0
$10^{-1} > B \geq 10^{-2}$	strong evidence against H_0
$10^{-2} > B$	decisive evidence against H_0

in context

of testing one hypothesis with equal prior odds

Coefficients

```
beta = coef(poll.bma)  
par(mfrow=c(2,3)); plot(beta, subset=2:7, ask=F)
```



Information Paradox

The Bayes factor for comparing \mathcal{M}_γ to the null model:

$$BF(\mathcal{M}_\gamma : \mathfrak{M}_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R^2))^{-(n-1)/2}$$

Information Paradox

The Bayes factor for comparing \mathcal{M}_γ to the null model:

$$BF(\mathcal{M}_\gamma : \mathcal{M}_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R^2))^{-(n-1)/2}$$

- ▶ Let g be a fixed constant and take n fixed.

Information Paradox

The Bayes factor for comparing \mathcal{M}_γ to the null model:

$$BF(\mathcal{M}_\gamma : \mathcal{M}_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R^2))^{-(n-1)/2}$$

- ▶ Let g be a fixed constant and take n fixed.
- ▶ Let $F = \frac{R_\gamma^2 / p_\gamma}{(1 - R_\gamma^2) / (n - 1 - p_\gamma)}$

Information Paradox

The Bayes factor for comparing \mathcal{M}_γ to the null model:

$$BF(\mathcal{M}_\gamma : \mathcal{M}_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R^2))^{-(n-1)/2}$$

- ▶ Let g be a fixed constant and take n fixed.
- ▶ Let $F = \frac{R_\gamma^2 / p_\gamma}{(1 - R_\gamma^2) / (n - 1 - p_\gamma)}$
- ▶ As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject \mathcal{M}_0 where F is the usual F statistic for comparing model \mathcal{M}_γ to \mathcal{M}_0

Information Paradox

The Bayes factor for comparing \mathcal{M}_γ to the null model:

$$BF(\mathcal{M}_\gamma : \mathcal{M}_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R^2))^{-(n-1)/2}$$

- ▶ Let g be a fixed constant and take n fixed.
- ▶ Let $F = \frac{R_\gamma^2 / p_\gamma}{(1 - R_\gamma^2) / (n - 1 - p_\gamma)}$
- ▶ As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject \mathcal{M}_0 where F is the usual F statistic for comparing model \mathcal{M}_γ to \mathcal{M}_0
- ▶ BF converges to a fixed constant $(1 + g)^{-p_\gamma/2}$ (does not go to infinity)

“Information Inconsistency” see Liang et al JASA 2008

Mixtures of g priors & Information consistency

Need $BF \rightarrow \infty$ if $R^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges (proof in Liang et al)

Mixtures of g priors & Information consistency

Need $BF \rightarrow \infty$ if $R^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges (proof in Liang et al)

- ▶ Zellner-Siow Cauchy prior

Mixtures of g priors & Information consistency

Need $BF \rightarrow \infty$ if $R^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges (proof in Liang et al)

- ▶ Zellner-Siow Cauchy prior
- ▶ hyper- g prior or hyper- g/n (Liang et al JASA 2008)

Mixtures of g priors & Information consistency

Need $BF \rightarrow \infty$ if $R^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges (proof in Liang et al)

- ▶ Zellner-Siow Cauchy prior
- ▶ hyper- g prior or hyper- g/n (Liang et al JASA 2008)
- ▶ robust prior (Bayarri et al Annals of Statistics 2012)

Mixtures of g priors & Information consistency

Need $BF \rightarrow \infty$ if $R^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges (proof in Liang et al)

- ▶ Zellner-Siow Cauchy prior
- ▶ hyper- g prior or hyper- g/n (Liang et al JASA 2008)
- ▶ robust prior (Bayarri et al Annals of Statistics 2012)

All have tails that behave like a Cauchy distribution

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities
- ▶ response Mortality

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities
- ▶ response Mortality
- ▶ measures of HC, NOX, SO2

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities
- ▶ response Mortality
- ▶ measures of HC, NOX, SO2
- ▶ Is pollution associated with mortality after adjusting for other socio-economic and meteorological factors?

Mortality & Pollution





- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities
- ▶ response Mortality
- ▶ measures of HC, NOX, SO2
- ▶ Is pollution associated with mortality after adjusting for other socio-economic and meteorological factors?
- ▶ 15 predictor variables implies $2^{15} = 32,768$ possible models

Mortality & Pollution

- ▶ Data from Statistical Sleuth 12.17
- ▶ 60 cities
- ▶ response Mortality
- ▶ measures of HC, NOX, SO2
- ▶ Is pollution associated with mortality after adjusting for other socio-economic and meteorological factors?
- ▶ 15 predictor variables implies $2^{15} = 32,768$ possible models
- ▶ Use Zellner-Siow Cauchy prior $1/g \sim G(1/2, n/2)$

```
mort.bma = bas.lm(MORTALITY ~ ., data=mortality,  
                  prior="ZS-null",  
                  alpha=60, n.models=2^15,  
                  update=100, initprobs="eplogp")
```

Posterior Distributions



Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0
> sum(poll.in * mort.bma$postprob)
[1] 0.9889641
```
- ▶ Posterior probability is 0.011

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```

- ▶ Posterior probability is 0.011
- ▶ Odds that there is an effect $(1 - .011)/(.011) = 89$.

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```

- ▶ Posterior probability is 0.011
- ▶ Odds that there is an effect $(1 - .011)/(.011) = 89$.
- ▶ Prior Odds $7 = (1 - .5^3)/.5^3$

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```

- ▶ Posterior probability is 0.011
- ▶ Odds that there is an effect $(1 - .011)/(.011) = 89$.
- ▶ Prior Odds $7 = (1 - .5^3)/.5^3$
- ▶ Bayes Factor for a pollution effect $89.9/7 = 12.8$

Posterior Probabilities

- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```

- ▶ Posterior probability is 0.011
- ▶ Odds that there is an effect $(1 - .011)/(.011) = 89$.
- ▶ Prior Odds $7 = (1 - .5^3)/.5^3$
- ▶ Bayes Factor for a pollution effect $89.9/7 = 12.8$
- ▶ Bayes Factor for NOX based on marginal inclusion probability $0.917/(1 - 0.917) = 11.0$

Posterior Probabilities


- ▶ What is the probability that there is no pollution effect?
- ▶ Sum posterior model probabilities over all models that include no pollution variables

```
> which.mat = list2matrix.which(mort.bma,1:(2^15))  
> poll.in = (which.mat[, 14:16] %*% rep(1, 3)) > 0  
> sum(poll.in * mort.bma$postprob)  
[1] 0.9889641
```









- ▶ Posterior probability is 0.011
- ▶ Odds that there is an effect $(1 - .011)/(.011) = 89$.
- ▶ Prior Odds $7 = (1 - .5^3)/.5^3$
- ▶ Bayes Factor for a pollution effect $89.9/7 = 12.8$
- ▶ Bayes Factor for NOX based on marginal inclusion probability $0.917/(1 - 0.917) = 11.0$
- ▶ Marginal inclusion probability for logHC = 0.427144
- ▶ Marginal inclusion probability for logSO2 = 0.218978

Bayes Factors are not additive! Better to work with probabilities.









Model Space



Coefficients



Coefficients



Effect Estimation

- ▶ Coefficients in each model are adjusted for other variables in the model
- ▶ OLS: leave out a predictor with a non-zero coefficient then estimates are biased!
- ▶ Model Selection in the presence of high correlation, may leave out "redundant" variables;
- ▶ improved MSE for prediction (Bias-variance tradeoff)
- ▶ Bayes is biased anyway so should we care?
- ▶ What is meaning of $\sum_{\gamma} \beta_{j\gamma} \gamma_j P(\mathcal{M}_{\gamma} | \mathbf{Y})$

Problem with confounding! Need to change prior?

Other Problems

- ▶ Computational

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations
 - ▶ Metropolis Hastings algorithms more flexibility

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations
 - ▶ Metropolis Hastings algorithms more flexibility
 - ▶ "Stochastic Search" (no guarantee samples represent posterior)

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations
 - ▶ Metropolis Hastings algorithms more flexibility
 - ▶ "Stochastic Search" (no guarantee samples represent posterior)
 - ▶ in BMA all variables are included, but coefficients are shrunk to 0; alternative is to use Shrinkage methods

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations
 - ▶ Metropolis Hastings algorithms more flexibility
 - ▶ "Stochastic Search" (no guarantee samples represent posterior)
 - ▶ in BMA all variables are included, but coefficients are shrunk to 0; alternative is to use Shrinkage methods
- ▶ Prior Choice: Choice of prior distributions on β and on γ

Other Problems

- ▶ Computational if $p > 35$ enumeration is difficult
 - ▶ Gibbs sampler or Random-Walk algorithm on γ
 - ▶ poor convergence/mixing with high correlations
 - ▶ Metropolis Hastings algorithms more flexibility
 - ▶ "Stochastic Search" (no guarantee samples represent posterior)
 - ▶ in BMA all variables are included, but coefficients are shrunk to 0; alternative is to use Shrinkage methods
- ▶ Prior Choice: Choice of prior distributions on β and on γ

Model averaging versus Model Selection – what are objectives?