

# Maximum Likelihood Estimation

Merlise Clyde

STA721 Linear Models

Duke University

September 2, 2014

## Topics

- Projections
- Maximum Likelihood Estimates
- Spectral Decomposition

Readings: Continue reading Wakefield 5.6.1 or for more details Christensen Chapter 1-2, Appendix A, and Appendix B

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $P_{\mathbf{X}}\mathbf{Y}$
- $P_{\mathbf{X}}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$
- $P = P^2$  (idempotent)

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\end{aligned}$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned} \mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X} \mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X} \end{aligned}$$



# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned} \mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X} \mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X} \end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\mathbf{P}_\mathbf{X}^T = (\mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T((\mathbf{X}^T\mathbf{X})^{-1})^T(\mathbf{X})^T\end{aligned}$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T((\mathbf{X}^T\mathbf{X})^{-1})^T(\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\end{aligned}$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T((\mathbf{X}^T\mathbf{X})^{-1})^T(\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

# Models & MLEs

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X}\mathbf{Y}$
- $\mathbf{P}_\mathbf{X}$  is the orthogonal projection operator on the column space of  $\mathbf{X}$ ; e.g.  $\mathbf{X}$  full rank  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$
- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T((\mathbf{X}^T\mathbf{X})^{-1})^T(\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $C(\mathbf{X}) = C(\mathbf{P}_\mathbf{X})$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent



Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}})$$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 &= (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \\ &= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\mathbf{P}_{\mathbf{X}}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 &= (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 &= (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 &= (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_{\mathbf{X}} = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})^T$

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_\mathbf{X} = (\mathbf{I} - \mathbf{P}_\mathbf{X})^T$
- $\mathbf{u} \in C(\mathbf{X})^\perp \Rightarrow \mathbf{u} \perp C(\mathbf{X})$  and  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{u} = \mathbf{u}$  (projection)

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_\mathbf{X} = (\mathbf{I} - \mathbf{P}_\mathbf{X})^T$
- $\mathbf{u} \in C(\mathbf{X})^\perp \Rightarrow \mathbf{u} \perp C(\mathbf{X})$  and  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{u} = \mathbf{u}$  (projection)
- if  $\mathbf{v} \in C(\mathbf{X})$ ,  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{v} = \mathbf{v} - \mathbf{v} = \mathbf{0}$

- Log Likelihood:



- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_X \mathbf{Y} + (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}$

- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_X \mathbf{Y} + (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}$
- Use  $\mathbf{P}_X \boldsymbol{\mu} = \boldsymbol{\mu}$

- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_X \mathbf{Y} + (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}$
- Use  $\mathbf{P}_X \boldsymbol{\mu} = \boldsymbol{\mu}$
- and Simplify  $\|\mathbf{Y} - \boldsymbol{\mu}\|^2$

# Expand

$$\|\mathbf{Y} - \boldsymbol{\mu}\|^2 = \|\mathbf{Y} - \mathbf{P}_x \mathbf{Y} + \mathbf{P}_x \mathbf{Y} - \boldsymbol{\mu}\|^2$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2 \\ &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \textcolor{red}{P}_\mathbf{X}\boldsymbol{\mu}\|^2\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2 \\ &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{P}_\mathbf{X}\boldsymbol{\mu}\|^2 \\ &= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y} + \mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2\end{aligned}$$



# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\ &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\ &= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\ &= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

Crossproduct term is zero

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

Crossproduct term is zero

$$\mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) = \mathbf{P}_X (\mathbf{I} - \mathbf{P}_X)$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_X \mathbf{Y} + \mathbf{P}_X \mathbf{Y} - \mathbf{P}_X \boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y} + \mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X (\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X \mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

Crossproduct term is zero

$$\begin{aligned}\mathbf{P}_X^T (\mathbf{I} - \mathbf{P}_X) &= \mathbf{P}_X (\mathbf{I} - \mathbf{P}_X) \\&= \mathbf{P}_X - \mathbf{P}_X \mathbf{P}_X\end{aligned}$$

# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{P}_\mathbf{X}\boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y} + \mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_\mathbf{X}^T (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

Crossproduct term is zero

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) &= \mathbf{P}_\mathbf{X} (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} \mathbf{P}_\mathbf{X} \\&= \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X}\end{aligned}$$



# Expand

$$\begin{aligned}\|\mathbf{Y} - \boldsymbol{\mu}\|^2 &= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2 \\&= \|\mathbf{Y} - \mathbf{P}_\mathbf{X}\mathbf{Y} + \mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{P}_\mathbf{X}\boldsymbol{\mu}\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y} + \mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 + 2(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{P}_\mathbf{X}^T (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y} \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}(\mathbf{Y} - \boldsymbol{\mu})\|^2 + 0 \\&= \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2\end{aligned}$$

Crossproduct term is zero

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) &= \mathbf{P}_\mathbf{X} (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} \mathbf{P}_\mathbf{X} \\&= \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} \\&= 0\end{aligned}$$

Substitute decomposition into log likelihood

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\ &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right)\end{aligned}$$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\text{}}\end{aligned}$$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} \\&= \text{constant with respect to } \boldsymbol{\mu}\end{aligned}$$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\leq 0} \\&= \text{constant with respect to } \boldsymbol{\mu} \quad \leq 0\end{aligned}$$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\leq 0} \\&= \text{constant with respect to } \boldsymbol{\mu} \leq 0\end{aligned}$$

Maximize with respect to  $\boldsymbol{\mu}$  for each  $\sigma^2$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\leq 0} \\&= \text{constant with respect to } \boldsymbol{\mu} \leq 0\end{aligned}$$

Maximize with respect to  $\boldsymbol{\mu}$  for each  $\sigma^2$

RHS is largest when  $\boldsymbol{\mu} = \mathbf{P}_\mathbf{X}\mathbf{Y}$  for any choice of  $\sigma^2$



Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\leq 0} \\&= \text{constant with respect to } \boldsymbol{\mu} \leq 0\end{aligned}$$

Maximize with respect to  $\boldsymbol{\mu}$  for each  $\sigma^2$

RHS is largest when  $\boldsymbol{\mu} = \mathbf{P}_\mathbf{X}\mathbf{Y}$  for any choice of  $\sigma^2$

$$\therefore \hat{\boldsymbol{\mu}} = \mathbf{P}_\mathbf{X}\mathbf{Y}$$

is the MLE of  $\boldsymbol{\mu}$

Substitute decomposition into log likelihood

$$\begin{aligned}\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \\&= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\&= \underbrace{-\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2}}_{\text{constant with respect to } \boldsymbol{\mu}} + \underbrace{-\frac{1}{2} \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}}_{\leq 0} \\&= \text{constant with respect to } \boldsymbol{\mu} \leq 0\end{aligned}$$

Maximize with respect to  $\boldsymbol{\mu}$  for each  $\sigma^2$

RHS is largest when  $\boldsymbol{\mu} = \mathbf{P}_\mathbf{X}\mathbf{Y}$  for any choice of  $\sigma^2$

$$\therefore \hat{\boldsymbol{\mu}} = \mathbf{P}_\mathbf{X}\mathbf{Y}$$

is the MLE of  $\boldsymbol{\mu}$  (yields fitted values  $\hat{\mathbf{Y}} = \mathbf{P}_\mathbf{X}\mathbf{Y}$ )

$$\mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right)$$

$$\begin{aligned}\mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\ \mathcal{L}(\boldsymbol{\beta}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}\|^2}{\sigma^2} \right)\end{aligned}$$

$$\begin{aligned}\mathcal{L}(\boldsymbol{\mu}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right) \\ \mathcal{L}(\boldsymbol{\beta}, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}\|^2}{\sigma^2} \right)\end{aligned}$$

Similar argument to show that RHS is maximized by minimizing

$$\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}\|^2$$

$$\begin{aligned}\mathcal{L}(\mu, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mu\|^2}{\sigma^2} \right) \\ \mathcal{L}(\beta, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\beta\|^2}{\sigma^2} \right)\end{aligned}$$

Similar argument to show that RHS is maximized by minimizing

$$\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\beta\|^2$$

Therefore  $\hat{\beta}$  is a MLE of  $\beta$  if and only if satisfies

$$\mathbf{P}_\mathbf{X}\mathbf{Y} = \mathbf{X}\hat{\beta}$$

$$\begin{aligned}\mathcal{L}(\mu, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mu\|^2}{\sigma^2} \right) \\ \mathcal{L}(\beta, \sigma^2) &= -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \left( \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{\sigma^2} + \frac{\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\beta\|^2}{\sigma^2} \right)\end{aligned}$$

Similar argument to show that RHS is maximized by minimizing

$$\|\mathbf{P}_\mathbf{X}\mathbf{Y} - \mathbf{X}\beta\|^2$$

Therefore  $\hat{\beta}$  is a MLE of  $\beta$  if and only if satisfies

$$\mathbf{P}_\mathbf{X}\mathbf{Y} = \mathbf{X}\hat{\beta}$$

If  $\mathbf{X}^T\mathbf{X}$  is full rank, the MLE of  $\beta$  is

$$(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y} = \hat{\beta}$$

- Plug-in MLE of  $\hat{\mu}$  for  $\mu$  and differentiate with respect to  $\sigma^2$



- Plug-in MLE of  $\hat{\boldsymbol{\mu}}$  for  $\boldsymbol{\mu}$  and differentiate with respect to  $\sigma^2$

$$\log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2) = -\frac{n}{2} \log \sigma^2 - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{\sigma^2}$$

- Plug-in MLE of  $\hat{\boldsymbol{\mu}}$  for  $\boldsymbol{\mu}$  and differentiate with respect to  $\sigma^2$

$$\begin{aligned}\log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2) &= -\frac{n}{2} \log \sigma^2 - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{\sigma^2} \\ \frac{\partial \log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2)}{\partial \sigma^2} &= -\frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\sigma^2} \right)^2\end{aligned}$$

- Plug-in MLE of  $\hat{\boldsymbol{\mu}}$  for  $\boldsymbol{\mu}$  and differentiate with respect to  $\sigma^2$

$$\begin{aligned}\log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2) &= -\frac{n}{2} \log \sigma^2 - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{\sigma^2} \\ \frac{\partial \log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2)}{\partial \sigma^2} &= -\frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\sigma^2} \right)^2\end{aligned}$$

- Set derivative to zero and solve for MLE

$$0 = -\frac{n}{2} \frac{1}{\hat{\sigma}^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\hat{\sigma}^2} \right)^2$$

- Plug-in MLE of  $\hat{\boldsymbol{\mu}}$  for  $\boldsymbol{\mu}$  and differentiate with respect to  $\sigma^2$

$$\begin{aligned}\log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2) &= -\frac{n}{2} \log \sigma^2 - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{\sigma^2} \\ \frac{\partial \log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2)}{\partial \sigma^2} &= -\frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\sigma^2} \right)^2\end{aligned}$$

- Set derivative to zero and solve for MLE

$$\begin{aligned}0 &= -\frac{n}{2} \frac{1}{\hat{\sigma}^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\hat{\sigma}^2} \right)^2 \\ \frac{n}{2} \hat{\sigma}^2 &= \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2\end{aligned}$$

- Plug-in MLE of  $\hat{\boldsymbol{\mu}}$  for  $\boldsymbol{\mu}$  and differentiate with respect to  $\sigma^2$

$$\begin{aligned}\log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2) &= -\frac{n}{2} \log \sigma^2 - \frac{1}{2} \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{\sigma^2} \\ \frac{\partial \log \mathcal{L}(\hat{\boldsymbol{\mu}}, \sigma^2)}{\partial \sigma^2} &= -\frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\sigma^2} \right)^2\end{aligned}$$

- Set derivative to zero and solve for MLE

$$\begin{aligned}0 &= -\frac{n}{2} \frac{1}{\hat{\sigma}^2} + \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \left( \frac{1}{\hat{\sigma}^2} \right)^2 \\ \frac{n}{2} \hat{\sigma}^2 &= \frac{1}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2 \\ \hat{\sigma}^2 &= \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{n}\end{aligned}$$

Maximum Likelihood Estimate of  $\sigma^2$

$$\hat{\sigma}^2 = \frac{\|(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}\|^2}{n}$$

Maximum Likelihood Estimate of  $\sigma^2$

$$\begin{aligned}\hat{\sigma}^2 &= \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{n} \\ &= \frac{\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}}{n}\end{aligned}$$

Maximum Likelihood Estimate of  $\sigma^2$

$$\begin{aligned}\hat{\sigma}^2 &= \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{n} \\ &= \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}}{n} \\ &= \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}}{n}\end{aligned}$$



Maximum Likelihood Estimate of  $\sigma^2$

$$\begin{aligned}\hat{\sigma}^2 &= \frac{\|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2}{n} \\&= \frac{\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}}{n} \\&= \frac{\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}}{n} \\&= \frac{\mathbf{e}^T\mathbf{e}}{n}\end{aligned}$$

Maximum Likelihood Estimate of  $\sigma^2$

$$\begin{aligned}\hat{\sigma}^2 &= \frac{\|(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}\|^2}{n} \\&= \frac{\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}}{n} \\&= \frac{\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}}{n} \\&= \frac{\mathbf{e}^T\mathbf{e}}{n}\end{aligned}$$

where  $\mathbf{e} = (\mathbf{I} - \mathbf{P}_X)\mathbf{Y}$  **residuals** from the regression of  $\mathbf{Y}$  on  $\mathbf{X}$

# Geometric View

- Fitted Values  $\hat{\mathbf{Y}} = \mathbf{P}_{\mathbf{X}}\mathbf{Y} = \mathbf{X}\hat{\boldsymbol{\beta}}$

# Geometric View

- Fitted Values  $\hat{\mathbf{Y}} = \mathbf{P}_\mathbf{X}\mathbf{Y} = \mathbf{X}\hat{\boldsymbol{\beta}}$
- Residuals  $\mathbf{e} = (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}$

# Geometric View

- Fitted Values  $\hat{\mathbf{Y}} = \mathbf{P}_\mathbf{X}\mathbf{Y} = \mathbf{X}\hat{\boldsymbol{\beta}}$
- Residuals  $\mathbf{e} = (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}$
- $\mathbf{Y} = \hat{\mathbf{Y}} + \mathbf{e}$

# Geometric View


- Fitted Values  $\hat{\mathbf{Y}} = \mathbf{P}_\mathbf{X}\mathbf{Y} = \mathbf{X}\hat{\boldsymbol{\beta}}$
- Residuals  $\mathbf{e} = (\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}$
- $\mathbf{Y} = \hat{\mathbf{Y}} + \mathbf{e}$

$$\|\mathbf{Y}\|^2 = \|(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}\|^2 + \|\mathbf{P}_\mathbf{X}\mathbf{Y}\|^2$$

# Geometric View

- Fitted Values  $\hat{\mathbf{Y}} = \mathbf{P}_X \mathbf{Y} = \mathbf{X} \hat{\boldsymbol{\beta}}$
- Residuals  $\mathbf{e} = (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}$
- $\mathbf{Y} = \hat{\mathbf{Y}} + \mathbf{e}$

$$\|\mathbf{Y}\|^2 = \|(\mathbf{I} - \mathbf{P}_X) \mathbf{Y}\|^2 + \|\mathbf{P}_X \mathbf{Y}\|^2$$



# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$



# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$E[\hat{\mathbf{Y}}] = E[\mathbf{P}_\mathbf{X}\mathbf{Y}]$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \end{aligned}$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \end{aligned}$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$E[\mathbf{e}] = E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}]$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$\begin{aligned} E[\mathbf{e}] &= E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})E[\mathbf{Y}] \end{aligned}$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$\begin{aligned} E[\mathbf{e}] &= E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})E[\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \end{aligned}$$



# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$\begin{aligned} E[\mathbf{e}] &= E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})E[\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\ &= \mathbf{0} \end{aligned}$$

# Properties

$\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

$$\begin{aligned} E[\hat{\mathbf{Y}}] &= E[\mathbf{P}_\mathbf{X}\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}E[\mathbf{Y}] \\ &= \mathbf{P}_\mathbf{X}\boldsymbol{\mu} \\ &= \boldsymbol{\mu} \end{aligned}$$

$E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$\begin{aligned} E[\mathbf{e}] &= E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})E[\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\ &= \mathbf{0} \end{aligned}$$

Will not be  $\mathbf{0}$  if  $\boldsymbol{\mu} \notin C(\mathbf{X})$

# Estimate of $\sigma^2$

MLE of  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{\mathbf{e}^T \mathbf{e}}{n} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}}{n}$$

# Estimate of $\sigma^2$

MLE of  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{\mathbf{e}^T \mathbf{e}}{n} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}}{n}$$

Is this an unbiased estimate of  $\sigma^2$ ?

# Estimate of $\sigma^2$

MLE of  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{\mathbf{e}^T \mathbf{e}}{n} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_X) \mathbf{Y}}{n}$$

Is this an unbiased estimate of  $\sigma^2$ ?

Need expectations of quadratic forms  $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  for  $\mathbf{A}$  an  $n \times n$  matrix  
 $\mathbf{Y}$  a random vector in  $\mathbb{R}^n$

Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

- $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  is a scalar

Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

- $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  is a scalar
- $\mathbf{Y}^T \mathbf{A} \mathbf{Y} = (\mathbf{Y}^T \mathbf{A} \mathbf{Y})^T = \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}$



Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

- $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  is a scalar
- $\mathbf{Y}^T \mathbf{A} \mathbf{Y} = (\mathbf{Y}^T \mathbf{A} \mathbf{Y})^T = \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}$

$$\frac{\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}}{2} = \mathbf{Y}^T \mathbf{A} \mathbf{Y}$$

Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

- $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  is a scalar
- $\mathbf{Y}^T \mathbf{A} \mathbf{Y} = (\mathbf{Y}^T \mathbf{A} \mathbf{Y})^T = \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}$

$$\frac{\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}}{2} = \mathbf{Y}^T \mathbf{A} \mathbf{Y}$$

$$\mathbf{Y}^T \frac{(\mathbf{A} + \mathbf{A}^T)}{2} \mathbf{Y} = \mathbf{Y}^T \mathbf{A} \mathbf{Y}$$

Without loss of generality we can assume that  $\mathbf{A} = \mathbf{A}^T$

- $\mathbf{Y}^T \mathbf{A} \mathbf{Y}$  is a scalar
- $\mathbf{Y}^T \mathbf{A} \mathbf{Y} = (\mathbf{Y}^T \mathbf{A} \mathbf{Y})^T = \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}$

$$\frac{\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \mathbf{Y}^T \mathbf{A}^T \mathbf{Y}}{2} = \mathbf{Y}^T \mathbf{A} \mathbf{Y}$$

$$\mathbf{Y}^T \frac{(\mathbf{A} + \mathbf{A}^T)}{2} \mathbf{Y} = \mathbf{Y}^T \mathbf{A} \mathbf{Y}$$

- may take  $\mathbf{A} = \mathbf{A}^T$

# Expectations of Quadratic Forms

## Theorem

*Let  $\mathbf{Y}$  be a random vector in  $\mathbb{R}^n$  with  $E[\mathbf{Y}] = \boldsymbol{\mu}$  and  $\text{Cov}(\mathbf{Y}) = \boldsymbol{\Sigma}$ .*

# Expectations of Quadratic Forms

## Theorem

Let  $\mathbf{Y}$  be a random vector in  $\mathbb{R}^n$  with  $E[\mathbf{Y}] = \boldsymbol{\mu}$  and  $\text{Cov}(\mathbf{Y}) = \boldsymbol{\Sigma}$ .  
Then  $E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] = \text{tr} \mathbf{A} \boldsymbol{\Sigma} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu}$ .

# Expectations of Quadratic Forms

## Theorem

Let  $\mathbf{Y}$  be a random vector in  $\mathbb{R}^n$  with  $E[\mathbf{Y}] = \boldsymbol{\mu}$  and  $\text{Cov}(\mathbf{Y}) = \boldsymbol{\Sigma}$ .  
Then  $E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] = \text{tr} \mathbf{A} \boldsymbol{\Sigma} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu}$ .

Result useful for finding expected values of Mean Squares; no normality required!

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] = E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}]$$



# Proof

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

# Proof

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] = E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr} \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr} \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} E[\mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr} \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} E[\mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} \mathbf{A} E[(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$



Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr} \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} E[\mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} \mathbf{A} E[(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} \mathbf{A} \boldsymbol{\Sigma} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Start with  $(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})$ , expand and take expectations

$$\begin{aligned} E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \mathbf{Y} - \mathbf{Y}^T \mathbf{A} \boldsymbol{\mu}] \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] - \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

Rearrange

$$\begin{aligned} E[\mathbf{Y}^T \mathbf{A} \mathbf{Y}] &= E[(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= E[\text{tr} \mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} E[\mathbf{A}(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} \mathbf{A} E[(\mathbf{Y} - \boldsymbol{\mu})(\mathbf{Y} - \boldsymbol{\mu})^T] + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \\ &= \text{tr} \mathbf{A} \boldsymbol{\Sigma} + \boldsymbol{\mu}^T \mathbf{A} \boldsymbol{\mu} \end{aligned}$$

$$\text{tr} \mathbf{A} \equiv \sum_{i=1}^n a_{ii}$$

# Expectation of $\hat{\sigma}^2$

Use the theorem:

# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] = \text{tr}(\mathbf{I} - \mathbf{P}_X)\sigma^2 + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_X)\boldsymbol{\mu}$$

# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$\begin{aligned} E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] &= \text{tr}(\mathbf{I} - \mathbf{P}_X)\sigma^2\mathbf{I} + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_X)\boldsymbol{\mu} \\ &= \sigma^2\text{tr}(\mathbf{I} - \mathbf{P}_X) \end{aligned}$$

# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$\begin{aligned} E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] &= \text{tr}(\mathbf{I} - \mathbf{P}_X)\sigma^2\mathbf{I} + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_X)\boldsymbol{\mu} \\ &= \sigma^2\text{tr}(\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2 r(\mathbf{I} - \mathbf{P}_X) \end{aligned}$$

# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$\begin{aligned}E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] &= \text{tr}(\mathbf{I} - \mathbf{P}_\mathbf{X})\sigma^2\mathbf{I} + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\&= \sigma^2\text{tr}(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \sigma^2r(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \sigma^2(n - r(\mathbf{X}))\end{aligned}$$

# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$\begin{aligned} E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] &= \text{tr}(\mathbf{I} - \mathbf{P}_X)\sigma^2\mathbf{I} + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_X)\boldsymbol{\mu} \\ &= \sigma^2\text{tr}(\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2 r(\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2(n - r(\mathbf{X})) \end{aligned}$$

Therefore an unbiased estimate of  $\sigma^2$  is

$$\frac{\mathbf{e}^T \mathbf{e}}{n - r(\mathbf{X})}$$



# Expectation of $\hat{\sigma}^2$

Use the theorem:

$$\begin{aligned}E[\mathbf{Y}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] &= \text{tr}(\mathbf{I} - \mathbf{P}_\mathbf{X})\sigma^2\mathbf{I} + \boldsymbol{\mu}^T(\mathbf{I} - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\&= \sigma^2\text{tr}(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \sigma^2r(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \sigma^2(n - r(\mathbf{X}))\end{aligned}$$

Therefore an unbiased estimate of  $\sigma^2$  is

$$\frac{\mathbf{e}^T\mathbf{e}}{n - r(\mathbf{X})}$$

## Trace of a Projection Matrix

If  $\mathbf{P}$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(\mathbf{P}) = \sum_i(\lambda_i) = r(\mathbf{P})$

# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$

# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$

# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$

# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)
- a square root of  $\mathbf{A} \geq 0$  is  $\mathbf{U} \mathbf{\Lambda}^{1/2} \mathbf{U}^T$



## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

- $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)
- a square root of  $\mathbf{A} \geq 0$  is  $\mathbf{U} \mathbf{\Lambda}^{1/2} \mathbf{U}^T$

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

- $P = \mathbf{U}\mathbf{U}^T$

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

- $P = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$
- $P = P^2 \Rightarrow \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{U}\mathbf{\Lambda}\mathbf{U}^T = \mathbf{U}\mathbf{\Lambda}^2\mathbf{U}^T$

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

- $P = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$
- $P = P^2 \Rightarrow \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{U}\mathbf{\Lambda}\mathbf{U}^T = \mathbf{U}\mathbf{\Lambda}^2\mathbf{U}^T$
- $\mathbf{\Lambda} = \mathbf{\Lambda}^2$  is true only for  $\lambda_i = 1$  or  $\lambda_i = 0$

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

- $P = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$
- $P = P^2 \Rightarrow \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{U}\mathbf{\Lambda}\mathbf{U}^T = \mathbf{U}\mathbf{\Lambda}^2\mathbf{U}^T$
- $\mathbf{\Lambda} = \mathbf{\Lambda}^2$  is true only for  $\lambda_i = 1$  or  $\lambda_i = 0$
- Since  $r(P)$  is the number of non-zero eigenvalues,  
 $r(P) = \sum \lambda_i = \text{tr}(P)$

# Projections

## Projection Matrix

If  $P$  is an orthogonal projection matrix, then its eigenvalues  $\lambda_i$  are either zero or one with  $\text{tr}(P) = \sum_i (\lambda_i) = r(P)$

- $P = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$
- $P = P^2 \Rightarrow \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{U}\mathbf{\Lambda}\mathbf{U}^T = \mathbf{U}\mathbf{\Lambda}^2\mathbf{U}^T$
- $\mathbf{\Lambda} = \mathbf{\Lambda}^2$  is true only for  $\lambda_i = 1$  or  $\lambda_i = 0$
- Since  $r(P)$  is the number of non-zero eigenvalues,  
 $r(P) = \sum \lambda_i = \text{tr}(P)$

$$P = [\mathbf{U}_P \mathbf{U}_{P^\perp}] \begin{bmatrix} \mathbf{I}_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{n-r} \end{bmatrix} \begin{bmatrix} \mathbf{U}_P^T \\ \mathbf{U}_{P^\perp}^T \end{bmatrix} = \mathbf{U}_P \mathbf{U}_P^T$$

$$P = \sum_{i=1}^r \mathbf{u}_i \mathbf{u}_i^T$$

sum of  $r$  rank 1 projections.

# Prostate Example

```
> library(lasso2)
> summary(lm(lcavol ~ ., data=Prostate))
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	-2.260101	1.259683	-1.794	0.0762	.
lweight	-0.073166	0.174450	-0.419	0.6759	
age	0.022736	0.010964	2.074	0.0410	*
lbph	-0.087449	0.058084	-1.506	0.1358	
svi	-0.153591	0.253932	-0.605	0.5468	
lcp	0.367300	0.081689	4.496	2.10e-05	***
gleason	0.190759	0.154283	1.236	0.2196	
pgg45	-0.007158	0.004326	-1.654	0.1016	
lpsa	0.572797	0.085790	6.677	2.11e-09	***

---

Residual standard error: 0.6998 on 88 degrees of freedom  
Multiple R-squared: 0.6769, Adjusted R-squared: 0.6475  
F-statistic: 23.04 on 8 and 88 DF, p-value: < 2.2e-16