

# Checking Assumptions: Residuals and Influential Observations

Merlise Clyde

STA721 Linear Models

Duke University

October 3, 2013

## Topics

- Distribution of Residuals
- Leverage
- Standardized Residuals
- Cook's Distance
- Example: Stackloss data

Readings: Christensen Chapter 13

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$$

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

What could go wrong?

- Wrong mean

# Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

What could go wrong?

- Wrong mean
- Wrong covariance



# Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

What could go wrong?

- Wrong mean
- Wrong covariance
- Wrong distribution for  $\boldsymbol{\epsilon}$

Residuals (MLE of  $\epsilon$ ) are

$$\mathbf{e} = \mathbf{Y} - \mathbf{X}\hat{\beta}$$

Residuals (MLE of  $\epsilon$ ) are

$$\begin{aligned}\mathbf{e} &= \mathbf{Y} - \mathbf{X}\hat{\beta} \\ &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\mathbf{Y}\end{aligned}$$

Residuals (MLE of  $\epsilon$ ) are

$$\begin{aligned}\mathbf{e} &= \mathbf{Y} - \mathbf{X}\hat{\beta} \\ &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\mathbf{Y}\end{aligned}$$

$$E[\mathbf{e}] = (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\boldsymbol{\mu}$$

Residuals (MLE of  $\epsilon$ ) are

$$\begin{aligned}\mathbf{e} &= \mathbf{Y} - \mathbf{X}\hat{\beta} \\ &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\mathbf{Y}\end{aligned}$$

$$\begin{aligned}\mathbb{E}[\mathbf{e}] &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\ &= \mathbf{0}_n \text{ if } \boldsymbol{\mu} \in C(\mathbf{X})\end{aligned}$$

Residuals (MLE of  $\epsilon$ ) are

$$\begin{aligned}\mathbf{e} &= \mathbf{Y} - \mathbf{X}\hat{\beta} \\ &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\mathbf{Y}\end{aligned}$$

$$\begin{aligned}\mathbb{E}[\mathbf{e}] &= (\mathbf{I}_n - \mathbf{P}_\mathbf{X})\boldsymbol{\mu} \\ &= \mathbf{0}_n \text{ if } \boldsymbol{\mu} \in C(\mathbf{X})\end{aligned}$$

Mean will not be zero if we have left out terms

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\text{Cov} = [\mathbf{e}] = \text{Cov}[(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}]$$



# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_X)^T \text{Cov}[\mathbf{Y}](\mathbf{I} - \mathbf{P}_X)\end{aligned}$$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_X)^T \text{Cov}[\mathbf{Y}](\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_X)^T (\mathbf{I} - \mathbf{P}_X)\end{aligned}$$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_X)\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_X)^T \text{Cov}[\mathbf{Y}](\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_X)^T (\mathbf{I} - \mathbf{P}_X) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_X)\end{aligned}$$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})^T \text{Cov}[\mathbf{Y}] (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})\end{aligned}$$

$$\mathbf{e} \sim \text{N}(\mathbf{0}, \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X}))$$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})^T \text{Cov}[\mathbf{Y}] (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})\end{aligned}$$

$$\mathbf{e} \sim \text{N}(\mathbf{0}, \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X}))$$

- $\mathbf{P}_\mathbf{X}$  is the “hat” matrix (sometimes called  $\mathbf{H} = [h_{ij}]$ )

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})^T \text{Cov}[\mathbf{Y}](\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})\end{aligned}$$

$$\mathbf{e} \sim \text{N}(\mathbf{0}, \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X}))$$

- $\mathbf{P}_\mathbf{X}$  is the “hat” matrix (sometimes called  $\mathbf{H} = [h_{ij}]$ )
- diagonal elements of  $\mathbf{P}_\mathbf{X}$  are denoted as  $h_{ii}$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})^T \text{Cov}[\mathbf{Y}] (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})\end{aligned}$$

$$\mathbf{e} \sim \text{N}(\mathbf{0}, \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X}))$$

- $\mathbf{P}_\mathbf{X}$  is the “hat” matrix (sometimes called  $\mathbf{H} = [h_{ij}]$ )
- diagonal elements of  $\mathbf{P}_\mathbf{X}$  are denoted as  $h_{ii}$
- leverage of the  $i$ th observation is  $h_{ii}$

# Variance Residuals

Under model with constant variances  $\text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$

$$\begin{aligned}\text{Cov} = [\mathbf{e}] &= \text{Cov}[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}] \\ &= (\mathbf{I} - \mathbf{P}_\mathbf{X})^T \text{Cov}[\mathbf{Y}] (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X})\end{aligned}$$

$$\mathbf{e} \sim \text{N}(\mathbf{0}, \sigma^2 (\mathbf{I} - \mathbf{P}_\mathbf{X}))$$

- $\mathbf{P}_\mathbf{X}$  is the “hat” matrix (sometimes called  $\mathbf{H} = [h_{ij}]$ )
- diagonal elements of  $\mathbf{P}_\mathbf{X}$  are denoted as  $h_{ii}$
- leverage of the  $i$ th observation is  $h_{ii}$

$$\text{var}(e_i) = \sigma^2(1 - h_i)$$



# Parameterization

Let  $\mathbf{X} = [\mathbf{1}_n \mathbf{Z}]$  and  $\mathbf{P}_1 = \mathbf{1}(\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T$  denote the projection on  $C(\mathbf{1})$

# Parameterization

Let  $\mathbf{X} = [\mathbf{1}_n \mathbf{Z}]$  and  $\mathbf{P}_1 = \mathbf{1}(\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T$  denote the projection on  $C(\mathbf{1})$

$$\mathbf{Y} = \mathbf{X}\beta + \epsilon$$

# Parameterization

Let  $\mathbf{X} = [\mathbf{1}_n \mathbf{Z}]$  and  $\mathbf{P}_1 = \mathbf{1}(\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T$  denote the projection on  $C(\mathbf{1})$

$$\begin{aligned}\mathbf{Y} &= \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon} \\ &= \mathbf{1}_n \alpha_0 + \mathbf{Z}\boldsymbol{\alpha} + \boldsymbol{\epsilon}\end{aligned}$$

# Parameterization

Let  $\mathbf{X} = [\mathbf{1}_n \mathbf{Z}]$  and  $\mathbf{P}_1 = \mathbf{1}(\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T$  denote the projection on  $C(\mathbf{1})$

$$\begin{aligned}\mathbf{Y} &= \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon} \\ &= \mathbf{1}_n \alpha_0 + \mathbf{Z}\boldsymbol{\alpha} + \boldsymbol{\epsilon} \\ &= \mathbf{1}_n \alpha_0 + \mathbf{P}_1 \mathbf{Z}\boldsymbol{\alpha} + (\mathbf{I}_n - \mathbf{P}_1) \mathbf{Z}\boldsymbol{\alpha} + \boldsymbol{\epsilon}\end{aligned}$$

# Parameterization

Let  $\mathbf{X} = [\mathbf{1}_n \mathbf{Z}]$  and  $\mathbf{P}_1 = \mathbf{1}(\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T$  denote the projection on  $C(\mathbf{1})$

$$\begin{aligned}\mathbf{Y} &= \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon} \\ &= \mathbf{1}_n \alpha_0 + \mathbf{Z}\boldsymbol{\alpha} + \boldsymbol{\epsilon} \\ &= \mathbf{1}_n \alpha_0 + \mathbf{P}_1 \mathbf{Z}\boldsymbol{\alpha} + (\mathbf{I}_n - \mathbf{P}_1) \mathbf{Z}\boldsymbol{\alpha} + \boldsymbol{\epsilon}\end{aligned}$$

$$= \mathbf{1}_n \alpha_0^* + (\mathbf{Z} - \mathbf{1}_n \bar{\mathbf{Z}}^T) \boldsymbol{\alpha} + \boldsymbol{\epsilon}$$

# Mahalanobis Distance & Leverage

Let  $z_i^T$  denote the vector of explanatory variables for the  $i$ th case

# Mahalanobis Distance & Leverage

Let  $z_i^T$  denote the vector of explanatory variables for the  $i$ th case

$$P_{\mathbf{x}} = P_1 + P_{\mathbf{Z}_{-1}} \bar{\mathbf{z}}^T$$

# Mahalanobis Distance & Leverage

Let  $z_i^T$  denote the vector of explanatory variables for the  $i$ th case

$$P_{\mathbf{X}} = P_{\mathbf{1}} + P_{\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T}$$

$$h_{ii} = \frac{1}{n} + (z_i - \bar{\mathbf{Z}})^T \left( (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T)^T (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T) \right)^{-1} (z_i - \bar{\mathbf{Z}})$$



# Mahalanobis Distance & Leverage

Let  $z_i^T$  denote the vector of explanatory variables for the  $i$ th case

$$P_{\mathbf{X}} = P_1 + P_{\mathbf{Z}-1\bar{\mathbf{Z}}^T}$$

$$\begin{aligned} h_{ii} &= \frac{1}{n} + (z_i - \bar{\mathbf{Z}})^T \left( (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T)^T (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T) \right)^{-1} (z_i - \bar{\mathbf{Z}}) \\ &= \frac{1}{n} + \frac{D_i^2}{n-1} \end{aligned}$$

# Mahalanobis Distance & Leverage

Let  $z_i^T$  denote the vector of explanatory variables for the  $i$ th case

$$P_{\mathbf{X}} = P_{\mathbf{1}} + P_{\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T}$$

$$\begin{aligned} h_{ii} &= \frac{1}{n} + (z_i - \bar{\mathbf{Z}})^T \left( (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T)^T (\mathbf{Z} - \mathbf{1}\bar{\mathbf{Z}}^T) \right)^{-1} (z_i - \bar{\mathbf{Z}}) \\ &= \frac{1}{n} + \frac{D_i^2}{n-1} \end{aligned}$$

Leverage is a function of the estimated Mahalanobis distance  $D_i^2$  of  $z_i$  from  $\bar{\mathbf{Z}}$

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent.

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

- $E[\mathbf{e}] = \mathbf{0}_n$ : scatter plots of residuals versus any combination of terms in the mean function will have a constant mean of  $\mathbf{0}$

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

- $E[\mathbf{e}] = \mathbf{0}_n$ : scatter plots of residuals versus any combination of terms in the mean function will have a constant mean of  $\mathbf{0}$
- $\text{var}(e_i) = \sigma^2(1 - h_{ii})$ , so even if fitted model is correct, the variances are not constant. Lower variability for points with high leverage with  $h_{ii} \approx 1$

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

- $E[\mathbf{e}] = \mathbf{0}_n$ : scatter plots of residuals versus any combination of terms in the mean function will have a constant mean of  $\mathbf{0}$
- $\text{var}(e_i) = \sigma^2(1 - h_{ii})$ , so even if fitted model is correct, the variances are not constant. Lower variability for points with high leverage with  $h_{ii} \approx 1$
- The residuals are correlated, but this correlation is usually not visible in residual plots

# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

- $E[\mathbf{e}] = \mathbf{0}_n$ : scatter plots of residuals versus any combination of terms in the mean function will have a constant mean of  $\mathbf{0}$
- $\text{var}(e_i) = \sigma^2(1 - h_{ii})$ , so even if fitted model is correct, the variances are not constant. Lower variability for points with high leverage with  $h_{ii} \approx 1$
- The residuals are correlated, but this correlation is usually not visible in residual plots

When the model is not correct we hope to see structure in the residual plots to help us re-model the data



# Residual Plots

Under normality and constant variance assumptions, the residuals  $\mathbf{e}$  and fitted values  $\hat{\mathbf{Y}}$  are jointly normal but independent. Under the correct model

- $E[\mathbf{e}] = \mathbf{0}_n$ : scatter plots of residuals versus any combination of terms in the mean function will have a constant mean of  $\mathbf{0}$
- $\text{var}(e_i) = \sigma^2(1 - h_{ii})$ , so even if fitted model is correct, the variances are not constant. Lower variability for points with high leverage with  $h_{ii} \approx 1$
- The residuals are correlated, but this correlation is usually not visible in residual plots

When the model is not correct we hope to see structure in the residual plots to help us re-model the data

Use standardized residuals

# Standardized Residuals

Since  $\text{var}(e_i) = \sigma^2(1 - h_{ii})$  has non-constant variance

# Standardized Residuals

Since  $\text{var}(e_i) = \sigma^2(1 - h_{ii})$  has non-constant variance

- Divide by standard deviation:

$$\frac{e_i}{\sigma\sqrt{1 - h_{ii}}}$$

(constant variance)

# Standardized Residuals

Since  $\text{var}(e_i) = \sigma^2(1 - h_{ii})$  has non-constant variance

- Divide by standard deviation:

$$\frac{e_i}{\sigma\sqrt{1 - h_{ii}}}$$

(constant variance)

- (Internally) Standardized residuals

$$r_i = \frac{e_i}{\hat{\sigma}\sqrt{1 - h_{ii}}}$$

where  $\hat{\sigma}$  is root MSE

# Standardized Residuals

Since  $\text{var}(e_i) = \sigma^2(1 - h_{ii})$  has non-constant variance

- Divide by standard deviation:

$$\frac{e_i}{\sigma\sqrt{1 - h_{ii}}}$$

(constant variance)

- (Internally) Standardized residuals

$$r_i = \frac{e_i}{\hat{\sigma}\sqrt{1 - h_{ii}}}$$

where  $\hat{\sigma}$  is root MSE

# Properties of Leverage

- $\sum_i h_{ij} = 1$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\mathbf{1}^T \mathbf{P}_X = \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T\end{aligned}$$



# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}_{-1} \bar{\mathbf{Z}}^T} \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-\mathbf{1}\bar{\mathbf{z}}^T} \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ij} \leq 1$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-\mathbf{1}\bar{\mathbf{z}}^T} \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ij} \leq 1$

$$\mathbf{P}_X = \mathbf{P}_X^2$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ii} \leq 1$

$$\begin{aligned}\mathbf{P}_X &= \mathbf{P}_X^2 \\ h_{ii} &= \sum_j h_{ij} h_{ji}\end{aligned}$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_{\mathbf{X}} &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ii} \leq 1$

$$\begin{aligned}\mathbf{P}_{\mathbf{X}} &= \mathbf{P}_{\mathbf{X}}^2 \\ h_{ii} &= \sum_j h_{ij} h_{ji} \\ &= \sum_j h_{jj}^2\end{aligned}$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ii} \leq 1$

$$\begin{aligned}\mathbf{P}_X &= \mathbf{P}_X^2 \\ h_{ii} &= \sum_j h_{ij} h_{ji} \\ &= \sum_j h_{jj}^2 \\ &= h_{ii}^2 + \sum_{j \neq i} h_{jj}^2\end{aligned}$$

# Properties of Leverage

- $\sum_i h_{ij} = 1$

$$\begin{aligned}\mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T \mathbf{P}_1 + \mathbf{1}^T \mathbf{P}_{\mathbf{Z}-1} \bar{\mathbf{z}}^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T\end{aligned}$$

- $0 \leq h_{ii} \leq 1$

$$\begin{aligned}\mathbf{P}_X &= \mathbf{P}_X^2 \\ h_{ii} &= \sum_j h_{ij} h_{ji} \\ &= \sum_j h_{jj}^2 \\ &= h_{ii}^2 + \sum_{j \neq i} h_{jj}^2\end{aligned}$$

$$h_{ii}(1 - h_{ii}) \geq 0$$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$



$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\mathbf{1}^T \mathbf{P}_{\mathbf{X}} = \mathbf{1}^T (\mathbf{P}_1 + \mathbf{P}_{\mathbf{X}}) (\mathbf{P}_1 + \mathbf{P}_{\mathbf{X}})^T$$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\begin{aligned} \mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T (\mathbf{P}_1 + \mathbf{P}_X) (\mathbf{P}_1 + \mathbf{P}_X)^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \end{aligned}$$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\begin{aligned} \mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T (\mathbf{P}_1 + \mathbf{P}_X) (\mathbf{P}_1 + \mathbf{P}_X)^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T \end{aligned}$$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\begin{aligned} \mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T (\mathbf{P}_1 + \mathbf{P}_X) (\mathbf{P}_1 + \mathbf{P}_X)^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T \end{aligned}$$

- $h_{ij} \approx 0$  for  $i \neq j$  if  $h_{ii} \approx 1$

$$\hat{Y}_i = h_{ii} Y_i + \sum_{j \neq i} h_{ij} Y_j$$

If  $h_{ii} \approx 1$  then  $\hat{Y}_i \approx Y_i$

- $\sum_j h_{ij} = 1$
- $\sum_i h_{ij}^2 = 1$

$$\begin{aligned} \mathbf{1}^T \mathbf{P}_X &= \mathbf{1}^T (\mathbf{P}_1 + \mathbf{P}_X) (\mathbf{P}_1 + \mathbf{P}_X)^T \\ &= \mathbf{1}^T \mathbf{1} (\mathbf{1}^T \mathbf{1})^{-1} \mathbf{1}^T + \mathbf{0}^T \\ &= \mathbf{1}^T \end{aligned}$$

- $h_{ij} \approx 0$  for  $i \neq j$  if  $h_{ii} \approx 1$
- $\text{var}(e_i) = \sigma^2(1 - h_{ii}) \approx 0$  if  $h_{ii} \approx 1$



# Illustration

# Predicted Residuals

Let  $\mathbf{X}_{(i)}$  and  $\mathbf{Y}_{(i)}$  denote the design matrix and response vector with the  $i$ th row  $\mathbf{x}_i^T$  deleted

Let  $\mathbf{X}_{(i)}$  and  $\mathbf{Y}_{(i)}$  denote the design matrix and response vector with the  $i$ th row  $\mathbf{x}_i^T$  deleted

- $\hat{\beta}_{(i)} = (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)}$

Let  $\mathbf{X}_{(i)}$  and  $\mathbf{Y}_{(i)}$  denote the design matrix and response vector with the  $i$ th row  $\mathbf{x}_i^T$  deleted

- $\hat{\beta}_{(i)} = (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)}$
- Predicted residual  $\mathbf{e}_{(i)} = y_i - \mathbf{x}_{(i)}^T \hat{\beta}_{(i)}$

# Predicted Residuals

Let  $\mathbf{X}_{(i)}$  and  $\mathbf{Y}_{(i)}$  denote the design matrix and response vector with the  $i$ th row  $\mathbf{x}_i^T$  deleted

- $\hat{\beta}_{(i)} = (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)}$
- Predicted residual  $\mathbf{e}_{(i)} = y_i - \mathbf{x}_{(i)}^T \hat{\beta}_{(i)}$
- Variance  $\text{var}(e_{(i)}) = \sigma^2(1 + \mathbf{x}_{(i)}^T (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{x}_{(i)})$

# Predicted Residuals

Let  $\mathbf{X}_{(i)}$  and  $\mathbf{Y}_{(i)}$  denote the design matrix and response vector with the  $i$ th row  $\mathbf{x}_i^T$  deleted

- $\hat{\beta}_{(i)} = (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)}$
- Predicted residual  $\mathbf{e}_{(i)} = y_i - \mathbf{x}_{(i)}^T \hat{\beta}_{(i)}$
- Variance  $\text{var}(e_{(i)}) = \sigma^2(1 + \mathbf{x}_{(i)}^T (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{x}_{(i)})$

# Updating Formula

How to compute without re-fitting model for each case deletion?

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{x}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{x}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^t (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$



# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{x}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{x}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^t (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{x}_{(i)}^T \mathbf{x}_{(i)} = \mathbf{x}^T \mathbf{x} + \frac{(\mathbf{x}^T \mathbf{x})^{-1} \mathbf{x}_i \mathbf{x}_i^t (\mathbf{x}^T \mathbf{x})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{x} \mathbf{x})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

$$\hat{\beta}_{(i)} = (\mathbf{x}_{(i)}^T \mathbf{x}_{(i)})^{-1} \mathbf{x}_{(i)}^T \mathbf{y}_{(i)}$$

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{X}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{X}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^t (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} (\mathbf{X}^T \mathbf{Y} - \mathbf{x}_i y_i)\end{aligned}$$

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{X}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{X}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} (\mathbf{X}^T \mathbf{Y} - \mathbf{x}_i y_i) \\ &\quad \text{some algebra}\end{aligned}$$

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{X}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{X}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} (\mathbf{X}^T \mathbf{Y} - \mathbf{x}_i y_i) \\ &\quad \text{some algebra} \\ &= \hat{\beta} + \frac{\mathbf{X}^T \mathbf{X} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

# Updating Formula

How to compute without re-fitting model for each case deletion?

$$\mathbf{X}_{(i)}^T \mathbf{X}_{(i)} = \mathbf{X}^T \mathbf{X} + \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1}}{1 - \mathbf{x}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i}$$

Special case of the Binomial Inverse Theorem or Woodbury Theorem

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} (\mathbf{X}^T \mathbf{Y} - \mathbf{x}_i y_i) \\ &\quad \text{some algebra} \\ &= \hat{\beta} + \frac{\mathbf{X}^T \mathbf{X} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

# Predicted Residuals

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}(i) = \frac{e_i}{1 - h_{ii}}$$

# Predicted Residuals

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}(i) = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$



# Predicted Residuals

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}(i) = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$

Standardized predicted residual is

$$\frac{e_{(i)}}{\sqrt{\text{var}(e_{(i)})}} = \frac{e_i / (1 - h_{ii})}{\sigma / \sqrt{1 - h_{ii}}} = \frac{e_i}{\sigma \sqrt{1 - h_{ii}}}$$

# Predicted Residuals

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}(i) = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$

Standardized predicted residual is

$$\frac{e_{(i)}}{\sqrt{\text{var}(e_{(i)})}} = \frac{e_i / (1 - h_{ii})}{\sigma / \sqrt{1 - h_{ii}}} = \frac{e_i}{\sigma \sqrt{1 - h_{ii}}}$$

Same as before!

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

$$\text{SSE}_{(i)} = \text{SSE} - \frac{e_i^2}{1 - h_{ii}}$$

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

Externally Standardized residuals

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

$$\begin{aligned}\text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1}\end{aligned}$$

Externally Standardized residuals

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}}$$

# External estimate of $\sigma^2$

Estimate  $\hat{\sigma}_{(i)}^2$  using data with case  $i$  deleted

$$\begin{aligned}\text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1}\end{aligned}$$

Externally Standardized residuals

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}}$$

May still miss extreme points with high leverage, but will pick up unusual  $y_i$ s



# Outlier Test

$$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} \text{ versus } H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$$

$$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} \text{ versus } H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$$

- Show that t-test for testing  $H_0: \alpha_i = 0$  is equal to  $t_i$  (HW)

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$  versus  $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing  $H_0: \alpha_i = 0$  is equal to  $t_i$  (HW)
- if p-value is small declare the  $i$ th case to be an outlier:  $E[Y_i]$  not given by  $\mathbf{X}\boldsymbol{\beta}$  but  $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$  versus  $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing  $H_0: \alpha_i = 0$  is equal to  $t_i$  (HW)
- if p-value is small declare the  $i$ th case to be an outlier:  $E[Y_i]$  not given by  $\mathbf{X}\boldsymbol{\beta}$  but  $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- control for multiple testing

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$  versus  $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing  $H_0: \alpha_i = 0$  is equal to  $t_i$  (HW)
- if p-value is small declare the  $i$ th case to be an outlier:  $E[Y_i]$  not given by  $\mathbf{X}\boldsymbol{\beta}$  but  $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- control for multiple testing
- Extreme case  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} + \mathbf{I}_n\boldsymbol{\alpha}$  all points have their own mean!

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$  versus  $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing  $H_0: \alpha_i = 0$  is equal to  $t_i$  (HW)
- if p-value is small declare the  $i$ th case to be an outlier:  $E[Y_i]$  not given by  $\mathbf{X}\boldsymbol{\beta}$  but  $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- control for multiple testing
- Extreme case  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} + \mathbf{I}_n\boldsymbol{\alpha}$  all points have their own mean!

Cook's Distance measure of how much predictions change with  $i$ th case deleted

# Cook's Distance

Cook's Distance measure of how much predictions change with  $i$ th case deleted

$$D_i = \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2}$$



# Cook's Distance

Cook's Distance measure of how much predictions change with  $i$ th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

# Cook's Distance

Cook's Distance measure of how much predictions change with  $i$ th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

Flag cases where  $D_i > 1$  or large relative to other cases

# Cook's Distance


Cook's Distance measure of how much predictions change with  $i$ th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

Flag cases where  $D_i > 1$  or large relative to other cases Influential




Cases

# Stackloss Data







# Stackloss Added Variable Plot

Added-Variable Plots



# Stackloss Data



- Leverage 0.285
- Cooks'd Distance .69
- p-value  $t_{21}$  is 0.0042
- Bonferonni adjusted p-value is 0.0024
- Other points? Masking?
- Refit without Case 21

Others have suggested that cases (1, 3, 4, 21) are outliers

# To Remove or Not?

- For suspicious cases, check data sources



# To Remove or Not?

- For suspicious cases, check data sources
- Check that points are not outliers because of wrong mean function or distributional assumptions

# To Remove or Not?

- For suspicious cases, check data sources
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Influential cases - report results with and without cases (results may change)

# To Remove or Not?

- For suspicious cases, check data sources
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Influential cases - report results with and without cases (results may change)
- Outlier test - suggests alternative population; if not influential may in keep analysis, but will inflate  $\hat{\sigma}^2$  and interval estimates

# To Remove or Not?

- For suspicious cases, check data sources
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Influential cases - report results with and without cases (results may change)
- Outlier test - suggests alternative population; if not influential may in keep analysis, but will inflate  $\hat{\sigma}^2$  and interval estimates
- Document steps - reproducibility!
- Robust Regression Methods

# To Remove or Not?

- For suspicious cases, check data sources
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Influential cases - report results with and without cases (results may change)
- Outlier test - suggests alternative population; if not influential may in keep analysis, but will inflate  $\hat{\sigma}^2$  and interval estimates
- Document steps - reproducibility!
- Robust Regression Methods

Are there outliers in the Stackloss Data?