

Checking Assumptions

Merlise Clyde

STA721 Linear Models

Duke University

October 8, 2012

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\boldsymbol{\mu} \in C(\mathbf{X}) \Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$$

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Focus on

- Wrong mean for a case or cases

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Focus on

- Wrong mean for a case or cases
- Wrong distribution for $\boldsymbol{\epsilon}$

Linear Model

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Focus on

- Wrong mean for a case or cases
- Wrong distribution for $\boldsymbol{\epsilon}$
- Cases that influence the mean

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

Focus on

- Wrong mean for a case or cases
- Wrong distribution for $\boldsymbol{\epsilon}$
- Cases that influence the mean

If $\mu_i \neq \mathbf{x}_i^T \boldsymbol{\beta}$ then expected value of $e_i = Y_i - \hat{Y}_i$ is not zero;

Standardized residuals

standardized residuals

$$r_i = e_i / \sqrt{\sigma^2(1 - h_{ii})}$$

- Under correct model have mean 0 and scale 1

standardized residuals

$$r_i = e_i / \sqrt{\sigma^2(1 - h_{ii})}$$

- Under correct model have mean 0 and scale 1
- Use estimate of σ^2

standardized residuals

$$r_i = e_i / \sqrt{\sigma^2(1 - h_{ii})}$$

- Under correct model have mean 0 and scale 1
- Use estimate of σ^2
- The distribution of r_i is not a t

standardized residuals

$$r_i = e_i / \sqrt{\sigma^2(1 - h_{ii})}$$

- Under correct model have mean 0 and scale 1
- Use estimate of σ^2
- The distribution of r_i is not a t
- if h_{ii} (leverage) is close to 1, then \hat{Y}_i is close to Y_i so e_i is approximately 0

standardized residuals

$$r_i = e_i / \sqrt{\sigma^2(1 - h_{ii})}$$

- Under correct model have mean 0 and scale 1
- Use estimate of σ^2
- The distribution of r_i is not a t
- if h_{ii} (leverage) is close to 1, then \hat{Y}_i is close to Y_i so e_i is approximately 0
- Variance is also almost 0, so standardize value may not flag “outliers”

Predicted Residuals

Estimates without Case (i):

$$\hat{\beta}_{(i)} = (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)}$$

Predicted Residuals

Estimates without Case (i):

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= \hat{\beta} - \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

Predicted Residuals

Estimates without Case (i):

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= \hat{\beta} - \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\beta}_{(i)} = \frac{e_i}{1 - h_{ii}}$$

Predicted Residuals

Estimates without Case (i):

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= \hat{\beta} - \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\beta}_{(i)} = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$

Predicted Residuals

Estimates without Case (i):

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= \hat{\beta} - \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\beta}_{(i)} = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$

Standardized predicted residual is

$$\frac{e_{(i)}}{\sqrt{\text{var}(e_{(i)})}} = \frac{e_i / (1 - h_{ii})}{\sigma / \sqrt{1 - h_{ii}}} = \frac{e_i}{\sigma \sqrt{1 - h_{ii}}}$$

Predicted Residuals

Estimates without Case (i):

$$\begin{aligned}\hat{\beta}_{(i)} &= (\mathbf{X}_{(i)}^T \mathbf{X}_{(i)})^{-1} \mathbf{X}_{(i)}^T \mathbf{Y}_{(i)} \\ &= \hat{\beta} - \frac{(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_i e_i}{1 - h_{ii}}\end{aligned}$$

Predicted residual

$$e_{(i)} = y_i - \mathbf{x}_i^T \hat{\beta}_{(i)} = \frac{e_i}{1 - h_{ii}}$$

with variance

$$\text{var}(e_{(i)}) = \frac{\sigma^2}{1 - h_{ii}}$$

Standardized predicted residual is

$$\frac{e_{(i)}}{\sqrt{\text{var}(e_{(i)})}} = \frac{e_i / (1 - h_{ii})}{\sigma / \sqrt{1 - h_{ii}}} = \frac{e_i}{\sigma \sqrt{1 - h_{ii}}}$$

Can show that these are the same as standardized residual!

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\text{SSE}_{(i)} = \text{SSE} - \frac{e_i^2}{1 - h_{ii}}$$

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

Externally Standardized residuals

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\begin{aligned}\text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1}\end{aligned}$$

Externally Standardized residuals

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}}$$

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

Externally Standardized residuals

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = r_i \left(\frac{n - p - 1}{n - p - r_i^2} \right)^{1/2}$$

External estimate of σ^2

Estimate $\hat{\sigma}_{(i)}^2$ using data with case i deleted

$$\begin{aligned} \text{SSE}_{(i)} &= \text{SSE} - \frac{e_i^2}{1 - h_{ii}} \\ \hat{\sigma}_{(i)}^2 = \text{MSE}_{(i)} &= \frac{\text{SSE}_{(i)}}{n - p - 1} \end{aligned}$$

Externally Standardized residuals

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = r_i \left(\frac{n - p - 1}{n - p - r_i^2} \right)^{1/2}$$

May still miss extreme points with high leverage, but will pick up unusual y_i s

Distribution of Externally Studentized Residual

$$t_i = \frac{e_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} = \frac{y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_{(i)}}{\sqrt{\hat{\sigma}_{(i)}^2/(1 - h_{ii})}} \sim \text{St}(n - p - 1)$$

Outlier Test

$$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} \text{ versus } H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$$

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i

Outlier Test

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i
- if p-value is small declare the i th case to be an outlier: $E[Y_i]$ not given by $\mathbf{X}\boldsymbol{\beta}$ but $\mathbf{X}\boldsymbol{\beta} + \delta_i \alpha_i$

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i
- if p-value is small declare the i th case to be an outlier: $E[Y_i]$ not given by $\mathbf{X}\boldsymbol{\beta}$ but $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- Can extend to include multiple δ_i and δ_j to test that case i and j are both outliers

Outlier Test

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i
- if p-value is small declare the i th case to be an outlier: $E[Y_i]$ not given by $\mathbf{X}\boldsymbol{\beta}$ but $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- Can extend to include multiple δ_i and δ_j to test that case i and j are both outliers
- Extreme case $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} + \mathbf{I}_n\boldsymbol{\alpha}$ all points have their own mean!

Outlier Test

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i
- if p-value is small declare the i th case to be an outlier: $E[Y_i]$ not given by $\mathbf{X}\boldsymbol{\beta}$ but $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- Can extend to include multiple δ_i and δ_j to test that case i and j are both outliers
- Extreme case $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} + \mathbf{I}_n\boldsymbol{\alpha}$ all points have their own mean!
- Control for multiple testing

Outlier Test

$H_0: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$ versus $H_a: \mu_i = \mathbf{x}_i^T \boldsymbol{\beta} + \alpha_i$

- Show that t-test for testing $H_0: \alpha_i = 0$ is equal to t_i
- if p-value is small declare the i th case to be an outlier: $E[Y_i]$ not given by $\mathbf{X}\boldsymbol{\beta}$ but $\mathbf{X}\boldsymbol{\beta} + \delta_i\alpha_i$
- Can extend to include multiple δ_i and δ_j to test that case i and j are both outliers
- Extreme case $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} + \mathbf{I}_n\boldsymbol{\alpha}$ all points have their own mean!
- Control for multiple testing

look to see if the $\max |t_i|$ is larger than expected under the null

Multiple Testing

look to see if the $\max |t_i|$ is larger than expected under the null

- Need distribution of the max of Student t random variables (simulate)

Multiple Testing

look to see if the $\max |t_i|$ is larger than expected under the null

- Need distribution of the max of Student t random variables (simulate)
- Conservative is Bonferroni Correction: For n tests of size α the probability of falsely labeling at least one case as an outlier is no greater than $n\alpha$; e.g. with 21 cases and $\alpha = 0.05$, the probability is no greater than 1.05!

Multiple Testing

look to see if the $\max |t_i|$ is larger than expected under the null

- Need distribution of the max of Student t random variables (simulate)
- Conservative is Bonferroni Correction: For n tests of size α the probability of falsely labeling at least one case as an outlier is no greater than $n\alpha$; e.g. with 21 cases and $\alpha = 0.05$, the probability is no greater than 1.05!
- Adjust $\alpha^* = \alpha/n$ so that the probability of falsely labeling at least one point an outlier is α $\alpha/n = .00238$ with $n = 21$

Multiple Testing

look to see if the $\max |t_i|$ is larger than expected under the null

- Need distribution of the max of Student t random variables (simulate)
- Conservative is Bonferroni Correction: For n tests of size α the probability of falsely labeling at least one case as an outlier is no greater than $n\alpha$; e.g. with 21 cases and $\alpha = 0.05$, the probability is no greater than 1.05!
- Adjust $\alpha^* = \alpha/n$ so that the probability of falsely labeling at least one point an outlier is α $\alpha/n = .00238$ with $n = 21$

Cook's Distance measure of how much predictions change with i th case deleted

Cook's Distance

Cook's Distance measure of how much predictions change with i th case deleted

$$D_i = \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2}$$

Cook's Distance

Cook's Distance measure of how much predictions change with i th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

Cook's Distance

Cook's Distance measure of how much predictions change with i th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

Flag cases where $D_i > 1$ or large relative to other cases

Cook's Distance


Cook's Distance measure of how much predictions change with i th case deleted

$$\begin{aligned} D_i &= \frac{\|\mathbf{Y}_{(i)} - \mathbf{Y}\|^2}{p\hat{\sigma}^2} = \frac{(\hat{\beta}_{(i)} - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}_{(i)} - \hat{\beta})}{p\hat{\sigma}^2} \\ &= \frac{r_i^2}{p} \frac{h_{ii}}{1 - h_{ii}} \end{aligned}$$

Flag cases where $D_i > 1$ or large relative to other cases




Influential Cases

Stackloss Data







Stackloss Added Variable Plot

Added-Variable Plots



Stackloss Data Again



- Leverage 0.285 (compare to $p/n = .19$)

Case 21

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042
- Bonferroni adjusted p-value is 0.0024

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042
- Bonferroni adjusted p-value is 0.0024
- Cooks' Distance .69

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042
- Bonferroni adjusted p-value is 0.0024
- Cooks' Distance .69
- Other points? Masking?

Case 21

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042
- Bonferroni adjusted p-value is 0.0024
- Cooks' Distance .69
- Other points? Masking?
- Refit without Case 21

- Leverage 0.285 (compare to $p/n = .19$)
- p-value t_{21} is 0.0042
- Bonferroni adjusted p-value is 0.0024
- Cooks' Distance .69
- Other points? Masking?
- Refit without Case 21

Other analyses have suggested that cases (1, 2, 3, 4, 21) are outliers

Multiple Outliers

- Hoeting, Madigan and Raftery (in various permutations) consider the problem of simultaneous variable selection and outlier identification.

Multiple Outliers

- Hoeting, Madigan and Raftery (in various permutations) consider the problem of simultaneous variable selection and outlier identification.
- This is implemented in the library(BMA) in the function MC3.REG. This has the advantage that more than 2 points may be considered as outliers at the same time.

Multiple Outliers

- Hoeting, Madigan and Raftery (in various permutations) consider the problem of simultaneous variable selection and outlier identification.
- This is implemented in the library(BMA) in the function MC3.REG. This has the advantage that more than 2 points may be considered as outliers at the same time.
- The function uses a Markov chain to identify both important variables and potential outliers, but is coded in Fortran so should run reasonably quickly.

Multiple Outliers

- Hoeting, Madigan and Raftery (in various permutations) consider the problem of simultaneous variable selection and outlier identification.
- This is implemented in the library(BMA) in the function MC3.REG. This has the advantage that more than 2 points may be considered as outliers at the same time.
- The function uses a Markov chain to identify both important variables and potential outliers, but is coded in Fortran so should run reasonably quickly.
- Can also use BAS or other variable selection programs

Multiple Outliers





- Hoeting, Madigan and Raftery (in various permutations) consider the problem of simultaneous variable selection and outlier identification.
- This is implemented in the library(BMA) in the function MC3.REG. This has the advantage that more than 2 points may be considered as outliers at the same time.
- The function uses a Markov chain to identify both important variables and potential outliers, but is coded in Fortran so should run reasonably quickly.
- Can also use BAS or other variable selection programs


```
library(MASS)
data(stackloss)
n = nrow(stackloss)
stack.out = cbind(stackloss, diag(n))
```

```
library(BAS)
BAS.stack = bas.lm(stack.loss ~ ., data=stack.out,
                    prior="hyper-g-n", a=3,
                    modelprior=beta.binomial(1, ncol(stack.out)),
                    n.models=2^20,
                    method="MCMC", MCMC.it=2^21)
```


Output

```
> summary(BAS.stack)
      Intercept Air.Flow Water.Temp Acid.Conc. '1' '2' '3' '4' '5' '6' '7' '8'
[1,]          1          1          0          0  0  0  0  0  1  0  0  0  0
[2,]          1          1          0          0  0  0  0  0  0  0  0  0  0
[3,]          1          1          0          0  0  0  0  0  0  0  0  0  0
[4,]          1          1          1          0  0  0  0  0  0  0  0  0  0
[5,]          1          1          1          0  0  0  0  0  0  0  0  0  0
      '9' '10' '11' '12' '13' '14' '15' '16' '17' '18' '19' '20' '21'
[1,]  0  0  0  0  0  0  0  0  0  0  0  0  0  1
[2,]  0  0  0  0  0  0  0  0  0  0  0  0  0  1
[3,]  0  0  0  0  0  0  0  0  0  0  0  0  0  0
[4,]  0  0  0  0  0  0  0  0  0  0  0  0  0  0
[5,]  0  0  0  0  0  0  0  0  0  0  0  0  0  1
      BF PostProbs      R2 dim logmarg
[1,] 1.00000000000 0.2940 0.9605  4 22.16893
[2,] 0.0644223528  0.2904 0.9271  3 19.42664
[3,] 0.0008704905  0.0922 0.8458  2 15.12248
[4,] 0.0092248619  0.0416 0.9088  3 17.48308
[5,] 0.0845728129  0.0249 0.9466  4 19.69879
>
```





Renormalized likelihood Estimates vs MCMC frequencies



To Remove or Not?

- For suspicious cases, check data sources for errors

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)
- Outlier test - suggests alternative population for the case(s); if not influential may in keep analysis, but will inflate $\hat{\sigma}^2$ and interval estimates

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)
- Outlier test - suggests alternative population for the case(s); if not influential may in keep analysis, but will inflate $\hat{\sigma}^2$ and interval estimates
- Document how you handle any case deletions - reproducibility!

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)
- Outlier test - suggests alternative population for the case(s); if not influential may in keep analysis, but will inflate $\hat{\sigma}^2$ and interval estimates
- Document how you handle any case deletions - reproducibility!
- Robust Regression Methods

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)
- Outlier test - suggests alternative population for the case(s); if not influential may in keep analysis, but will inflate $\hat{\sigma}^2$ and interval estimates
- Document how you handle any case deletions - reproducibility!
- Robust Regression Methods