

# Models & Estimation

Merlise Clyde

STA721 Linear Models

Duke University

August 28, 2014

Readings: Christensen Chapter 1-2, Appendix A

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed)

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0}$

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)



Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)
- $\text{Cov}[\boldsymbol{\epsilon}] = \sigma^2 \mathbf{I}_n$

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)
- $\text{Cov}[\boldsymbol{\epsilon}] = \sigma^2 \mathbf{I}_n \Rightarrow \text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$  (errors are uncorrelated)

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)
- $\text{Cov}[\boldsymbol{\epsilon}] = \sigma^2 \mathbf{I}_n \Rightarrow \text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$  (errors are uncorrelated)
- $\boldsymbol{\epsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I})$

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)
- $\text{Cov}[\boldsymbol{\epsilon}] = \sigma^2 \mathbf{I}_n \Rightarrow \text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$  (errors are uncorrelated)
- $\boldsymbol{\epsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I}) \Rightarrow \mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I})$

Take an random vector  $\mathbf{Y} \in \mathbb{R}^n$  which is observable and decompose

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

into  $\boldsymbol{\mu} \in \mathbb{R}^n$  (unknown, fixed) and  $\boldsymbol{\epsilon} \in \mathbb{R}^n$  unobservable error vector (random)

Usual assumptions?

- $E[\boldsymbol{\epsilon}] = \mathbf{0} \Rightarrow E[\mathbf{Y}] = \boldsymbol{\mu}$  (mean vector)
- $\text{Cov}[\boldsymbol{\epsilon}] = \sigma^2 \mathbf{I}_n \Rightarrow \text{Cov}[\mathbf{Y}] = \sigma^2 \mathbf{I}_n$  (errors are uncorrelated)
- $\boldsymbol{\epsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I}) \Rightarrow \mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I})$

The distribution assumption allows us to write down a likelihood function

# Likelihood Functions

The likelihood function for  $\mu, \sigma^2$  is proportional to the sampling distribution of the data

# Likelihood Functions

The likelihood function for  $\boldsymbol{\mu}, \sigma^2$  is proportional to the sampling distribution of the data

$$\mathcal{L}(\boldsymbol{\mu}, \sigma^2) \propto (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{(\mathbf{Y} - \boldsymbol{\mu})^T (\mathbf{Y} - \boldsymbol{\mu})}{\sigma^2} \right\}$$

# Likelihood Functions

The likelihood function for  $\boldsymbol{\mu}, \sigma^2$  is proportional to the sampling distribution of the data

$$\begin{aligned}\mathcal{L}(\boldsymbol{\mu}, \sigma^2) &\propto (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{(\mathbf{Y} - \boldsymbol{\mu})^T (\mathbf{Y} - \boldsymbol{\mu})}{\sigma^2} \right\} \\ &\propto (\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right\}\end{aligned}$$



# Likelihood Functions

The likelihood function for  $\boldsymbol{\mu}, \sigma^2$  is proportional to the sampling distribution of the data

$$\begin{aligned}\mathcal{L}(\boldsymbol{\mu}, \sigma^2) &\propto (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{(\mathbf{Y} - \boldsymbol{\mu})^T (\mathbf{Y} - \boldsymbol{\mu})}{\sigma^2} \right\} \\ &\propto (\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right\}\end{aligned}$$

Find values of  $\hat{\boldsymbol{\mu}}$  and  $\hat{\sigma}^2$  that maximize the likelihood  $\mathcal{L}(\boldsymbol{\mu}, \sigma^2)$  for  $\boldsymbol{\mu} \in \mathbb{R}^n$  and  $\sigma^2 \in \mathbb{R}^+$

# Likelihood Functions

The likelihood function for  $\boldsymbol{\mu}, \sigma^2$  is proportional to the sampling distribution of the data

$$\begin{aligned}\mathcal{L}(\boldsymbol{\mu}, \sigma^2) &\propto (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{(\mathbf{Y} - \boldsymbol{\mu})^T (\mathbf{Y} - \boldsymbol{\mu})}{\sigma^2} \right\} \\ &\propto (\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2} \right\}\end{aligned}$$

Find values of  $\hat{\boldsymbol{\mu}}$  and  $\hat{\sigma}^2$  that maximize the likelihood  $\mathcal{L}(\boldsymbol{\mu}, \sigma^2)$  for  $\boldsymbol{\mu} \in \mathbb{R}^n$  and  $\sigma^2 \in \mathbb{R}^+$

Clearly,  $\hat{\boldsymbol{\mu}} = \mathbf{Y}$  but  $\hat{\sigma}^2 = 0$  is outside the parameter space

# Restrictions on $\mu$

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$   
restrict the possible values for  $\mu$

# Restrictions on $\mu$

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$   
restrict the possible values for  $\mu$

- Oneway Anova Model  $j = 1, \dots, J$  groups with  $n_j$  replicates in each group  $j$

# Restrictions on $\mu$

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$   
restrict the possible values for  $\mu$

- Oneway Anova Model  $j = 1, \dots, J$  groups with  $n_j$  replicates in each group  $j$
- “cell means”  $Y_{ij} = \mu_j + \epsilon_{ij}$

$$\mu = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{0}_{n_1} & \dots & \mathbf{0}_{n_1} \\ \mathbf{0}_{n_2} & \mathbf{1}_{n_2} & \dots & \mathbf{0}_{n_2} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}_{n_J} & \dots & \mathbf{0}_{n_J} & \mathbf{1}_{n_J} \end{bmatrix} \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_J \end{pmatrix}$$

- $\mathbf{1}_{n_j}$  is a vector of length  $n_j$  of ones.
- $\mathbf{0}_{n_j}$  is a vector of length  $n_j$  of zeros.

# Equivalent Model

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$

# Equivalent Model

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$

- Oneway Anova Model  $j = 1, \dots, J$  groups with  $n_j$  replicates in each group  $j$

# Equivalent Model

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$

- Oneway Anova Model  $j = 1, \dots, J$  groups with  $n_j$  replicates in each group  $j$
- “Treatment effects”  $Y_{ij} = \mu + \tau_j + \epsilon_{ij}$

$$\mu = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{1}_{n_1} & \mathbf{0}_{n_1} & \dots & \mathbf{0}_{n_1} \\ \mathbf{1}_{n_2} & \mathbf{0}_{n_2} & \mathbf{1}_{n_2} & \dots & \mathbf{0}_{n_2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{1}_{n_J} & \mathbf{0}_{n_J} & \dots & \mathbf{0}_{n_J} & \mathbf{1}_{n_J} \end{bmatrix} \begin{pmatrix} \mu \\ \tau_1 \\ \vdots \\ \tau_J \end{pmatrix}$$



# Equivalent Model

Linear model  $\mu = \mathbf{X}\beta$  for fixed  $\mathbf{X}$  ( $n \times p$ ) and some  $\beta \in \mathbb{R}^p$

- Oneway Anova Model  $j = 1, \dots, J$  groups with  $n_j$  replicates in each group  $j$
- “Treatment effects”  $Y_{ij} = \mu + \tau_j + \epsilon_{ij}$

$$\mu = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{1}_{n_1} & \mathbf{0}_{n_1} & \dots & \mathbf{0}_{n_1} \\ \mathbf{1}_{n_2} & \mathbf{0}_{n_2} & \mathbf{1}_{n_2} & \dots & \mathbf{0}_{n_2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{1}_{n_J} & \mathbf{0}_{n_J} & \dots & \mathbf{0}_{n_J} & \mathbf{1}_{n_J} \end{bmatrix} \begin{pmatrix} \mu \\ \tau_1 \\ \vdots \\ \tau_J \end{pmatrix}$$

- Equivalent means  $\mu_j = \mu + \tau_j$
- Should our inference for  $\mu$  depend on how we represent or parameterize  $\mu$ ?

Many equivalent ways to represent the same mean vector –  
inference should be independent of the coordinate system used

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$
- The set of all linear combinations of  $\mathbf{X}_1, \dots, \mathbf{X}_p$  is the space spanned by  $\mathbf{X}_1, \dots, \mathbf{X}_p \equiv S(\mathbf{X}_1, \dots, \mathbf{X}_p)$

# Column Space

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$
- The set of all linear combinations of  $\mathbf{X}_1, \dots, \mathbf{X}_p$  is the space spanned by  $\mathbf{X}_1, \dots, \mathbf{X}_p \equiv S(\mathbf{X}_1, \dots, \mathbf{X}_p)$
- Let  $\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_p]$  be a  $n \times p$  matrix with columns  $\mathbf{X}_j$ ; then the column space of  $\mathbf{X}$ ,  $C(\mathbf{X}) = S(\mathbf{X}_1, \dots, \mathbf{X}_p)$  space spanned by the (column) vectors of  $\mathbf{X}$

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$
- The set of all linear combinations of  $\mathbf{X}_1, \dots, \mathbf{X}_p$  is the space spanned by  $\mathbf{X}_1, \dots, \mathbf{X}_p \equiv S(\mathbf{X}_1, \dots, \mathbf{X}_p)$
- Let  $\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_p]$  be a  $n \times p$  matrix with columns  $\mathbf{X}_j$ ; then the column space of  $\mathbf{X}$ ,  $C(\mathbf{X}) = S(\mathbf{X}_1, \dots, \mathbf{X}_p)$  space spanned by the (column) vectors of  $\mathbf{X}$
- $\boldsymbol{\mu} \in C(\mathbf{X}) : C(\mathbf{X}) = \{\boldsymbol{\mu} \mid \boldsymbol{\mu} \in \mathbb{R}^n \text{ such that } \mathbf{X}\boldsymbol{\beta} = \boldsymbol{\mu} \text{ for some } \boldsymbol{\beta} \in \mathbb{R}^p\}$  (also called the Range of  $\mathbf{X}$ ,  $R(\mathbf{X})$ )

# Column Space

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$
- The set of all linear combinations of  $\mathbf{X}_1, \dots, \mathbf{X}_p$  is the space spanned by  $\mathbf{X}_1, \dots, \mathbf{X}_p \equiv S(\mathbf{X}_1, \dots, \mathbf{X}_p)$
- Let  $\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_p]$  be a  $n \times p$  matrix with columns  $\mathbf{X}_j$ ; then the column space of  $\mathbf{X}$ ,  $C(\mathbf{X}) = S(\mathbf{X}_1, \dots, \mathbf{X}_p)$  space spanned by the (column) vectors of  $\mathbf{X}$
- $\boldsymbol{\mu} \in C(\mathbf{X}) : C(\mathbf{X}) = \{\boldsymbol{\mu} \mid \boldsymbol{\mu} \in \mathbb{R}^n \text{ such that } \mathbf{X}\boldsymbol{\beta} = \boldsymbol{\mu} \text{ for some } \boldsymbol{\beta} \in \mathbb{R}^p\}$  (also called the Range of  $\mathbf{X}$ ,  $R(\mathbf{X})$ )
- $\boldsymbol{\beta}$  are the “coordinates” of  $\boldsymbol{\mu}$  in this space

# Column Space

Many equivalent ways to represent the same mean vector – inference should be independent of the coordinate system used

- Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_p \in \mathbb{R}^n$
- The set of all linear combinations of  $\mathbf{X}_1, \dots, \mathbf{X}_p$  is the space spanned by  $\mathbf{X}_1, \dots, \mathbf{X}_p \equiv S(\mathbf{X}_1, \dots, \mathbf{X}_p)$
- Let  $\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_p]$  be a  $n \times p$  matrix with columns  $\mathbf{X}_j$ ; then the column space of  $\mathbf{X}$ ,  $C(\mathbf{X}) = S(\mathbf{X}_1, \dots, \mathbf{X}_p)$  space spanned by the (column) vectors of  $\mathbf{X}$
- $\boldsymbol{\mu} \in C(\mathbf{X}) : C(\mathbf{X}) = \{\boldsymbol{\mu} \mid \boldsymbol{\mu} \in \mathbb{R}^n \text{ such that } \mathbf{X}\boldsymbol{\beta} = \boldsymbol{\mu} \text{ for some } \boldsymbol{\beta} \in \mathbb{R}^p\}$  (also called the Range of  $\mathbf{X}$ ,  $R(\mathbf{X})$ )
- $\boldsymbol{\beta}$  are the “coordinates” of  $\boldsymbol{\mu}$  in this space
- $C(\mathbf{X})$  is a subspace of  $\mathbb{R}^n$



# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- 1  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- 1  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- 2  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- 1  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- 2  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- 3 there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- 1  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- 2  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- 3 there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$
- 4 for every  $\mathbf{x} \in V$ , there exists a unique vector  $-\mathbf{x}$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- ①  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- ②  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- ③ there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$
- ④ for every  $\mathbf{x} \in V$ , there exists a unique vector  $-\mathbf{x}$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
- ⑤  $\alpha(\beta\mathbf{x}) = (\alpha\beta)\mathbf{x}$  (multiplication by scalars is associative)

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- ①  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- ②  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- ③ there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$
- ④ for every  $\mathbf{x} \in V$ , there exists a unique vector  $-\mathbf{x}$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
- ⑤  $\alpha(\beta\mathbf{x}) = (\alpha\beta)\mathbf{x}$  (multiplication by scalars is associative)
- ⑥  $1\mathbf{x} = \mathbf{x}$  for every  $\mathbf{x}$

# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- ①  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- ②  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- ③ there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$
- ④ for every  $\mathbf{x} \in V$ , there exists a unique vector  $-\mathbf{x}$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
- ⑤  $\alpha(\beta\mathbf{x}) = (\alpha\beta)\mathbf{x}$  (multiplication by scalars is associative)
- ⑥  $1\mathbf{x} = \mathbf{x}$  for every  $\mathbf{x}$
- ⑦  $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}$  (multiplication by scalars is distributive with respect vector addition)



# Vector spaces

A collection of vectors  $V$  is a real **vector space** if the following conditions hold: for any pair  $\mathbf{x}$  and  $\mathbf{y}$  of vectors in  $V$  there corresponds a vector  $\mathbf{x} + \mathbf{y}$  and scalars  $\alpha, \beta \in \mathbb{R}$  such that:

- ①  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  (vector addition is commutative)
- ②  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  (vector addition is associative)
- ③ there exists a unique vector  $\mathbf{0} \in V$  (the origin) such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every vector  $\mathbf{x}$
- ④ for every  $\mathbf{x} \in V$ , there exists a unique vector  $-\mathbf{x}$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
- ⑤  $\alpha(\beta\mathbf{x}) = (\alpha\beta)\mathbf{x}$  (multiplication by scalars is associative)
- ⑥  $1\mathbf{x} = \mathbf{x}$  for every  $\mathbf{x}$
- ⑦  $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}$  (multiplication by scalars is distributive with respect vector addition)
- ⑧  $(\alpha + \beta)\mathbf{x} = \alpha\mathbf{x} + \beta\mathbf{x}$  (multiplication by scalars is distributive with respect to vector addition)

## Definition

Let  $V$  be a vector space and let  $V_0$  be a set with  $V_0 \subseteq V$ .  $V_0$  is a *subspace* of  $V$  if and only if  $V_0$  is a vector space.

# Subspaces

## Definition

Let  $V$  be a vector space and let  $V_0$  be a set with  $V_0 \subseteq V$ .  $V_0$  is a *subspace* of  $V$  if and only if  $V_0$  is a vector space.

## Theorem

*Let  $V$  be a vector space, and let  $V_0$  be a non-empty subset of  $V$ . If  $V_0$  is closed on vector addition and scalar multiplication, then  $V_0$  is a subspace of  $V$*

# Subspaces

## Definition

Let  $V$  be a vector space and let  $V_0$  be a set with  $V_0 \subseteq V$ .  $V_0$  is a *subspace* of  $V$  if and only if  $V_0$  is a vector space.

## Theorem

*Let  $V$  be a vector space, and let  $V_0$  be a non-empty subset of  $V$ . If  $V_0$  is closed on vector addition and scalar multiplication, then  $V_0$  is a subspace of  $V$*

## Theorem

*Let  $V$  be a vector space, and let  $\mathbf{x}_1, \dots, \mathbf{x}_r$  be vectors in  $V$ . The set of all linear combinations of  $\mathbf{x}_1, \dots, \mathbf{x}_r$  is a subspace of  $V$ .*

# Subspaces

## Definition

Let  $V$  be a vector space and let  $V_0$  be a set with  $V_0 \subseteq V$ .  $V_0$  is a *subspace* of  $V$  if and only if  $V_0$  is a vector space.

## Theorem

*Let  $V$  be a vector space, and let  $V_0$  be a non-empty subset of  $V$ . If  $V_0$  is closed on vector addition and scalar multiplication, then  $V_0$  is a subspace of  $V$*

## Theorem

*Let  $V$  be a vector space, and let  $\mathbf{x}_1, \dots, \mathbf{x}_r$  be vectors in  $V$ . The set of all linear combinations of  $\mathbf{x}_1, \dots, \mathbf{x}_r$  is a subspace of  $V$ .*

The Column space of  $\mathbf{X}$ ,  $C(\mathbf{X})$ , is a subspace of  $\mathbb{R}^n$

## Definition

A finite set of vectors  $\{\mathbf{x}_i\}$  is *linearly dependent* if there exist scalars  $\{a_i\}$  (not all zero) such that  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$ .

## Definition

A finite set of vectors  $\{\mathbf{x}_i\}$  is *linearly dependent* if there exist scalars  $\{a_i\}$  (not all zero) such that  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$ . Conversely  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$  implies that  $a_i = 0 \forall i$  the set is *linearly independent*.

## Definition

A finite set of vectors  $\{\mathbf{x}_i\}$  is *linearly dependent* if there exist scalars  $\{a_i\}$  (not all zero) such that  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$ . Conversely  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$  implies that  $a_i = 0 \forall i$  the set is *linearly independent*.

## Definition

If  $V_0$  is a subspace of  $V$  and if  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a linearly independent spanning set for  $V_0$ , then  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a basis for  $V_0$



## Definition

A finite set of vectors  $\{\mathbf{x}_i\}$  is *linearly dependent* if there exist scalars  $\{a_i\}$  (not all zero) such that  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$ . Conversely  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$  implies that  $a_i = 0 \forall i$  the set is *linearly independent*.

## Definition

If  $V_0$  is a subspace of  $V$  and if  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a linearly independent spanning set for  $V_0$ , then  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a basis for  $V_0$

## Theorem

*If  $V_0$  is a subspace of  $V$ , then all bases for  $V_0$  have the same number of vectors*

## Definition

A finite set of vectors  $\{\mathbf{x}_i\}$  is *linearly dependent* if there exist scalars  $\{a_i\}$  (not all zero) such that  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$ . Conversely  $\sum_i a_i \mathbf{x}_i = \mathbf{0}$  implies that  $a_i = 0 \forall i$  the set is *linearly independent*.

## Definition

If  $V_0$  is a subspace of  $V$  and if  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a linearly independent spanning set for  $V_0$ , then  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is a basis for  $V_0$

## Theorem

*If  $V_0$  is a subspace of  $V$ , then all bases for  $V_0$  have the same number of vectors*

Can both the collection of vectors in the cell means and the treatment effects parameterizations be a basis?

## Definition

The rank of a subspace  $V_0$  is the number of elements in a basis for  $V_0$  and is written as  $r(V_0)$ . Similarly if  $\mathbf{A}$  is a matrix, the rank of  $C(\mathbf{A})$  is called the rank of  $\mathbf{A}$  and is written  $r(\mathbf{A})$ .

## Definition

The rank of a subspace  $V_0$  is the number of elements in a basis for  $V_0$  and is written as  $r(V_0)$ . Similarly if  $\mathbf{A}$  is a matrix, the rank of  $C(\mathbf{A})$  is called the rank of  $\mathbf{A}$  and is written  $r(\mathbf{A})$ .

What is the rank of the subspace in the Oneway ANOVA model?

## Definition

An inner product space is a vector space  $V$  equipped with an inner product:  $\langle \cdot, \cdot \rangle$  is a mapping  $V \times V \rightarrow \mathbb{R}$ . Two vectors are orthogonal if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ , written  $\mathbf{x} \perp \mathbf{y}$

# Orthogonality

## Definition

An inner product space is a vector space  $V$  equipped with an inner product:  $\langle \cdot, \cdot \rangle$  is a mapping  $V \times V \rightarrow \mathbb{R}$ . Two vectors are orthogonal if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ , written  $\mathbf{x} \perp \mathbf{y}$  (usual  $\mathbf{x}^T \mathbf{y} = 0$ )

## Definition

Two subspaces are orthogonal if  $\mathbf{x} \in M$  and  $\mathbf{y} \in N$  imply that  $\mathbf{x} \perp \mathbf{y}$

# Orthogonality

## Definition

An inner product space is a vector space  $V$  equipped with an inner product:  $\langle \cdot, \cdot \rangle$  is a mapping  $V \times V \rightarrow \mathbb{R}$ . Two vectors are orthogonal if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ , written  $\mathbf{x} \perp \mathbf{y}$  (usual  $\mathbf{x}^T \mathbf{y} = 0$ )

## Definition

Two subspaces are orthogonal if  $\mathbf{x} \in M$  and  $\mathbf{y} \in N$  imply that  $\mathbf{x} \perp \mathbf{y}$

## Definition

Orthonormal basis:  $\{\mathbf{x}_1, \dots, \mathbf{x}_r\}$  is an orthonormal basis (ONB) for  $M$  if  $\langle \mathbf{x}_i, \mathbf{x}_j \rangle = 0$  for  $i \neq j$  and  $\langle \mathbf{x}_i, \mathbf{x}_i \rangle = 1$  for all  $i$ . Length  $\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$ . Distance of two vectors is  $\|\mathbf{x} - \mathbf{y}\|$

Find an orthonormal basis for the oneway ANOVA model. Is it unique?



Find an orthonormal basis for the oneway ANOVA model. Is it unique?

Can also use Gram-Schmidt sequential orthogonalization

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$
- $M + N = V$  where  $M + N = \{\mathbf{z} \mid \mathbf{z} = \mathbf{x} + \mathbf{y}, \mathbf{x} \in M, \mathbf{y} \in N\}$   
( $M \oplus N = V$ ) Direct Sum

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$
- $M + N = V$  where  $M + N = \{\mathbf{z} \mid \mathbf{z} = \mathbf{x} + \mathbf{y}, \mathbf{x} \in M, \mathbf{y} \in N\}$   
( $M \oplus N = V$ ) Direct Sum

then  $M$  and  $N$  are complementary spaces.

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$
- $M + N = V$  where  $M + N = \{\mathbf{z} \mid \mathbf{z} = \mathbf{x} + \mathbf{y}, \mathbf{x} \in M, \mathbf{y} \in N\}$   
( $M \oplus N = V$ ) Direct Sum

then  $M$  and  $N$  are complementary spaces.  $N$  is the complement of  $M$

# Decomposition

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$
- $M + N = V$  where  $M + N = \{\mathbf{z} \mid \mathbf{z} = \mathbf{x} + \mathbf{y}, \mathbf{x} \in M, \mathbf{y} \in N\}$   
( $M \oplus N = V$ ) Direct Sum

then  $M$  and  $N$  are complementary spaces.  $N$  is the complement of  $M$

## Definition

If  $M$  and  $N$  are complementary spaces  $M + N = \mathbb{R}^n$  and  $M$  and  $N$  are orthogonal subspaces, then  $M$  is the orthogonal complement of  $N$ ,  $N^\perp$ .

## Definition

If  $M$  and  $N$  are subspaces of  $V$  that satisfy

- $M \cap N = \{\mathbf{0}\}$
- $M + N = V$  where  $M + N = \{\mathbf{z} \mid \mathbf{z} = \mathbf{x} + \mathbf{y}, \mathbf{x} \in M, \mathbf{y} \in N\}$   
( $M \oplus N = V$ ) Direct Sum

then  $M$  and  $N$  are complementary spaces.  $N$  is the complement of  $M$

## Definition

If  $M$  and  $N$  are complementary spaces  $M + N = \mathbb{R}^n$  and  $M$  and  $N$  are orthogonal subspaces, then  $M$  is the orthogonal complement of  $N$ ,  $N^\perp$ .

If  $\mathbf{z} \in \mathbb{R}^n = N + M$ , then we can uniquely decompose it into a part  $\mathbf{x} \in M$  and  $\mathbf{y} \in N$  and  $r(M) + r(N) = n$

## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if



## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if

- $\mathbf{u} \in C(\mathbf{X})$  implies  $P\mathbf{u} = \mathbf{u}$  (projection)

## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if

- $\mathbf{u} \in C(\mathbf{X})$  implies  $P\mathbf{u} = \mathbf{u}$  (projection)
- $\mathbf{v} \perp C(\mathbf{X})$  implies  $P\mathbf{v} = \mathbf{0}$  (perpendicular)

## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if

- $\mathbf{u} \in C(\mathbf{X})$  implies  $P\mathbf{u} = \mathbf{u}$  (projection)
- $\mathbf{v} \perp C(\mathbf{X})$  implies  $P\mathbf{v} = \mathbf{0}$  (perpendicular)
- $C(\mathbf{X})$  and  $C(\mathbf{X})^\perp$  are complementary spaces

## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if

- $\mathbf{u} \in C(\mathbf{X})$  implies  $P\mathbf{u} = \mathbf{u}$  (projection)
  - $\mathbf{v} \perp C(\mathbf{X})$  implies  $P\mathbf{v} = \mathbf{0}$  (perpendicular)
- 
- $C(\mathbf{X})$  and  $C(\mathbf{X})^\perp$  are complementary spaces
  - $\mathbf{y} = \mathbf{u} + \mathbf{v}$  for  $\mathbf{y} \in \mathbb{R}^n$ ,  $\mathbf{u} \in C(\mathbf{X})$  and  $\mathbf{v} \perp C(\mathbf{X})$  then  
 $P\mathbf{y} = P\mathbf{u} + P\mathbf{v} = \mathbf{u}$

## Definition

$P$  is an orthogonal projection operator onto  $C(\mathbf{X})$  if and only if

- $\mathbf{u} \in C(\mathbf{X})$  implies  $P\mathbf{u} = \mathbf{u}$  (projection)
  - $\mathbf{v} \perp C(\mathbf{X})$  implies  $P\mathbf{v} = \mathbf{0}$  (perpendicular)
- 
- $C(\mathbf{X})$  and  $C(\mathbf{X})^\perp$  are complementary spaces
  - $\mathbf{y} = \mathbf{u} + \mathbf{v}$  for  $\mathbf{y} \in \mathbb{R}^n$ ,  $\mathbf{u} \in C(\mathbf{X})$  and  $\mathbf{v} \perp C(\mathbf{X})$  then  
 $P\mathbf{y} = P\mathbf{u} + P\mathbf{v} = \mathbf{u}$
  - $\mathbf{u}$  is the orthogonal projection of  $\mathbf{y}$  onto  $C(\mathbf{X})$

Prop

$P$  is an orthogonal projection operator on  $C(\mathbf{X})$  then  $C(\mathbf{X}) = C(P)$

# More on Projections

## Prop

$P$  is an orthogonal projection operator on  $C(\mathbf{X})$  then  $C(\mathbf{X}) = C(P)$

## Theorem

*$P$  is an orthogonal projection operator on  $C(P)$  if and only if*

## Prop

$P$  is an orthogonal projection operator on  $C(\mathbf{X})$  then  $C(\mathbf{X}) = C(P)$

## Theorem

*$P$  is an orthogonal projection operator on  $C(P)$  if and only if*

- $P = P^2$  (idempotent)



## Prop

$P$  is an orthogonal projection operator on  $C(\mathbf{X})$  then  $C(\mathbf{X}) = C(P)$

## Theorem

*$P$  is an orthogonal projection operator on  $C(P)$  if and only if*

- $P = P^2$  (idempotent)
- $P = P^T$  (symmetry)

# More on Projections

## Prop

$P$  is an orthogonal projection operator on  $C(\mathbf{X})$  then  $C(\mathbf{X}) = C(P)$

## Theorem

*$P$  is an orthogonal projection operator on  $C(P)$  if and only if*

- $P = P^2$  (idempotent)
- $P = P^T$  (symmetry)

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$   
operator onto  $C(\mathbf{X})$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $P = P^2$  (idempotent)

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $P = P^2$  (idempotent)

$$P_{\mathbf{X}}^2 = P_{\mathbf{X}} P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $P = P^2$  (idempotent)

$$\begin{aligned} P_{\mathbf{X}}^2 = P_{\mathbf{X}} P_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \end{aligned}$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $P = P^2$  (idempotent)

$$\begin{aligned} P_{\mathbf{X}}^2 = P_{\mathbf{X}} P_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= P_{\mathbf{X}} \end{aligned}$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $P_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $P = P^2$  (idempotent)

$$\begin{aligned} P_{\mathbf{X}}^2 = P_{\mathbf{X}} P_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= P_{\mathbf{X}} \end{aligned}$$

- $P = P^T$  (symmetry)



# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $\mathbf{P}_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_{\mathbf{X}}^2 = \mathbf{P}_{\mathbf{X}} \mathbf{P}_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_{\mathbf{X}}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\mathbf{P}_{\mathbf{X}}^T = (\mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $\mathbf{P}_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_{\mathbf{X}}^2 = \mathbf{P}_{\mathbf{X}} \mathbf{P}_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_{\mathbf{X}}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_{\mathbf{X}}^T &= (\mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T ((\mathbf{X}^T \mathbf{X})^{-1})^T (\mathbf{X})^T\end{aligned}$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $\mathbf{P}_{\mathbf{X}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_{\mathbf{X}}^2 = \mathbf{P}_{\mathbf{X}} \mathbf{P}_{\mathbf{X}} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_{\mathbf{X}}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_{\mathbf{X}}^T &= (\mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T ((\mathbf{X}^T \mathbf{X})^{-1})^T (\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T\end{aligned}$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T((\mathbf{X}^T\mathbf{X})^{-1})^T(\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

# Projections

Claim: If  $\mathbf{X}$  is  $n \times p$  and  $r(\mathbf{X}) = p$  then,  $\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$  is an orthogonal projection operator onto  $C(\mathbf{X})$

- $\mathbf{P} = \mathbf{P}^2$  (idempotent)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^2 = \mathbf{P}_\mathbf{X} \mathbf{P}_\mathbf{X} &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $\mathbf{P} = \mathbf{P}^T$  (symmetry)

$$\begin{aligned}\mathbf{P}_\mathbf{X}^T &= (\mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\ &= (\mathbf{X}^T)^T ((\mathbf{X}^T \mathbf{X})^{-1})^T (\mathbf{X})^T \\ &= \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \\ &= \mathbf{P}_\mathbf{X}\end{aligned}$$

- $C(\mathbf{X}) = C(\mathbf{P}_\mathbf{X})$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 = (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}})$$



Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\ &= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_{\mathbf{X}}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_{\mathbf{X}})^2 &= (\mathbf{I} - \mathbf{P}_{\mathbf{X}})(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\mathbf{P}_{\mathbf{X}} \\&= \mathbf{I} - \mathbf{P}_{\mathbf{X}} - \mathbf{P}_{\mathbf{X}} + \mathbf{P}_{\mathbf{X}}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_\mathbf{X} = (\mathbf{I} - \mathbf{P}_\mathbf{X})^T$

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_\mathbf{X} = (\mathbf{I} - \mathbf{P}_\mathbf{X})^T$
- $\mathbf{u} \in C(\mathbf{X})^\perp \Rightarrow \mathbf{u} \perp C(\mathbf{X})$  and  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{u} = \mathbf{u}$  (projection)

Claim:  $\mathbf{I} - \mathbf{P}_\mathbf{X}$  is an orthogonal projection onto  $C(\mathbf{X})^\perp$

- idempotent

$$\begin{aligned}(\mathbf{I} - \mathbf{P}_\mathbf{X})^2 &= (\mathbf{I} - \mathbf{P}_\mathbf{X})(\mathbf{I} - \mathbf{P}_\mathbf{X}) \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X}\mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X} - \mathbf{P}_\mathbf{X} + \mathbf{P}_\mathbf{X} \\&= \mathbf{I} - \mathbf{P}_\mathbf{X}\end{aligned}$$

- Symmetry  $\mathbf{I} - \mathbf{P}_\mathbf{X} = (\mathbf{I} - \mathbf{P}_\mathbf{X})^T$
- $\mathbf{u} \in C(\mathbf{X})^\perp \Rightarrow \mathbf{u} \perp C(\mathbf{X})$  and  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{u} = \mathbf{u}$  (projection)
- if  $\mathbf{v} \in C(\mathbf{X})$ ,  $(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{v} = \mathbf{v} - \mathbf{v} = \mathbf{0}$

# Null Space

- Column space  $C(\mathbf{X})$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = 0\}$



# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = 0\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = 0\}$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = 0\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = 0\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = 0\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = 0\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(P_{\mathbf{X}}) = C(\mathbf{X})^\perp$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = \mathbf{0}\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = \mathbf{0}\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(\mathbf{P}_\mathbf{X}) = C(\mathbf{X})^\perp$

$$\mathbf{u} \in N(\mathbf{P}) \Rightarrow \mathbf{P}_\mathbf{X}\mathbf{u} = \mathbf{0}$$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = \mathbf{0}\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = \mathbf{0}\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(\mathbf{P}_\mathbf{X}) = C(\mathbf{X})^\perp$

$$\begin{aligned}\mathbf{u} \in N(\mathbf{P}) &\Rightarrow \mathbf{P}_\mathbf{X}\mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{u} = \mathbf{0}\end{aligned}$$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = \mathbf{0}\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = \mathbf{0}\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(\mathbf{P}_\mathbf{X}) = C(\mathbf{X})^\perp$

$$\begin{aligned}\mathbf{u} \in N(\mathbf{P}) &\Rightarrow \mathbf{P}_\mathbf{X}\mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{u}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T = \mathbf{0}^T\end{aligned}$$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = \mathbf{0}\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = \mathbf{0}\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(P_{\mathbf{X}}) = C(\mathbf{X})^\perp$

$$\begin{aligned}\mathbf{u} \in N(P) &\Rightarrow P_{\mathbf{X}}\mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{u}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T = \mathbf{0}^T \\ &\Leftrightarrow \mathbf{u}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\boldsymbol{\beta} = 0\end{aligned}$$

# Null Space

- Column space  $C(\mathbf{X})$
- $N(\mathbf{A})$ : Null space of  $\mathbf{A}$  is  $\{\mathbf{u} \mid \mathbf{A}\mathbf{u} = 0\}$
- Null space of  $\mathbf{X}^T$  is  $\{\mathbf{u} \mid \mathbf{X}^T\mathbf{u} = 0\}$
- $N(\mathbf{X}^T) = C(\mathbf{X})^\perp$  the orthogonal complement of  $C(\mathbf{X})$
- $N(P_X) = C(\mathbf{X})^\perp$

$$\begin{aligned}\mathbf{u} \in N(P) &\Rightarrow P_X \mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{u} = \mathbf{0} \\ &\Leftrightarrow \mathbf{u}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T = \mathbf{0}^T \\ &\Leftrightarrow \mathbf{u}^T \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \boldsymbol{\beta} = 0 \\ &\Leftrightarrow \mathbf{u}^T \mathbf{v} = 0 \quad \mathbf{v} \in C(\mathbf{X})\end{aligned}$$



- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_X \mathbf{Y}$

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_X \mathbf{Y}$
- Log Likelihood:

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X} \mathbf{Y}$
- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X} \mathbf{Y}$
- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_\mathbf{X} \mathbf{Y} + (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}$

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X} \mathbf{Y}$
- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_\mathbf{X} \mathbf{Y} + (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}$
- Use  $\mathbf{P}_\mathbf{X} \boldsymbol{\mu} = \boldsymbol{\mu}$

# Models Again

- $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  with  $\boldsymbol{\mu} \in C(\mathbf{X})$
- Claim: Maximum Likelihood Estimator (MLE) of  $\boldsymbol{\mu}$  is  $\mathbf{P}_\mathbf{X} \mathbf{Y}$
- Log Likelihood:

$$\log \mathcal{L}(\boldsymbol{\mu}, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \frac{\|\mathbf{Y} - \boldsymbol{\mu}\|^2}{\sigma^2}$$

- Decompose  $\mathbf{Y} = \mathbf{P}_\mathbf{X} \mathbf{Y} + (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}$
- Use  $\mathbf{P}_\mathbf{X} \boldsymbol{\mu} = \boldsymbol{\mu}$
- and Simplify

$$\|\mathbf{Y} - \boldsymbol{\mu}\|^2$$