

Checking Assumptions

Merlise Clyde

STA721 Linear Models

Duke University

November 16, 2015

Topics

- Normality
- Brain Weights and Body Mass
- Box-Cox

Readings: Christensen Chapter 13

Linear Model:

$$\mathbf{Y} = \boldsymbol{\mu} + \boldsymbol{\epsilon}$$

Assumptions:

$$\begin{aligned}\boldsymbol{\mu} \in C(\mathbf{X}) &\Leftrightarrow \boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta} \\ \boldsymbol{\epsilon} &\sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)\end{aligned}$$

What could go wrong?

- Wrong mean for a case or cases (outliers)
- Cases that influence the mean (influential cases)
- Both of the above
- Wrong distribution for $\boldsymbol{\epsilon}$

Recall

$$\begin{aligned}\mathbf{e} &= (\mathbf{I} - \mathbf{P}_X)\mathbf{Y} \\ &= (\mathbf{I} - \mathbf{P}_X)(\mathbf{X}\hat{\beta} + \epsilon) \\ &= (\mathbf{I} - \mathbf{P}_X)\epsilon\end{aligned}$$


$$e_i = \epsilon_i - \sum_{j=1}^n h_{ij}\epsilon_j$$

Lyapunov CLT implies that residuals will be approximately normal (even for modest n), if the errors are not normal


“Supernormality of residuals”

- Order e_i : $e_{(1)} \leq e_{(2)} \dots \leq e_{(n)}$ sample order statistics or sample quantiles
- Let $z_{(1)} \leq z_{(2)} \dots z_{(n)}$ denote the expected order statistics of a sample of size n from a standard normal distribution “theoretical quantiles”
- If the e_i are normal then $E[e_{(i)}] = \sigma z_{(i)}$
- Expect that points in a scatter plot of $e_{(i)}$ and $z_{(i)}$ should be on a straight line.
- Judgment call - use simulations to gain experience!

Animal Example



Residual Plots



Box-Cox Transformation

Box and Cox (1964) suggested a family of power transformations for $Y > 0$

$$U(\mathbf{Y}, \lambda) = Y^{(\lambda)} = \begin{cases} \frac{(Y^\lambda - 1)}{\lambda} & \lambda \neq 0 \\ \log(Y) & \lambda = 0 \end{cases}$$


- Estimate λ by maximum Likelihood

$$\mathcal{L}(\lambda, \beta, \sigma^2) \propto \prod f(y_i | \lambda, \beta, \sigma^2)$$





- $U(\mathbf{Y}, \lambda) = Y^{(\lambda)} \sim N(\mathbf{X}\beta, \sigma^2)$
- Jacobian term is $\prod_i y_i^{\lambda-1}$ for all λ
- Profile Likelihood based on substituting MLE β and σ^2 for each value of λ is

$$\log(\mathcal{L}(\lambda)) \propto (\lambda - 1) \sum_i \log(Y_i) - \frac{n}{2} \log(\text{SSE}(\lambda))$$





Profile Likelihood



Residuals After Transformation of Response



Residuals After Transformation of Both



Transformed Data

Test that Dinos are Outliers

| | Res.Df | RSS | Df | Sum of Sq | F | Pr(>F) |
|---|--------|-------|----|-----------|-------|--------|
| 1 | 23 | 12.12 | | | | |
| 2 | 26 | 60.99 | -3 | -48.87 | 30.92 | 0.0000 |

| | Estimate | Std. Error | t value | Pr(> t) |
|---------------|----------|------------|---------|----------|
| (Intercept) | 2.1504 | 0.2006 | 10.72 | 0.0000 |
| log(body) | 0.7523 | 0.0457 | 16.45 | 0.0000 |
| Triceratops | -4.7839 | 0.7913 | -6.05 | 0.0000 |
| Brachiosaurus | -5.6662 | 0.8328 | -6.80 | 0.0000 |
| Dipliodocus | -5.2851 | 0.7949 | -6.65 | 0.0000 |

Dinosaurs come from a different population from mammals

To Remove or Not?

- For suspicious cases, check data sources for errors
- Check that points are not outliers because of wrong mean function or distributional assumptions
- Investigate need for transformations (use EDA at several stages)
- Influential cases - report results with and without cases (results may change - are differences meaningful?)
- Outlier test - suggests alternative population for the case(s); if not influential may in keep analysis, but will inflate $\hat{\sigma}^2$ and interval estimates
- Document how you handle any case deletions - reproducibility!
- Consider BMA with outliers (See BMA package) to address model uncertainty
- Robust Regression Methods

Variance Stabilizing Transformations

- If $Y - \mu$ (approximately) $N(0, h(\mu))$
- Delta Method implies that

$$g(Y) \dot{\sim} N(g(\mu), g'(\mu)^2 h(\mu))$$

- Find function g such that $g'(\mu)^2/h(\mu)$ is constant

$$g(Y) \sim N(g(\mu), c)$$

- Poisson Counts: g is square root transformation
- Binomial: $\arcsin(\sqrt{Y})$