



Short communication

The importance of non-academic coauthors in bridging the conservation genetics gap

Meghan Britt^a, Sarah E. Haworth^a, Justin B. Johnson^b, Daria Martchenko^b, Aaron B.A. Shafer^{a,b,*}^a Forensics Program, Trent University, 7K9J 7B8 Peterborough, Canada^b Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough, Canada

ARTICLE INFO

Keywords:
Meta-analysis
Genomics
Management
Wildlife

ABSTRACT

DNA sequence data from genetic (traditional) and genomic (high-throughput) instruments have been highlighted as important information for biodiversity conservation. Despite a high number of publications, a gap has emerged between fundamental research and real-world application, meaning genetic studies often do not have a substantive link to policy or conservation science. We conducted a meta-analysis on conservation studies using genetic and genomic datasets and found the majority of publications were on species that were data deficient or of low conservation priority (66%); only 38% of studies specifically identified how the data could inform conservation or management through clearly stated recommendations for action or policy. Importantly, our mixed-model found a 2.5-fold increase in the odds of a publication making a specific recommendation when non-academic affiliated authors were included. This suggests the conservation genetics gap has arisen, in part, by academic authors failing to make explicit links to policy and conservation outcomes. Conservation geneticists should refrain from hyperbole and collaborate with agency and non-government scientists to accelerate development and implementation, and ultimately help bridge the conservation genetics gap.

1. Introduction

Conserving biodiversity is a difficult task fraught with competing interests and often limited knowledge of species and biological systems. Governments, academics, and conservation practitioners are faced with the immense challenge of stemming the rapid loss of species and habitat under limited budgets in an increasingly polarized world. Working in an interdisciplinary science, conservation biologists are quick to try new tools, and any promising technological advancements garner much attention in this regard (Pimm et al., 2014). Genetics and genomics, the latter defined as high-throughput sampling of nucleic acids (McMahon et al., 2014), have been highlighted as important tools for conservation (Frankham, 2003; Ouborg, 2010; Ouborg et al., 2010). Genetics has numerous applications of conservation relevance such as mark-recapture, parentage and population assignment, and estimates of effective population size; these data have made clear inroads with well-known examples like Atlantic Salmon management (Nielsen, 1998), seed collection programs (Silva et al., 2008), and the restoration of the Florida Panther (Johnson et al., 2010). The impact of genomic data on conservation is less clear and debated (see Cronin, 2007; Shafer et al., 2015, 2016; Garner et al., 2016); in principle, genomic data should

increase the precision and accuracy of demographic estimates and allow for the identification of adaptive loci (McMahon et al., 2014). It has also been suggested that analysis of whole genomes will become routine task in conservation biology (Fuentes-Pardo and Ruzzante, n.d.).

Despite the high number of publications and noteworthy examples, a gap has emerged between basic genetic and genomic research, and real-world applications; this has been dubbed the conservation genetics gap and is reflected by the limited integration of genetic data into conservation management (Hoban et al., 2013b; Shafer et al., 2015; Haig et al., 2015; Taylor et al., 2017). European researchers were among the first to recognize the gap and in 2012 initiated the project known as Conservation Genetic Resources for Effective Species Survival (ConGRESS); ConGRESS had the stated goal of bridging the gap and providing (genetic) resources for biodiversity managers and policy makers (Hoban et al., 2013a). Following a European Science Foundation funded meeting on the role of genomics in conservation, Shafer et al. (2015) argued that the gap would only become exacerbated in the genomics era due to the infancy of the field and uncertainty surrounding datasets and bioinformatics tools. In response, Garner et al. (2016) provided a list of genomic case studies showing real-world

* Corresponding author at: 2140 East Bank Dr., Peterborough, Ontario K9J 7B8, Canada.

E-mail addresses: meghanbritt@trentu.ca (M. Britt), sarahhaworth@trentu.ca (S.E. Haworth), justinjohnson@trentu.ca (J.B. Johnson), dariamartchenko@trentu.ca (D. Martchenko), aaronshafer@trentu.ca (A.B.A. Shafer).

<https://doi.org/10.1016/j.biocon.2017.12.019>

Received 19 September 2017; Received in revised form 9 November 2017; Accepted 13 December 2017
0006-3207/ © 2017 Elsevier Ltd. All rights reserved.

application, although these examples were restricted to economically important species and the widespread implementation of genomic tools was still a limiting factor (Shafer et al., 2016).

It is clear that some disconnect exists between genetics research and its application to conservation issues. In a survey of 150 New Zealand conservation practitioners, Taylor et al. (2017) found that access to expertise and funding were the main barriers to uptake and implementation. While access to funds and expertise might be limiting factors, it does not explain the absence of genetic targets and tools in conservation priority and implementation plans (Carter et al., 2000; Groves et al., 2002; Groves, 2003; Knight et al., 2006). Similarly, only 1 of the 100 so-called questions of importance to the conservation of global biological diversity mentioned genetics, and this was in relation to genetically modified organisms (Sutherland et al., 2009). Collectively, this suggests that many, perhaps most, conservation problems do not need genetic or genomic solutions and leads to an alternative explanation of the conservation genetics gap that has to do with the perception of molecular tools having limited use in solving most conservation problems.

The idea of a perception problem is not unfounded (Amos and Balmford, 2001). When surveyed, stakeholders ranked access to genetic resources 18th out of a list of 35 priority issues (Nagulendran et al., 2016). Anecdotal, such a low ranking for molecular tools among conservation priorities and planning is not uncommon. Even though the majority of practitioners said they would use genetic data if it were available (Hoban et al., 2013b), the perceived lack of real-world applications suggests more than expertise and access being a barrier. When presented with a problem, practitioners and committees generally consult the scientific literature and there is an increasing value placed on peer-reviewed conservation science (Game et al., 2015). If publications provide nebulous applications and inaccessible tools with respect to a subdiscipline like conservation genetics, this could create a scenario where practitioners might simply not seek out tools like genetic and genomic assays because their real-world applications are not clear (sensu Soulé, 1985). Here, we conducted a meta-analysis of conservation genetic and genomic studies and collected data on author affiliation, the conservation status of studied species, the molecular marker used, and the conservation recommendations made in the study. The aim of this meta-analysis was to shed light on the actors and drivers behind the conservation genetic research-implementation gap, and offer solutions to better link genetic and genomic research to policy and conservation action.

2. Methods

We collected data from four peer-reviewed journals; two of which are conservation-oriented and publish genetic research, while the other two focus on genetics research but include papers of conservation interest. The journals were: Conservation Genetics (CG), Biological Conservation (BC), Evolutionary Applications (EA), and Molecular Ecology (ME). Additional conservation journals were screened but had a small number of genetic and genomic studies and were excluded. Using the keyword search “genetics” within CG, BC, and EA, original empirical research articles from the past ten years (2006–2017) were randomly collected and retained if they focused on a specific species, or multiple named species. The search term “genomics” was also entered within CG, BC, and EA and as many articles as possible were collected based on the aforementioned criteria. Within ME the keyword “conservation” was searched because the journal publishes more than conservation research. We aimed to retain 75 studies from each of the four journals to achieve a representative sample of the literature.

The following information was collected from each article: publication year, number of authors, author affiliations (i.e. government, academic, or non-governmental), focal species and taxonomic grouping (class or kingdom), the species' current global conservation status, the species' regional status if reported in the article (NA if not), the class of

molecular marker(s) used (genetics being microsatellites, single gene sequencing, low-throughput single nucleotide polymorphism (SNP) assays; genomics being tools using high-throughput sequencing or genotyping such as RADseq, SNP chips, whole-genome re-sequencing). In the event that more than one species was included in an article, only one, randomly selected species was included in our analysis. The current global conservation status of each species was obtained from the IUCN RedList or NatureServe. We merged the two ranking systems to create a conservation status scoring system ranging from 0 (i.e. least concern) to 2 (i.e. critically endangered; Table S1). In the event that a species was not listed in one of these collections or article they were given a NA in our database.

We scored the specificity of conservation recommendations provided in each article. We considered recommendations on a binary scale – generic or specific – where a generic recommendation would be broad statements with no explicit direction on how the data can be used to inform conservation and management, or reference to a change in policy or legislation (see also Table S2 for scoring criteria). For example, a generic recommendation would be “we propose maintaining genetic diversity of the species to ensure long-term viability.” This statement reflects a truism in conservation biology. An article was ranked as having a specific conservation recommendation if there was a clear course of action suggested, stated implementation methods, or policy changes that were advocated for. A specific example would be “we propose translocating 10 animals from the mainland to limit genetic erosion and maintain 90% of current genetic variation.” This statement gives direction to a conservation action informed by the genetic or genomic data. If an article proposed an Evolutionary Significant Unit (ESU) or Management Unit (MU) it was scored as specific. All articles were independently scored at least twice.

All summary statistics, statistical models, and data visualizations were conducted in R v. 3.4.1. The repeatability of scoring the conservation recommendation, meaning the same recommendation score was given by both evaluators, was assessed using Krippendorff's α -Reliability, where an α of 1 equals perfect reliability and 0 equates to random chance. As the data were subdivided into two groups (repeated and non-repeated conservation recommendations), we compared the authorship composition (i.e. number of academic, government, non-government authors) using a multivariate analysis of variance (MANOVA). The remaining analysis focused only on the repeated data set. A number of generalized linear mixed models were fitted to the data with the journal in which the study was published as a random effect (i.e., intercepts varying by journal). The four tested predictor variables used were proportion of non-academic authors (to simplify the model we merged government and non-government authors into one variable for non-academic affiliations), global conservation status (categorical values of 0, 1, 2, described in Table S1), marker type and taxonomic group (categories listed in Table 1). Factors that were significant based on an α of 0.05 in a univariate model with a random effect were added to the final model. To aid in interpretation of the logistic model coefficients were converted to odds-ratios, which is the relative odds of the outcome given a condition.

3. Results

In total we reviewed 300 publications from 2006 to 2017 in four scientific journals that focused specifically on, or have a section devoted to, conservation and wildlife management. The mean authorship breakdown and taxonomic affiliation of research organisms, seen in Table 1, shows that academic affiliated authors constituted the majority of coauthors (61–70%), followed by government (18–27%) and non-government (7–18%). We had a scoring repeatability of 77% (231 of 300 studies) which is better than chance (Krippendorff's α = 0.56). The authorship breakdown did not differ between repeated and non-repeated studies based on a MANOVA (Pillai's trace = 0.003, $F(1, 298)$ = 0.323, p = 0.808). The majority of studies were on species of low

Table 1
Summary of data collected from 300 genetic or genomic studies of conservation and management relevance. Authorship is the mean number (max-mix) and mean proportion of assigned affiliation. Taxonomic breakdown is the number of studies with the focal species in one of the following groups: Aves (A), Amphibia and Reptilia (A + R), Fish (F), Insecta (I), Mammalia (M), Plantae (P), Other (O).

Journal	Authorship breakdown				Taxonomic breakdown						
	Author number (range)	Academic	Government	Non-government	A	A + R	F	I	M	P	O
Conservation Genetics	5 (1 – 13)	70%	18%	13%	9	18	9	3	19	13	4
Biological Conservation	5 (1 – 12)	61%	21%	18%	10	17	5	6	24	13	1
Evolutionary Applications	6 (1–17)	64%	27%	9%	8	4	33	5	12	8	5
Molecular Ecology	5 (2–14)	68%	20%	12%	8	14	13	6	18	11	5




Fig. 1. Taxonomic and conservation breakdown of conservation genetic and genomic studies included in the analysis. The conservation status of 0, 1, or 2, reflect a scale from common to critically endangered and were derived from the IUCN and NatureServe criteria (see Table S1 for more detail). Species that were data deficient or not assessed were given an NA.

Table 2
Logistic regression analysis predicting the specificity of recommendation based on the species global conservation status and proportion of non-academic author affiliation with journal as a random effect ($n = 194$, random effect SD = 0.14). The probability of a baseline study with only academic authors and on a species of least concern status making a specific recommendation was 22%. Studies with only academic authors were 2.8 and 3.4 times more likely to make a conservation recommendation for threatened and critically endangered species, respectively, compared to species of least concern.

Variable	β (SE)	p-Value	Odds ratio (95% CI)
Intercept	− 1.28 (0.28)	< 0.001	
Proportion of non-academic authors	0.91 (0.47)	0.05	2.49 (1.00, 6.24)
Global conservation status			
1 (Vulnerable or threatened)	1.06 (0.35)	0.002	2.88 (1.46, 5.67)
2 (Critically endangered or possibly extinct)	1.21 (0.48)	0.012	3.35 (1.31, 8.57)

conservation concern or species yet to be assessed (Fig. 1). The author composition of generic and specific studies showed a positive effect of including government authors on the conservation recommendation (Table 2; Fig. 2) and there was no relationship between author affiliation and the conservation status of the focal taxa (Table S3).

Focusing on only those studies where we had consistent recommendation scores (i.e. both evaluators had the same recommendation), the proportion of non-academic authors and the global conservation status were significant in the univariate models (p -value < 0.05; Tables S3 and S4). We found no significant difference in the proportion of non-academic authors between the three levels of conservation concern based on a one-way ANOVA ($F(2191) = 1.288$, $p = 0.278$) so both variables were added to the final model (Table 2). Studies that included non-academic affiliated authors increased the odds by 2.5 of the paper making a specific conservation or management recommendation (Table 2).

4. Discussion

The conservation genetics gap reflects the limited integration of genetic data into policy and conservation action. Access to funds and expertise are thought to be among the driving factors (Taylor et al., 2017); however, our meta-analysis sheds some important light on the conservation genetics gap. The majority of conservation genetic and genomic studies we analyzed were on ubiquitous species and offered no specific conservation or management recommendation. This has created a sub-discipline replete with studies of questionable conservation relevance that do not directly inform action or policy. If conservation and management decisions are to be data driven, the scientific literature is critical when it comes to guiding implementation (Game et al., 2015). This is particularly problematic in the case of genomics, as such data cannot be easily applied by members of the conservation community. There is certainly value in studying common, widespread species (e.g. invasive species), but it is problematic when it constitutes the bulk of literature in an applied science like conservation genetics.

Most relevant to bridging the gap is the role of non-academic authors. In studies with generic conservation recommendations for example, the median number of authors with government affiliations was 0, whereas most studies with specific recommendations one or more government author. Consequently, we observed a 2.5-fold increase of recommendations being made in publications when non-academic authors were listed as coauthors (Table 2). This is further reflected in the density plots (Fig. 2), where we see a small, but notable shift towards increased non-academic participation in those articles making specific policy and conservation recommendations. The conservation genetics gap can therefore be partially bridged simply by the inclusion of non-academic collaborators that we surmise have a heightened understanding of policy and on-the-ground needs.

4.1. The conservation genetics gap

The application of genetics has clearly influenced wildlife




Fig. 2. Density plots showing the relationship between proportion of academic and non-academic authors in published articles making a generic or specific conservation and management recommendations.

conservation and management (Nielsen, 1998; Silva et al., 2008; Johnson et al., 2010). Still, a conservation genetics gap has emerged (Hoban et al., 2013a), and a summary of the 2007 international workshop on Population Genetics for Animal Conservation highlighted the field's disconnect from policy and practical conservation solutions (Vernesi et al., 2008). Almost 10 years later, our quantitative analysis shows the gap is still present and largely reiterates the commentary of Vernesi et al. (2008); few conservation genetic studies actually offer explicit ways to use or implement the data and generally disregard policy. Likewise, a recent review on landscape genetics (Bowman et al., 2016) found practitioners applied landscape genetics, and in fact in all genetic fields, less than expected based on the number of publications.

Bowman et al. (2016), like Taylor et al. (2017), suggested that the gap in application might be due to lack of understanding by practitioners, but also hypothesized that studies might be designed in such a way that they have limited policy and management relevance. We suggest an alternative explanation where the demands to publish have resulted in academic authors using the themes of conservation as a means of framing a thesis to fit the publisher's aims and scope. This has resulted in a conservation genetics literature reporting on largely ubiquitous species (Fig. 1) and generic conservation statements (Table 2) and is supported by the fact that the majority of publications with generic recommendations consisted of only academic authors (Fig. 2). Equivalent to the reproducibility crisis facing human and drug research (Prinz et al., 2011), conservation genetics faces an application crisis, where few studies appear to have real-world applications. We argue this has created a perception problem where the conservation community sees a reduced value in the tool because the scientific literature primarily offers examples with tangential applications to conservation.

4.2. The conservation genomics gap

Conservation science produces the data and information that leads to knowledge about biodiversity conservation; in contrast, conservation action is the process of using that information to make decisions that help conserve biodiversity (Sullivan et al., 2017). New technology and data streams received with a lot of attention in the scientific and popular literature, might not be relevant to conservation if they are too complicated or constrained by accessibility and perceived value (Sunderland et al., 2009; Cook et al., 2013). Finding only 16 (~5%) conservation genomic studies in our non-exhaustive literature review provides evidence that a gap exists between high-throughput molecular methods and their utility for conservation practice, despite the plethora of reviews and opinion pieces on the subject (see also Garner et al., 2016). Put another way, the number of instances where whole-genome sequencing is more valuable than aerial surveys or anti-poaching efforts for management and conservation is very small.

In the field of genomics, the quest to identify genes underlying traits has led to a tendency to 'tell stories', meaning trying to make biological sense out of false positives (Pavlidis et al., 2012). Similarly, we argue conservation genomics has fallen into the same trap of telling stories about saving biodiversity while offering few specifics on how to achieve conservation targets. It is important to recognize that more complex data and methods, including the use of gene-editing tools for conservation (see Taylor and Gemmell, 2016), are dramatically less approachable and much higher risk for wildlife managers and conservation practitioners. While increased resolution and novel genomic metrics (e.g. Kardos et al., 2016) are clearly valuable to some systems and questions, there is a need to identify explicit scenarios where genomic tools will have meaningful impact on conservation and management outcomes. Even the notion of improved taxonomic designations is complicated by the amount of noise seen with genomic data (i.e.

incomplete lineage sorting; Nater et al., 2015), as evidenced by the failure to resolve the controversial species status and highly relevant conservation listing of wolves in North America (vonHoldt et al., 2016; Hohenlohe et al., 2017). Touting genomics as a panacea might in fact worsen the conservation genomics gap by creating unrealistic expectations while actually holding limited relevance to stakeholders.

4.3. Bridging the gap

Managers and committees are tasked with using the best available scientific knowledge to make decisions. In this regard, the sheer volume of published genetic and genomic studies touted as impacting conservation and management, but offering no means of how the data can be used, must surely impact one's view of the tool. Faced with a choice of sequencing genomes versus purchasing radio-collars, the applications of the latter are clear and less nuanced than identifying putatively adaptive regions of the genome. Conservation genetics and genomics must refrain from hyperbole and focus on identifying clear and readily transferable applications, which might not align with academic funding and publishing models (Shafer et al., 2015). This builds on our most important finding in that increasing non-academic participation in research projects resulted in more studies offering specific conservation and management recommendations. Thus, research teams interested in achieving conservation action and actually informing policy should seek out non-academic collaborations; likewise, conservation-oriented journals should ensure that the genetic or genomic relevance to conservation practice or policy is clearly stated. Vernesi et al. (2008) concluded their assessment of conservation genetics by arguing for closer collaboration between theoreticians and field biologists; we argue for the same but between academics and non-academics. Otherwise the field of conservation genetics and conservation genomics are at risk of becoming only an academic exercise.

Acknowledgements

We thank the research group of Dr. Christopher Kyle for providing feedback to our research query. We thank Dr. Joe Northrup (Ontario Ministry of Natural Resources) and Kevin White (Alaska Department of Fish & Game) for providing feedback on this manuscript. Trent University start-up funds and an NSERC Discovery Grant (RGPIN-2017-03934) to ABAS provided financial support. MB is supported by an NSERC-USRA, and DM an Ontario Graduate Scholarship. We thank our reviewers, two anonymous and Eric Archer (NOAA Southwest Fisheries Science Center), for their useful feedback on this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.biocon.2017.12.019>.

References

- Amos, W., Balmford, A., 2001. When does conservation genetics matter? *Heredity* 87, 257–265.
- Bowman, J., Greenhorn, J.E., Marrotte, R.R., McKay, M.M., Morris, K.Y., Prentice, M.B., Wehtje, M., 2016. On applications of landscape genetics. *Conserv. Genet.* 1–10.
- Carter, M.F., Hunter, W.C., Pashley, D.N., Rosenberg, K.V., 2000. Conservation report: setting conservation priorities for landbirds in the United States: the partners in flight approach. *Auk* 117, 541–548.
- Cook, C.N., Mascia, M.B., Schwartz, M.W., Possingham, H.P., Fuller, R.A., 2013. Achieving conservation science that bridges the knowledge-action boundary. *Conserv. Biol. J. Soc. Conserv. Biol.* 27, 669–678.
- Cronin, M.A., 2007. Limitations of molecular genetics in conservation. *Nature* 447 (7145), 638.
- Frankham, R., 2003. Genetics and conservation biology. *C. R. Biol.* 326, 22–29.
- Fuentes-Pardo, A.P. & Ruzzante, D.E. (n.d.). Whole-genome sequencing approaches for conservation biology: advantages, limitations, and practical recommendations. *Mol. Ecol.*, (n/a–n/a).
- Game, E.T., Schwartz, M.W., Knight, A.T., 2015. Policy relevant conservation science. *Conserv. Lett.* 8, 309–311.
- Garner, B.A., Hand, B.K., Amish, S.J., Bernatchez, L., Foster, J.T., Miller, K.M., Morin, P.A., Narum, S.R., O'Brien, S.J., Roffler, G., Templin, W.D., Sunnucks, P., Strait, J., Warheit, K.L., Seamons, T.R., Wenburg, J., Olsen, J., Luikart, G., 2016. Genomics in conservation: case studies and bridging the gap between data and application. *Trends Ecol. Evol.* 31, 81–83.
- Groves, C., 2003. Drafting a Conservation Blueprint: A practitioner's Guide to Planning for Biodiversity. Island Press.
- Groves, C.R., Jensen, D.B., Valutis, L.L., Redford, K.H., Shaffer, M.L., Scott, J.M., Baumgartner, J.V., Higgins, J.V., Beck, M.W., Anderson, M.G., 2002. Planning for biodiversity conservation: putting conservation science into practice seven-step framework for developing regional plans to conserve biological diversity, based upon principles of conservation biology and ecology, is being used extensively by the nature conservancy to identify priority areas for conservation. *Bioscience* 52, 499–512.
- Haig, S.M., Miller, M.P., Bellinger, M.R., Draheim, H.M., Mercer, D.M., Mullins, T.D., 2015. The Conservation Genetics Juggling Act: integrating genetics and ecology, science and policy. *Evol. Appl.* 9, 181–195.
- Hoban, S., Arntzen, J.W., Bertorelle, G., Bryja, J., Fernandes, M., Frith, K., Gaggiotti, O., Galbusera, P., Godoy, J.A., Hauffe, H.C., Rus Hoelzel, A., Nichols, R.A., Pérez-Espona, S., Primmer, C., Russo, I.-R.M., Segelbacher, G., Siegmund, H.R., Sihvonen, M., Sjögren-Gulve, P., Vernesi, C., Vilà, C., Bruford, M.W., 2013a. Conservation genetic resources for effective species survival (congress): bridging the divide between conservation research and practice. *J. Nat. Conserv.* 21, 433–437.
- Hoban, S.M., Hauffe, H.C., Pérez-Espona, S., Arntzen, J.W., Bertorelle, G., Bryja, J., Frith, K., Gaggiotti, O.E., Galbusera, P., Godoy, J.A., Hoelzel, A.R., Nichols, R.A., Primmer, C.R., Russo, I.-R., Segelbacher, G., Siegmund, H.R., Sihvonen, M., Vernesi, C., Vilà, C., Bruford, M.W., 2013b. Bringing genetic diversity to the forefront of conservation policy and management. *Conserv. Genet. Resour.* 5, 593–598.
- Hohenlohe, P.A., Rutledge, L.Y., Waits, L.P., Andrews, K.R., Adams, J.R., Hinton, J.W., Nowak, R.M., Patterson, B.R., Wydeven, A.P., Wilson, P.A., White, B.N., 2017. Comment on “Whole-genome sequence analysis shows two endemic species of North American wolf are admixtures of the coyote and gray wolf.”. *Sci. Adv.* 3, e1602250.
- vonHoldt, B.M., Cahill, J.A., Fan, Z., Gronau, I., Robinson, J., Pollinger, J.P., Shapiro, B., Wall, J., Wayne, R.K., 2016. Whole-genome sequence analysis shows that two endemic species of North American wolf are admixtures of the coyote and gray wolf. *Sci. Adv.* 2, e1501714.
- Johnson, W.E., Onorato, D.P., Roelke, M.E., Land, E.D., Cunningham, M., Belden, R.C., McBride, R., Jansen, D., Lotz, M., Shindle, D., Howard, J., Wildt, D.E., Penfold, L.M., Hostetler, J.A., Oli, M.K., O'Brien, S.J., 2010. Genetic restoration of the Florida Panther. *Science* 329, 1641–1645.
- Kardos, M., Taylor, H.R., Ellegren, H., Luikart, G., Allendorf, F.W., 2016. Genomics advances the study of inbreeding depression in the wild. *Evol. Appl.* 9, 1205–1218.
- Knight, A.T., Driver, A., Cowling, R.M., Maze, K., Desmet, P.G., Lombard, A.T., Rouget, M., Botha, M.A., Boshoff, A.F., Castley, J.G., Goodman, P.S., Mackinnon, K., Pierce, S.M., Sims-Castley, R., Stewart, W.I., von Hase, A., 2006. Designing systematic conservation assessments that promote effective implementation: best practice from South Africa. *Conserv. Biol. J. Soc. Conserv. Biol.* 20, 739–750.
- McMahon, B.J., Teeling, E.C., Höglund, J., 2014. How and why should we implement genomics into conservation? *Evol. Appl.* 7, 999–1007.
- Nagulendran, K., Padfield, R., Aziz, S.A., Amir, A.A., Rahman, A.R.A., Latiff, M.A., Zafir, A., Quilter, A.G., Tan, A., Arifah, S., Awang, N., Azhar, N., Balu, P., Gan, P.C., Hii, N., Reza, M.I.H., Lavanya, R.I.L., Lim, T., Mahendra, S., Rayan, D.M., McGowan, S., Paxton, M., Mohamed, Z., Salleh, D.M., Abdullah, M.T., Ibrahim, N.A.N., Puan, C.L., Clements, G.R., Mohamed, I.S.M., Saw, L.G., Shashi, K., Sivananthan, E., Sharma, D.S.K., Surin, S., Vanitha, P., Wadey, J., Hasmah, W.M.W., Wong, E.P., Wong, P.M., Yeap, C.A., Campos-Arceiz, A., 2016. A multi-stakeholder strategy to identify conservation priorities in Peninsular Malaysia. *Cogent Environ. Sci.* 2, 1254078.
- Nater, A., Burri, R., Kawakami, T., Smeds, L., Ellegren, H., 2015. Resolving evolutionary relationships in closely related species with whole-genome sequencing data. *Syst. Biol.* 64, syv045–1017.
- Nielsen, J.L., 1998. Population genetics and the conservation and management of Atlantic salmon (*Salmo salar*). *Can. J. Fish. Aquat. Sci.* 55, 145–152.
- Ouborg, N.J., 2010. Integrating population genetics and conservation biology in the era of genomics. *Biol. Lett.* 6, 3–6.
- Ouborg, N.J., Angeloni, F., Vergeer, P., 2010. An essay on the necessity and feasibility of conservation genomics. *Conserv. Genet.* 11, 643–653.
- Pavlidis, P., Jensen, J.D., Stephan, W., Stamatakis, A., 2012. A critical assessment of storytelling: gene ontology categories and the importance of validating genomic scans. *Mol. Biol. Evol.* 29, 3237–3248.
- Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N., Raven, P.H., Roberts, C.M., Sexton, J.O., 2014. The biodiversity of species and their rates of extinction, distribution, and protection. *Science* 344, 1246752.
- Prinz, F., Schlange, T., Asadullah, K., 2011. Believe it or not: how much can we rely on published data on potential drug targets? *Nat. Rev. Drug Discov.* 10, 712.
- Shafer, A.B.A., Wolf, J.B.W., Alves, P.C., Bergström, L., Bruford, M.W., Brännström, I., Colling, G., Dalén, L., De Meester, L., Ekblom, R., Fawcett, K.D., Fior, S., Hajibabaei, M., Hill, J.A., Hoezel, A.R., Höglund, J., Jensen, E.L., Krause, J., Kristensen, T.N., Krützen, M., McKay, J.K., Norman, A.J., Ogden, R., Österling, E.M., Ouborg, N.J., Piccolo, J., Popović, D., Primmer, C.R., Reed, F.A., Roumet, M., Salmons, J., Schenekar, T., Schwartz, M.K., Segelbacher, G., Senn, H., Thaulow, J., Valtonen, M., Veale, A., Vergeer, P., Vijay, N., Vilà, C., Weissensteiner, M., Wennerström, L., Wheat, C.W., Zieliński, P., 2015. Genomics and the challenging translation into conservation practice. *Trends Ecol. Evol.* 30, 78–87.
- Shafer, A.B.A., Wolf, J.B.W., Alves, P.C., Bergström, L., Colling, G., Dalén, L., De Meester, L., Ekblom, R., Fior, S., Hajibabaei, M., Hoezel, A.R., Höglund, J., Jensen, E.L., Krützen, M., Norman, A.J., Österling, E.M., Ouborg, N.J., Piccolo, J., Primmer, C.R.,

- Reed, F.A., Roumet, M., Salmons, J., Schwartz, M.K., Segelbacher, G., Thaulow, J., Valtonen, M., Vergeer, P., Weissensteiner, M., Wheat, C.W., Vilà, C., Zieliński, P., 2016. Reply to Garner et al. *Trends Ecol. Evol.* 31, 83–84.
- Silva, M.B., Kanashiro, M., Ciampi, A.Y., Thompson, I., Sebbenn, A.M., 2008. Genetic effects of selective logging and pollen gene flow in a low-density population of the dioecious tropical tree *Bagassa guianensis* in the Brazilian Amazon. *For. Ecol. Manag.* 255, 1548–1558.
- Soulé, M.E., 1985. What is conservation biology? *Bioscience* 35, 727–734.
- Sullivan, B.L., Phillips, T., Dayer, A.A., Wood, C.L., Farnsworth, A., Iliff, M.J., Davies, I.J., Wiggins, A., Fink, D., Hochachka, W.M., Rodewald, A.D., Rosenberg, K.V., Bonney, R., Kelling, S., 2017. Using open access observational data for conservation action: a case study for birds. *Biol. Conserv.* 208, 5–14.
- Sunderland, T., Sunderland-Groves, J., Shanley, P., Campbell, B., 2009. Bridging the gap: how can information access and exchange between conservation biologists and field practitioners be improved for better conservation outcomes? *Biotropica* 41, 549–554.
- Sutherland, W.J., Adams, W.M., Aronson, R.B., Aveling, R., Blackburn, T.M., Broad, S., Ceballos, G., Côté, I.M., Cowling, R.M., Da Fonseca, G.A.B., Dinerstein, E., Ferraro, P.J., Fleishman, E., Gascon, C., Hunter, M., Hutton, J., Kareiva, P., Kuria, A., Macdonald, D.W., Mackinnon, K., Madgwick, F.J., Mascia, M.B., McNeely, J., Milner-Gulland, E.J., Moon, S., Morley, C.G., Nelson, S., Osborn, D., Pai, M., Parsons, E.C.M., Peck, L.S., Possingham, H., Prior, S.V., Pullin, A.S., Rands, M.R.W., Ranganathan, J., Redford, K.H., Rodriguez, J.P., Seymour, F., Sobel, J., Sodhi, N.S., Stott, A., Vance-Borland, K., Watkinson, A.R., 2009. One hundred questions of importance to the conservation of global biological diversity. *Conserv. Biol. J. Soc. Conserv. Biol.* 23, 557–567.
- Taylor, H.R., Gemmell, N.J., 2016. Emerging technologies to conserve biodiversity: further opportunities via genomics. Response to Pimm et al. *Trends Ecol. Evol.* 31, 171–172.
- Taylor, R.H., Dussex, N., van Heezik, Y., 2017. Bridging the conservation genetics gap by identifying barriers to implementation for conservation practitioners. *Glob. Ecol. Conserv.* 10, 231–242.
- Vernesi, C., Bruford, M.W., Bertorelle, G., Pecchioli, E., Rizzoli, A., Hauffe, H.C., 2008. Where's the conservation in conservation genetics? *Conserv. Biol.* 22, 802–804.