

# Math 100A: Homework 8

## Merrick Qiu

### Problem 1

Since the trace is the sum of the diagonal entries and each diagonal entry is the dot product of the associated row in  $X$  and column in  $Y$ , then

$$\begin{aligned}\operatorname{tr}(XY) &= \sum_{i=1}^n \sum_{j=1}^n X_{i,j} Y_{j,i} \\ &= \sum_{j=1}^n \sum_{i=1}^n Y_{j,i} X_{i,j} \\ &= \operatorname{tr}(YX)\end{aligned}$$

As a corollary, we have that

$$\operatorname{tr}(SXS^{-1}) = \operatorname{tr}(SS^{-1}X) = \operatorname{tr}(X)$$

## Problem 2

(  $\implies$  ) Suppose there existed nonzero  $w \in V$  such that  $\mathbf{C}[G](w) = W \neq V$ .  $W \neq \{0\}$  since  $W = \{0\}$  would imply  $\rho(g)(w) = 0$  for all  $g$ . Since  $\rho(g)$  is invertible, that implies  $w = 0$  which is a contradiction.

$W$  is a nontrivial  $G$ -invariant subspace since it is the span of  $g \cdot w$  for all  $g \in G$ . Applying a group action  $g$  to an element in that span will still yield an element in that span. Therefore if  $V$  is irreducible it must be that  $\mathbf{C}[G](v) = V$  for all nonzero  $v \in V$ .

(  $\impliedby$  ) If  $W \subseteq V$  was a nontrivial  $G$ -invariant subspace, then  $\rho(g)(w) \in W$  for all  $g$ , and so  $\mathbf{C}[G](w) \subseteq W$ . However this contradicts the fact that  $\mathbf{C}[G](v) = V$  for all  $v \neq 0$  in  $V$ . Therefore  $V$  does not contain any  $G$ -invariant subspaces other than  $\{0\}$  and  $V$ .

### Problem 3

Let  $W = \{(v_1, \dots, v_n) : v_1 + \dots + v_n = 0\}$ . Suppose  $w = (w_1, \dots, w_n) \in W$  is not zero. It is sufficient to show that  $\mathbf{C}[S_n](w) = W$  to show that  $W$  is irreducible by the previous question. There exists  $i < j$  so that  $w_i \neq w_j$  since  $w \neq 0$ . Let  $\tau$  be the permutation of  $S_n$  that only exchanges  $i$  with  $j$ .

Notice that  $\tau \cdot w - w \in \mathbf{C}[S_n](w)$  (since it is a linear combination of permutations of  $w$ ) and that  $\tau \cdot w - w = \alpha(e_i - e_j)$  for some  $\alpha \neq 0$ . Let  $\sigma_k$  be the permutation that exchanges  $j$  and  $n$  and then exchanges  $i$  with  $k$ . Then  $\sigma_k(\tau \cdot w - w) = \alpha(e_k - e_n) \in \mathbf{C}[S_n](w)$  too. Since any  $w \in W$  can be written as a linear combination of the  $e_k - e_n$  terms, and each  $e_k - e_n$  is in  $\mathbf{C}[S_n](w)$ , we have that  $\mathbf{C}[S_n](w) = W$ .