

# Math 120A: Homework 9

## Merrick Qiu

### 1. Page 159: Problem 1(a)(b)

$$f(z) = \frac{z^2}{z+3}$$

Since this function is the quotient of two polynomials, it is only not analytic at  $z = -3$  which is outside of the region bound by the contour. Therefore the function is analytic everywhere interior to and on  $C$  so the integral is zero by the Cauchy-Goursat theorem.

$$f(z) = ze^{-z}$$

Since  $z$  is analytic and  $e^{-z}$  is analytic, their product is analytic everywhere. Therefore the integral is zero by the Cauchy-Goursat theorem.

## 2. Page 159: Problem 2(a)

$$f(z) = \frac{1}{3z^2 + 1}$$

Since this function is the quotient of two polynomials, it is analytic everywhere except when  $3z^2 + 1 = 0$ , where  $z = \pm \frac{1}{\sqrt{3}}i$ , which is inside of the square and outside of the region bound by the circle and the square. Therefore by the corollary, the two integrals are equal.

### 3. Page 170: Problem 1(a)(b)

Since  $e^{-z}$  is analytic, by the cauchy integral formula

$$\int_C \frac{e^{-z} dz}{z - (\pi i/2)} = 2\pi i e^{-\pi i/2} = 2\pi$$

Since  $\frac{\cos z}{z^2+8}$  is analytic interior to  $C$ ,

$$\int_C \frac{\cos z}{z(z^2+8)} dz = 2\pi i \frac{\cos 0}{0^2+8} = \frac{2\pi i}{8}$$

#### 4. Page 170: Problem 2

Since  $\frac{1}{z+2i}$  is analytic inside the circle,

$$\int_C \frac{1}{(z-2i)(z+2i)} dz = 2\pi i \frac{1}{2i+2i} = \frac{\pi}{2}$$

By the cauchy extension,

$$\int_C \frac{1}{(z-2i)^2(z+2i)^2} dz = \frac{2\pi i}{1!} \left( -\frac{2}{(2i+2i)^3} \right) = \frac{\pi}{16}$$

## 5. Page 170: Problem 3

Since  $2s^2 - s - 2$  is analytic everywhere,

$$g(2) = \int_C \frac{2s^2 - s - 2}{s - 2} ds = 2\pi i(2(2)^2 - 2 - 2) = 8\pi i$$

If  $|z| > 3$  the cauchy-goursat theorem says that  $g(z) = 0$ .

## 6. Page 170: Problem 4

If  $z$  is inside then by the cauchy extension

$$\frac{d^2}{ds^2}s^3 + 2s = \frac{d}{ds}3s^2 + 2 = 6s$$

$$\begin{aligned}\int_C \frac{s^3 + 2s}{(s - z)^3} ds &= \frac{2\pi i}{2!} (6z) \\ &= 6\pi iz\end{aligned}$$

If  $z$  is outside then the cauchy-goursat theorem says that  $g(z) = 0$ .