

MATH 31AH - Homework 6

Merrick Qiu

November 11, 2021

1 Eigenvalues and Eigenvectors

A has eigenvalues when $\det(A - \lambda I) = 0$.

$$A - \lambda I = \begin{bmatrix} 7 - \lambda & -8 & 6 \\ 8 & -9 - \lambda & 6 \\ 0 & 0 & 1 - \lambda \end{bmatrix}$$

The characteristic polynomial of A is thus

$$(7 - \lambda)(-9 - \lambda)(1 - \lambda) - (-8)(8)(1 - \lambda) = (-63 + 2\lambda + \lambda^2)(1 - \lambda) + 64(1 - \lambda) \quad (1)$$

$$= (1 + 2\lambda + \lambda^2)(1 - \lambda) \quad (2)$$

$$= (1 + \lambda)^2(1 - \lambda) \quad (3)$$

$(1 + \lambda)^2(1 - \lambda) = 0$ when $\lambda = -1, 1$, so those are our eigenvalues. Using these eigenvalues, we can calculate the matrices for the eigenspaces.

$$E_{-1} = \begin{bmatrix} 8 & -8 & 6 \\ 8 & -8 & 6 \\ 0 & 0 & 2 \end{bmatrix} \text{ and } E_1 = \begin{bmatrix} 6 & -8 & 6 \\ 8 & -10 & 6 \\ 0 & 0 & 0 \end{bmatrix}$$

Using RREF, we have that

$$\text{rref}(E_{-1}) = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \text{ and } \text{rref}(E_1) = \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore a basis for E_{-1} is $\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$ and a basis for E_1 is $\left\{ \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix} \right\}$. Since there are not three eigenvalues, A is not diagonalizable.

2 More Eigenvalues and Eigenvectors

B has eigenvalues when $\det(B - \lambda I) = 0$.

$$B - \lambda I = \begin{bmatrix} 1 - \lambda & \frac{1}{4} & 0 \\ 0 & \frac{1}{2} - \lambda & 0 \\ 0 & \frac{1}{4} & -1 - \lambda \end{bmatrix}$$

The characteristic polynomial of B is thus $(1 - \lambda)(\frac{1}{2} - \lambda)(-1 - \lambda)$. The eigenvalues are thus $\lambda = -1, \frac{1}{2}, 1$. Using these eigenvalues, we can calculate the matrices for the eigenspaces.

$$E_{-1} = \begin{bmatrix} 2 & \frac{1}{4} & 0 \\ 0 & \frac{3}{2} & 0 \\ 0 & \frac{1}{4} & 0 \end{bmatrix} \text{ and } E_{\frac{1}{2}} = \begin{bmatrix} \frac{1}{2} & \frac{1}{4} & 0 \\ 0 & 0 & 0 \\ 0 & \frac{1}{4} & -\frac{3}{2} \end{bmatrix} \text{ and } E_1 = \begin{bmatrix} 0 & \frac{1}{4} & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & \frac{1}{4} & -2 \end{bmatrix}$$

Using RREF, we have that

$$\text{rref}(E_{-1}) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \text{ and } \text{rref}(E_{\frac{1}{2}}) = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & -6 \\ 0 & 0 & 0 \end{bmatrix} \text{ and } \text{rref}(E_1) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore a basis for E_{-1} is $\left\{ \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$, a basis for $E_{\frac{1}{2}}$ is $\left\{ \begin{bmatrix} -3 \\ 6 \\ 1 \end{bmatrix} \right\}$, and a basis for E_1 is $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$. Since there are three eigenvalues, B is diagonalizable.

3 \mathbb{R} and \mathbb{C} and diagonalizability

The matrix has eigenvalues when $\det(R_\theta - \lambda I) = 0$. We have that the characteristic polynomial is

$$(\cos(\theta) - \lambda)^2 + \sin^2(\theta) = \cos^2(\theta) - 2\lambda + \lambda^2 + \sin^2(\theta) \quad (4)$$

$$= 1 - 2\lambda \cos(\theta) + \lambda^2 \quad (5)$$

Using the quadratic formula, we get that our eigenvalues are

$$\lambda = \frac{2 \cos(\theta) \pm \sqrt{4 \cos^2(\theta) - 4}}{2} \quad (6)$$

$$= \cos(\theta) \pm \sqrt{\cos^2(\theta) - 1} \quad (7)$$

$$= \cos(\theta) \pm i \sin(\theta) \quad (8)$$

$$= e^{\pm i\theta} \quad (9)$$

Our eigenspace matrices are

$$E_{e^{-i\theta}} = \begin{bmatrix} i \sin(\theta) & -\sin(\theta) \\ \sin(\theta) & i \sin(\theta) \end{bmatrix}$$

$$E_{e^{i\theta}} = \begin{bmatrix} -i \sin(\theta) & -\sin(\theta) \\ \sin(\theta) & -i \sin(\theta) \end{bmatrix}$$

So the corresponding eigenvectors are $v_{e^{-i\theta}} = (-i, 1)$ and $v_{e^{i\theta}} = (i, 1)$. Over \mathbb{R} , R_θ is only diagonalizable when $\theta = 0, \pi$ since these are the only two angles that result in real eigenvalues. Over \mathbb{C} , R_θ is diagonalizable for all θ because there are always two eigenvalues.

4 Polynomials and diagonalizability

Proof. $f(A)$ is diagonalizable when A is diagonalizable.

Base Case: Since A is diagonalizable, let P be a matrix such that $A = PDP^{-1}$ for some D . A^n can thus be written as $A = PD^nP^{-1}$.¹ Therefore $c_nA^n = P(c_nD^n)P^{-1}$, so each of the terms in $f(A)$ is diagonalizable, including when $n = 0$.

Inductive Step: For $n > 0$, assume that the sum of the first $n - 1$ terms is diagonalizable with $f_{n-1}(A) = PBP^{-1}$ for some matrix B . Since the n th term can be written as $c_nA^n = P(c_nD^n)P^{-1}$, the sum of the first n terms is

$$f_{n-1}(A) + c_nA^n = PBP^{-1} + P(c_nD^n)P^{-1} \quad (10)$$

$$= P(BP^{-1} + c_nD^nP^{-1}) \quad (11)$$

$$= P(B + c_nD^n)P^{-1} \quad (12)$$

This completes the inductive step, and so $f(A)$ is diagonalizable when A is diagonalizable. \square

5 Direct sums and diagonalizability

Proof. If λ is an eigenvalue of T or U , then λ is an eigenvalue of $T \oplus U$

If λ is an eigenvalue of T and $v \in T$ is the corresponding eigenvector, then the corresponding eigenvector in $T \oplus U$ with eigenvalue λ would be $(v, 0)$ since

$$(T \oplus U)(v, 0) = (T(v), U(0)) \quad (13)$$

$$= (\lambda v, 0) \quad (14)$$

$$= \lambda(v, 0) \quad (15)$$

Similarly, if λ is an eigenvalue of U and $w \in U$ is the corresponding eigenvector, then the corresponding eigenvector in $T \oplus U$ with eigenvalue λ would be $(0, w)$ since

$$(T \oplus U)(0, w) = (T(0), U(w)) \quad (16)$$

$$= (0, \lambda w) \quad (17)$$

$$= \lambda(0, w) \quad (18)$$

Therefore, if λ is an eigenvalue of T or U , then λ is an eigenvalue of $T \oplus U$ \square

Proof. If T and U are diagonalizable, then $T \oplus U$ is diagonalizable.

Let V be m dimensional and let W be n dimensional. Since T and U are both diagonalizable, they have m and n eigenvalues respectively.

Since every eigenvalue of T or U is an eigenvalue of $T \oplus U$, $T \oplus U$ has $m + n$ eigenvalues. Since $T \oplus U$ has $m + n$ dimensions and $m + n$ eigenvalues, $T \oplus U$ is diagonalizable. \square

¹See problem 7 for proof

6 A 6x6 example

We have that

$$A - \lambda I = \begin{bmatrix} -\lambda & 0 & 0 & 0 & 0 & 1 \\ 1 & -\lambda & 0 & 0 & 0 & 0 \\ 0 & 1 & -\lambda & 0 & 0 & 0 \\ 0 & 0 & 1 & -\lambda & 0 & 0 \\ 0 & 0 & 0 & 1 & -\lambda & 0 \\ 0 & 0 & 0 & 0 & 1 & -\lambda \end{bmatrix}$$

There are only two possible permutations, or "rook placements" that result in non-negative terms, so

$$\det(A - \lambda I) = (-\lambda)^6 - 1^6 \quad (19)$$

$$= \lambda^6 - 1 \quad (20)$$

$\lambda^6 - 1 = 0$ when $\lambda = 1, e^{i\frac{\pi}{3}}, e^{i\frac{2\pi}{3}}, -1, e^{i\frac{4\pi}{3}}, e^{i\frac{5\pi}{3}}$, the 6 roots of unity. The row reduced matrix of $A - \lambda I$ is

$$\begin{bmatrix} -\lambda & 0 & 0 & 0 & 0 & 1 \\ 0 & -\lambda & 0 & 0 & 0 & \frac{1}{\lambda} \\ 0 & 0 & -\lambda & 0 & 0 & \frac{1}{\lambda^2} \\ 0 & 0 & 0 & -\lambda & 0 & \frac{1}{\lambda^3} \\ 0 & 0 & 0 & 0 & -\lambda & \frac{1}{\lambda^4} \\ 0 & 0 & 0 & 0 & 0 & \frac{-\lambda^6+1}{\lambda^5} \end{bmatrix}$$

Since $\lambda^6 - 1 = 0$, we have that $\frac{-\lambda^6+1}{\lambda^5} = 0$. The rref of $A - \lambda I$ is thus

$$\text{rref}(A - \lambda I) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & -\frac{1}{\lambda} \\ 0 & 1 & 0 & 0 & 0 & -\frac{1}{\lambda^2} \\ 0 & 0 & 1 & 0 & 0 & -\frac{1}{\lambda^3} \\ 0 & 0 & 0 & 1 & 0 & -\frac{1}{\lambda^4} \\ 0 & 0 & 0 & 0 & 1 & -\frac{1}{\lambda^5} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Using the rref, we can determine the eigenvectors of the eigenspace.

$$v_1 = \left\{ \left[\frac{1}{1}, \frac{1}{(1)^2}, \frac{1}{(1)^3}, \frac{1}{(1)^4}, \frac{1}{(1)^5}, 1 \right] \right\} \quad (21)$$

$$v_{e^{i\frac{\pi}{3}}} = \left\{ \left[\frac{1}{e^{i\frac{\pi}{3}}}, \frac{1}{(e^{i\frac{\pi}{3}})^2}, \frac{1}{(e^{i\frac{\pi}{3}})^3}, \frac{1}{(e^{i\frac{\pi}{3}})^4}, \frac{1}{(e^{i\frac{\pi}{3}})^5}, 1 \right] \right\} \quad (22)$$

$$v_{e^{i\frac{2\pi}{3}}} = \left\{ \left[\frac{1}{e^{i\frac{2\pi}{3}}}, \frac{1}{(e^{i\frac{2\pi}{3}})^2}, \frac{1}{(e^{i\frac{2\pi}{3}})^3}, \frac{1}{(e^{i\frac{2\pi}{3}})^4}, \frac{1}{(e^{i\frac{2\pi}{3}})^5}, 1 \right] \right\} \quad (23)$$

$$v_{-1} = \left\{ \left[\frac{1}{-1}, \frac{1}{(-1)^2}, \frac{1}{(-1)^3}, \frac{1}{(-1)^4}, \frac{1}{(-1)^5}, 1 \right] \right\} \quad (24)$$

$$v_{e^{i\frac{4\pi}{3}}} = \left\{ \left[\frac{1}{e^{i\frac{4\pi}{3}}}, \frac{1}{(e^{i\frac{4\pi}{3}})^2}, \frac{1}{(e^{i\frac{4\pi}{3}})^3}, \frac{1}{(e^{i\frac{4\pi}{3}})^4}, \frac{1}{(e^{i\frac{4\pi}{3}})^5}, 1 \right] \right\} \quad (25)$$

$$v_{e^{i\frac{5\pi}{3}}} = \left\{ \left[\frac{1}{e^{i\frac{5\pi}{3}}}, \frac{1}{(e^{i\frac{5\pi}{3}})^2}, \frac{1}{(e^{i\frac{5\pi}{3}})^3}, \frac{1}{(e^{i\frac{5\pi}{3}})^4}, \frac{1}{(e^{i\frac{5\pi}{3}})^5}, 1 \right] \right\} \quad (26)$$

7 A real sequence

Proof. $A^n = PD^nP^{-1}$ if $A = PDP^{-1}$.

We will induct on n . For $n = 1$, we have that $A^1 = PD^1P^{-1}$. For $n > 1$, assume that $A^{n-1} = PD^{n-1}P^{-1}$. We have that

$$A^n = A^{n-1}A \quad (27)$$

$$= (PD^{n-1}P^{-1})(PDP^{-1}) \quad (28)$$

$$= PD^nP^{-1} \quad (29)$$

This completes the inductive step, so $A^n = PD^nP^{-1}$. \square

Proof. When $a_1 = a_2 = 1$ and $a_n = 4a_{n-1} - 2a_{n-2}$, $a_n = \frac{(2+\sqrt{2})^{n-2} + (2-\sqrt{2})^{n-2}}{2}$ for $n > 2$.

The n th number in the sequence can be found using the following equation:

$$\begin{bmatrix} a_n \\ a_{n-1} \end{bmatrix} = \begin{bmatrix} 4 & -2 \\ 1 & 0 \end{bmatrix}^{n-2} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The matrix has characteristic polynomial of $(4-\lambda)(-\lambda)+2 = \lambda^2 - 4\lambda + 2$. Using the quadratic formula, we can find that the matrix has eigenvalues of $\lambda_1 = 2+\sqrt{2}$, $\lambda_2 = 2-\sqrt{2}$. The eigenvectors for these eigenvalues are $v_1 = (2+\sqrt{2}, 1)$ and $v_2 = (2-\sqrt{2}, 1)$ since they solve the corresponding homologous system of equations for the eigenspace.

Since there are two eigenvectors, we can diagonalize the matrix and simplify the expression to

$$\begin{bmatrix} a_n \\ a_{n-1} \end{bmatrix} = \begin{bmatrix} 2+\sqrt{2} & 2-\sqrt{2} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2+\sqrt{2} & 0 \\ 0 & 2-\sqrt{2} \end{bmatrix}^{n-2} \begin{bmatrix} \frac{1}{2\sqrt{2}} & \frac{\sqrt{2}-2}{2\sqrt{2}} \\ -\frac{1}{2\sqrt{2}} & \frac{\sqrt{2}+2}{2\sqrt{2}} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad (30)$$

$$= \frac{1}{2\sqrt{2}} \begin{bmatrix} 2+\sqrt{2} & 2-\sqrt{2} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} (2+\sqrt{2})^{n-2} & 0 \\ 0 & (2-\sqrt{2})^{n-2} \end{bmatrix} \begin{bmatrix} \sqrt{2}-1 \\ \sqrt{2}+1 \end{bmatrix} \quad (31)$$

$$= \frac{1}{2\sqrt{2}} \begin{bmatrix} 2+\sqrt{2} & 2-\sqrt{2} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} (\sqrt{2}-1)(2+\sqrt{2})^{n-2} \\ (\sqrt{2}+1)(2-\sqrt{2})^{n-2} \end{bmatrix} \quad (32)$$

$$= \frac{1}{2\sqrt{2}} \begin{bmatrix} \sqrt{2}(2+\sqrt{2})^{n-2} + \sqrt{2}(2-\sqrt{2})^{n-2} \\ (\sqrt{2}-1)(2+\sqrt{2})^{n-2} + (\sqrt{2}+1)(2-\sqrt{2})^{n-2} \end{bmatrix} \quad (33)$$

$$(34)$$

Therefore, $a_n = \frac{(2+\sqrt{2})^{n-2} + (2-\sqrt{2})^{n-2}}{2}$ for $n > 2$. \square

8 Commuting operators and eigenspaces

Proof. E_λ is U -invariant when T and U are commutative.

Let $T, U : V \rightarrow V$ be two linear transformations that commute. Let $v \in E_\lambda$ be an eigenvector. Since it is a T -eigenvector, we have that $T(v) = \lambda(v)$. Since T and U commute, we have that

$$U(T(v)) = U(\lambda v) \quad (35)$$

$$= \lambda U(v) \quad (36)$$

$$= T(U(v)) \quad (37)$$

Since $T(U(v)) = \lambda U(v)$, we know that $U(v) \in E_\lambda$, and so E_λ is U -invariant. \square

Proof. E_λ is not necessarily U -invariant when T and U are not commutative.

Let $T, U : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that T projects the vector onto the x-axis and U rotates the vector by $\pi/2$. E_λ would clearly not be U -invariant since the only eigenspace of T would be the x-axis, but U rotates vectors off of the x-axis. \square