H5N1 spatial risk: Australia

Technical report

Meryl Theng (on behalf of HASTE)

2026-10-30

1 Introduction

As part of the WOAH Datathon, we built an analytical workflow to map the risk of HPAI H5N1 (clade 2.3.4.4b) in a country of interest. We start with Australia, where H5N1 was first detected in wildlife in June 2026.

Datathon Challenge 1 deliverable: an integrated analytical workflow that produces early warning signals for specific diseases and provides corrected, real-time estimates of disease dynamics and short-term forecasts of disease evolution, with the level of uncertainty being explicitly quantified.

Aim: to estimate how far H5N1 spreads between areas of Australia, using outbreak records, and to combine that estimate with waterbird abundance to produce a risk surface for H5N1 in wild birds. Combining it with poultry density gives an indicative surface for spillover risk to domestic flocks.

How the aim maps onto the deliverable

- *Real-time estimates:* the workflow is fully reproducible (Snakemake), so the kernel and maps can be refitted to updated WAHIS and DAFF (Australian Department of Agriculture Fisheries and Forestry) outbreak data.
- *Correction for detection lag:* each infected area is treated as infectious from a set number of days *before* it was detected. This lag is assumed, not estimated.
- *Uncertainty:* kernel parameters are estimated with approximate confidence intervals, and the fitted curves are shown with uncertainty bands. Sensitivity to the assumed values is discussed in Section 5.

Two kinds of output. The distance kernel fitted to the outbreak data produces a statistical estimate of decay of transmission rate with increasing distance. The risk surfaces are not: they combine the kernel with covariates by fixed, assumed rules and give a *relative ranking* of cells.

The analysis has three parts:

1. fit a distance kernel to the outbreak data (Section 2);
2. prepare the covariate layers: waterbird abundance and poultry density (Section 3);
3. combine the kernel and covariates into risk maps (Section 4).

All steps run from a single command; see the repository README.

2 Distance transmission kernel

We estimate how the chance of H5N1 appearing in a new area changes with distance from areas where it has already been detected in wild birds. This relationship is the transmission kernel.

2.1 Outbreak data

WAHIS. Outbreak records for H5N1 in wild birds come from the WOAH World Animal Health Information System, event 7649. Each record has a location and a `startDate`, the declared start of the outbreak. For wild birds this is usually when carcasses were found or sampled, not when birds were infected. WAHIS records were exported manually from the event dashboard on 10 August 2026, as WAHIS offers no documented public interface for downloading event data. This is the only input the workflow does not retrieve itself.

DAFF supplement. The WAHIS event had not been updated since 7 August 2026 (edit: a later update on 10 October was not included), so more recent detections were taken from the Australian Government Department of Agriculture, Fisheries and Forestry (DAFF) "Events by species" table, retrieved through the public Power BI service behind the page. The table can only be retrieved as it stands on the day, so each retrieval is saved as a dated snapshot; this analysis uses the snapshot of 21 October 2026.

Locating DAFF records. DAFF reports locations only to locality (suburb or town). Each record was matched to a locality boundary from the Australian Bureau of Statistics (ABS Suburbs and Localities, 2021) on locality name *and* state, ignoring case. ABS adds a state suffix to names used in more than one state (e.g. "Richmond (Tas.)"); the suffix was removed before matching. Each matched record was given one random point within its locality (fixed seed). Unmatched records were left without coordinates.

Combining sources. The two sources overlap. Records with the same locality, date and species in both were flagged as probable duplicates but not removed at this stage.

2.2 Grid and risk set

WAHIS and DAFF record only where the virus *was* detected. The likelihood also needs areas that were at risk but not infected, so we built that set ourselves. Australia (GADM 4.1, excluding external territories) was divided into square cells 25 km wide on the Australian Albers equal-area projection (EPSG:3577). The same grid is used for every layer in the analysis.

Each cell is either *activated*, taking the date of its earliest detection, or never activated. Time runs in steps of 7 days.

2.3 Infectious window

A cell is detected some time after birds there were first infected, and it can keep transmitting after detection because birds shed virus before they die and uncollected carcasses remain infectious. Each activated cell is therefore given an infectious window relative to its detection date:

- it becomes infectious $\Delta = -7$ days relative to detection (negative means before detection);
- it stays infectious for $P = 21$ days.

A cell can therefore act as a source before it is itself detected. The data cannot estimate Δ or P, so both are assumptions.

2.4 Kernel model

On each time step t, every cell not yet activated is exposed to infection pressure from all cells infectious on that step. The infection pressure (force of infection) on cell i is

$$\lambda_i(t) = \beta_0 \sum_{j \in A(t),\, j \neq i} K(d_{ij}),$$

where

- $A(t)$ is the set of cells infectious at step t;
- d_{ij} is the distance (km) between the centres of cells i and j;
- $K(d)$ is the kernel, whose shape is estimated;
- β_0 scales the overall strength of spread.

The probability that cell i is activated on step t is $p_i(t) = 1 - e^{-\lambda_i(t)}$, which turns a rate into a probability between 0 and 1.

Illustrative example. One infectious cell lies 50 km from cell i. With a power-law kernel, $\rho_0 = 20$ km and $\psi = 2$: $K(50) = (1 + 50/20)^{-2} = 3.5^{-2} \approx 0.082$. With $\beta_0 = 0.1$, $\lambda_i = 0.0082$ and $p_i = 1 - e^{-0.0082} \approx 0.008$: about a 0.8% chance of activation in that week.

We fit and compare four kernel shapes, each with a distance scale ρ_0 and a shape parameter ψ:

Kernel	$K(d)$	Tail at long distances
Fat-tailed	$1/\left(1 + (d/\rho_0)^{\psi}\right)$	power law
Power law (type 2)	$(1 + d/\rho_0)^{-\psi}$	power law
Exponential (type 2)	$1 - \exp\left(-(d/\rho_0)^{-\psi}\right)$	power law
Exponential (type 3)	$\exp\left(-(d/\rho_0)^{\psi}\right)$	falls quickly to zero

> **Note**
>
> No covariates enter the kernel. It is fitted on distance and timing alone; the covariates are applied afterwards and are not estimated from these data.

Assumptions. Transmission depends only on distance between cell centres, equally in all directions; every infectious cell is equally infectious; all infections arise from other detected cells (there is no background rate for introductions from outside the detected set); and every infection is eventually detected, at the same delay everywhere.

2.5 Likelihood

The likelihood combines two kinds of evidence:

- each activated cell contributes the probability that it activated on the step it did, $1 - e^{-\lambda_i(t_i)}$, where t_i is its activation step;
- each cell contributes the probability that it did *not* activate on every earlier step while it was at risk, $e^{-\lambda_i(t)}$.

Taking logs:

$$\log L = \sum_{i \in S} \log\left(1 - e^{-\lambda_i(t_i)}\right) - \sum_{i=1}^{n} \sum_{t=1}^{r_i} \lambda_i(t),$$

where S is the set of activated cells, n the number of cells, and r_i the last step cell i was at risk (the step before activation, or the end of the study period for cells never activated). The first

term favours parameter values that put high pressure where and when detections happened. The second penalises values that put high pressure on cells that stayed free of detections; most of the information about how fast the kernel decays comes from this term.

Cells activated when no other cell was infectious have no source to attribute them to. These index cases are treated as given and excluded from the first term.

2.6 Estimation

Parameters were estimated by maximum likelihood, optimised on the log scale to keep them positive. Because the likelihood can have long, flat ridges (Section 2.7), each kernel was fitted from three starting distance scales ($\rho_0 = 5$, 20 and 80 km) and the best fit kept. Approximate confidence intervals come from the curvature of the likelihood at the optimum. Kernel shapes were compared by AIC. To keep computation feasible, distances in the second term were grouped into 1 km bins; the first term uses exact distances. No maximum distance was imposed.

2.7 Results




Figure 1: Fitted kernels: median transmission rate per pair of cells per week, with 50% and 95% uncertainty bands, from the shortest observed distance (one cell width). Bands extending off the plot are marked.

Table 1: Fitted kernels, ordered by AIC.

Kernel	ΔAIC	β_0	ρ_0 (km), 95% CI	ψ, 95% CI
Power law (type 2)	0.00	4.96	5.3 (0.3–91)	2.74 (2.31–3.25)
Exponential (type 3)	0.08	5.7×10^{25}	≈ 0	0.04
Fat-tailed	0.30	0.20	15.3 (3.0–78)	2.65 (2.36–2.98)

Kernel	ΔAIC	β_0	ρ_0 (km), 95% CI	ψ, 95% CI
Exponential (type 2)	0.36	0.13	17.6 (3.2–96)	2.64 (2.37–2.93)

Interpretation:

- **The four shapes cannot be distinguished.** All are within 0.4 AIC units. Over the observed distances (25 km and above) they give practically the same transmission rate per pair of cells per week: about 0.042 at 25 km, 0.008 at 50 km, 0.0013 at 100 km and 0.00002 at 500 km, agreeing to within about 5% at every distance.
- **The tail exponent ψ is the robust result.** It lies between 2.64 and 2.74 for the three power-law-tailed kernels, with overlapping 95% intervals spanning roughly 2.3 to 3.3, and between 2.6 and 3.2 across every alternative fit we ran.
- **The distance scale ρ_0 is not identifiable at this resolution.** Its 95% intervals span roughly 0.3–96 km. ρ_0 sets where the kernel flattens at short range, but no pair of cells is closer than 25 km, so the data constrain only the decline beyond that, governed mainly by ψ. ρ_0 and β_0 therefore trade off against each other (β_0 varies almost 40-fold between shapes giving the same curve), and neither is interpreted on its own.
- **The exponential (type 3) fit is degenerate.** Its parameters have moved to an extreme of the parameter space ($\rho_0 \to 0$, β_0 very large), where its curve over the observed distances mimics a power law. Its near-equal AIC is not support for a thin-tailed kernel, and its parameters have no physical meaning.
- **The transmission rate per pair of cells falls steeply with distance**, about 30-fold between 25 and 100 km. But the number of cells at a given distance grows in proportion to that distance, so the contribution of longer distances to new infections declines much more slowly (roughly as distance$^{-1.7}$ rather than distance$^{-2.7}$). Longer-range spread therefore matters more than the per-pair curve suggests.

2.8 National kernel layer

For every cell, the fitted kernel (power law type 2) was summed over all detected cells and divided by the largest value among undetected cells, so the layer runs from 0 to 1. Detected cells were assigned the maximum value, 1: a detected cell's own contribution at distance zero is not informed by the data, and would otherwise set the scale for every other cell.

The layer sums over every detection since June 2026 with equal weight. It therefore shows **where infection pressure has accumulated over the outbreak period**, not current risk. (Note: the infection period window is used only in fitting.)




Figure 2: Kernel-only relative risk. Detected cells in blue.

Interpretation. The layer sums the fitted power-law kernel over all detected cells (blue), rescaled so the highest undetected cell is 1. Pressure is highest (0.4–1) within 25–50 km of dense clusters of detections: the South Australian gulfs and Coorong, the Victorian coast and Tasmania. Isolated detections produce only small halos (≤ 0.25), because pressure decays steeply with distance ($\psi \approx 2.7$), but where detections are numerous their tails add up to a broad low band (0.01–0.05) across south-eastern Australia. The north and interior are below 0.01.

The layer reflects distance from detections only, not habitat or hosts, and weights all detections equally regardless of date, so it shows the (detected) outbreak's cumulative footprint rather than current pressure. The width of the halos depends on the poorly identified distance scale (ρ_0); the decay rate (ψ) is more robust.

3 Covariates

3.1 Waterbird relative abundance

This layer is a proxy for the abundance of potential waterbird "superspreaders": species likely to drive the spread of H5N1 across the country. Candidate species are assumed to:

- be highly abundant and highly or moderately dispersive;
- have had avian influenza (any strain) detected before, including recent H5N1 detections.

The species used are Greater Crested Tern, Silver Gull, Pacific Black Duck, Grey Teal, Black Swan, Australian Ibis, Pied Cormorant and Little Pied Cormorant. Australian Wood Duck and Australian Pelican were also candidates but have no eBird Status & Trends product.

Weekly relative abundance maps (median, 27 km; 2023 release) for each species were downloaded from eBird Status & Trends, reprojected onto the analysis grid (bilinear interpolation), and averaged over the weeks of each month. The layer is the **sum of the species' relative abundances** in each cell. eBird relative abundance is the expected count on a standardised checklist, so the sum approximates the expected number of these waterbirds seen on such a checklist. Two consequences:

- abundant, flocking species (notably Silver Gull and Grey Teal) dominate the sum;
- species differ in how easily they are detected, which the sum does not correct for.

The layer is a repeating seasonal pattern, not an observation of the current year. The maps use November (Figure 3).

3.2 Poultry density

Chicken density (birds per km^2) comes from the FAO Gridded Livestock of the World v4 (2020), at 5 arc-minutes (about 10 km). Each 25 km cell takes the mean density of the source cells it covers. Cells without a GLW4 estimate inside the boundary (mainly inland lakes and the coastal fringe) were set to zero. The layer totals about 114 million birds, consistent with the national flock.

The distribution is extremely skewed: half of the cells with any poultry hold less than one bird per cell, and these cells together hold 0.15% of all birds (Figure 4).




Figure 3: Summed relative abundance of multiple waterbird species. Relative abundance for individual species sourced from eBird Status & Trends.




Figure 4: Chicken density (FAO GLW4, 2020) on the 25 km grid, log colour scale. Cells with no chickens are left blank.

4 Risk

With few outbreaks, risk factors cannot be estimated from the data, so the layers are combined by fixed rules. For cell i:

Wild birds. Pressure from detections is weighted by how many waterbirds are present:

$$R_i^{\text{wild}} = \text{rescale}\left(K_i \cdot x_i^{\text{wb}}\right),$$

where K_i is the national kernel layer (0–1), x_i^{wb} is the waterbird layer rescaled to 0–1, and rescale divides by the range across cells so the result runs from 0 to 1.

Spillover to poultry. Spillover requires both virus arrival and birds at risk, so the two combine multiplicatively; high poultry density cannot make up for no virus pressure. Exposure rises with poultry density but with diminishing returns: the first farms in an area matter more than doubling an already dense one.

$$R_i^{\text{poultry}} = \text{rescale}\left(R_i^{\text{wild}} \cdot E_i\right), \qquad E_i = \frac{h_i}{h_i + k},$$

where h_i is poultry density (birds per km^2), E_i is exposure (0 with no poultry, approaching 1 when dense), and k is the half-saturation density at which exposure reaches half its maximum. We set k to the density above which half of all birds live, about 1,030 birds per km^2.

Illustrative example. A cell with $R^{\text{wild}} = 0.4$ and $h = 1{,}030$ has $E = 0.5$ and a spillover score of 0.20 before rescaling. With $h = 7{,}000$, $E = 0.87$ and the score is 0.35. With no poultry, the score is 0 whatever the wild-bird risk.

Assumptions. Exposure saturates with density; wild-bird risk and poultry exposure are both required; the waterbird month is the same everywhere. Nothing in the data estimates k: a different k changes how strongly the densest areas stand out. If the highest-ranked cells change substantially across a plausible range of k, the spillover ranking depends on this assumption.

The results are ordinal scores for ranking cells, not probabilities or expected numbers of detections.




Figure 5: Relative risk of H5N1 in wild birds (detected cells in blue) and of spillover to poultry, November.

Interpretation:

Both maps are relative: each is rescaled so that its highest cell is 1, so values rank cells within a map but cannot be compared between maps or read as probabilities. Most of the continent falls in the lowest class (< 0.01) on both.

Wild birds (November). Relative risk is concentrated along the south-east coast: coastal Victoria, with the highest values around Port Phillip Bay; the South Australian coast around the gulfs and the Coorong; and northern and eastern Tasmania. Lower but non-negligible values extend along the New South Wales coast and parts of the south-west of Western Australia. The interior is near zero, because the fitted kernel decays steeply with distance (tail exponent $\psi \approx 2.7$) and waterbird abundance is low there in November. Cells with existing detections carry the maximum kernel value, so the highest-ranked areas partly restate where H5N1 has already been found; the map is most informative for undetected cells near those areas.

Spillover to poultry. Under the multiplicative form, a cell ranks highly only if both its wild bird risk and its chicken density are high. This leaves a single dominant area around Port Phillip Bay (Melbourne–Geelong), where high wild bird risk coincides with dense commercial poultry, and low values near Adelaide and on the New South Wales coast. Because the surface is rescaled to its maximum, this one area compresses all others towards zero; *areas in the lowest classes are lower-ranked, not risk-free.*

What the maps do and do not show. They combine infection pressure from detections to date, as described by the fitted kernel, with waterbird abundance and chicken density, using weightings we chose rather than estimated. They have not been validated against later detections, and they are not forecasts. They also carry no uncertainty: the kernel's distance

scale (ρ_0) is poorly identified, and the spillover ranking depends on the poultry half-saturation density ($k \approx 1030$ birds/km^2) and on the multiplicative form.

5 Uncertainty and limitations

Not yet assessed. The sensitivity analyses below were planned but have not been run for this report. Until they are, the maps should be read as one plausible scenario.

- the infectious window: $\Delta \in \{-14, -7, -2\}$ days, $P \in \{21, 35, 60\}$ days;
- grid placement and resolution (e.g. offsets of a third of a cell; 10 and 50 km cells);
- removing probable duplicate records;
- the poultry half-saturation density k.

Detection, not just transmission. The kernel treats every pair of nearby detections in the same window as transmission. Nearby detections also arise from where people find and report carcasses, from one mass-mortality event reported from several localities, from remaining duplicates, and from DAFF records placed at random within their localities. Part of the steep short-range decline may therefore reflect clustered detection. Correcting for it needs a detection model, for example based on human population within travelling distance; more than half of the grid cells have no resident population, so such a model needs a baseline detection rate.

The kernel layer shows cumulative, not current, pressure. The infectious window is used in fitting but not in mapping: the layer sums over all detections to date. A map of current risk would sum only over cells infectious on a chosen date, and could then be expressed as a weekly probability of infection rather than a relative ranking. A series of weekly maps through the outbreak would show how pressure has shifted.

Locations. DAFF records are located only to locality, and some rural localities are comparable in size to a grid cell.

Time. The kernel layer weights all detections since June equally, and the waterbird layer is a long-term seasonal average. The maps are therefore a summary of the outbreak period, not a current-state estimate or a forecast.

Validation. The maps have not been tested against later detections. A direct test is to build the maps from data up to an earlier date and check whether the next detections fall in the highest-ranked cells.