

Announcements & Such

- *Fleet Foxes*
- Administrative Stuff
 - **Take-Home Mid-Term re-sub**s are due **Thursday**.
 - ☞ **When you turn in resubmissions, make sure that you staple them to your original homework submission.**
 - I will be discussing the grade curve for the course as soon as all of the mid-term grades are in (both the take-home and the in-class).
 - Branden will not be holding office hours this week.
- Today: Chapter 4 — Natural Deduction Proofs for LSL
 - We'll be done with the LSL *natural deduction rules* for \vdash this week.
 - **MacLogic** — a useful computer program for natural deduction.
 - * See <http://fitelson.org/maclogic.htm>.
 - ☞ **Natural deductions are the most challenging topic of the course.**

The Elimination Rule for \sim

Rule of \sim -Elimination: For any formula q , if ' $\sim q$ ' has been inferred at a line j in a proof and q at line k ($j < k$ or $j > k$) then we may infer ' \wedge ' at line m , labeling the line ' $j, k \sim E$ ' and writing on its left the numbers on the left at j and on the left at k . Schematically (with $j < k$):

$$\begin{array}{rcl} a_1, \dots, a_n & (j) & \sim q \\ & \vdots & \\ & & \\ b_1, \dots, b_u & (k) & q \\ & \vdots & \\ a_1, \dots, a_n, b_1, \dots, b_u & (m) & \wedge \quad j, k \sim E \end{array}$$

- Note: we have *added* the symbol ' \wedge ' to the language of LSL. It is treated as if it were an *atomic sentence* of LSL. We can now use it in compound sentences (*e.g.*, ' $A \rightarrow \wedge$ ', ' $\sim \sim \wedge$ ', *etc.*).

The Introduction Rule for \sim

Rule of \sim -Introduction: If ' \wedge ' has been inferred at line k in a proof and $\{a_1, \dots, a_n\}$ are the assumption and premise numbers ' \wedge ' depends upon, then if p is an assumption (or premise) at line j , ' $\sim p$ ' may be inferred at line m , labeling the line ' $j, k \sim I$ ' and writing on its left the numbers in the set $\{a_1, \dots, a_n\}/j$.

$$\begin{array}{rcl} j & (j) & p \quad \text{Assumption} \\ & \vdots & \\ a_1, \dots, a_n & (k) & \wedge \\ & \vdots & \\ \{a_1, \dots, a_n\}/j & (m) & \sim p \quad j, k \sim I \end{array}$$

- $\sim I$ is used (typically with $\sim E$) to deduce ' $\sim p$ ' via *reductio ad absurdum*, by (i) *assuming* p , (ii) deducing ' \wedge ', and (iii) *discharging* the assumption.

The Rule of Double Negation (DN)

- Negation is an odd connective in our system. It not only has an introduction rule and an elimination rule, but it also has an additional rule called the *double negation* (DN) rule.
- The DN rule says that we may infer p from ' $\sim \sim p$ '. Without this DN rule, we would not be able to prove certain valid LSL argument forms — *e.g.*, $\sim(A \& \sim B) \therefore (A \rightarrow B)$.

Rule of Double Negation: For any formula p , if ' $\sim \sim p$ ' has been inferred at a line j in a proof, then at line k we may infer p , labeling the line ' j ' and writing on its left the numbers to the left of j .

$$\begin{array}{rcl} a_1, \dots, a_n & (j) & \sim \sim p \\ a_1, \dots, a_n & (k) & p \quad j \text{ DN} \end{array}$$

Example Proof of a *Theorem*

- Using only the rules we have learned so far, we should be able to prove the following *theorem*: $\vdash \sim(A \& \sim A)$. Let's do this one by hand first.
- Here's a simple proof, generated using MacLogic (I'll show how):

Problem is: $\vdash \sim(A \& \sim A)$

1	(1)	$A \& \sim A$	Assumption (!)
1	(2)	$\sim A$	1 &E
1	(3)	A	1 &E
1	(4)	Δ	2,3 $\sim E$
	(5)	$\sim(A \& \sim A)$	1,4 $\sim I$

- This proof makes use of *no premises*, and its final line has *no numbers to its left* — indicating that we have succeeded in proving ' $\sim(A \& \sim A)$ ' from *nothing at all*. It's a *theorem* (i.e., a sequent with no premises)!

The Introduction Rule for \vee ($\vee I$)

Rule of \vee -Introduction: For any formula p , if p has been inferred at line j , then, for any formula q , *either* ' $p \vee q$ ' *or* ' $q \vee p$ ' may be inferred at line k , labeling the line ' $j \vee I$ ' and writing on its left the same premise and assumption numbers as appear on the left of j .

a_1, \dots, a_n	(j)	p		a_1, \dots, a_n	(j)	q
	\vdots		OR		\vdots	
a_1, \dots, a_n	(k)	$p \vee q$	$j \vee I$	a_1, \dots, a_n	(k)	$p \vee q$
						$j \vee I$

- The $\vee I$ rule is very simple and intuitive. Basically, it says that you may infer a disjunction from *either* of its disjuncts.
- The *elimination* rule ($\vee E$) for \vee , on the other hand, is considerably more complex to state and apply. It's the hardest of our rules.

The Elimination Rule for \vee ($\vee E$)

- First, the idea *behind* the \vee -elimination rule.
- The following argument form is valid (easily verified *via* truth-table):

$$\begin{array}{l} p \vee q \\ p \rightarrow r \\ q \rightarrow r \\ \hline \therefore r \end{array}$$

- This argument form is called the *constructive dilemma*. In essence, the $\vee E$ rule reflects the constructive dilemma form of reasoning and implements it in our system of natural deduction rules.
- The $\vee E$ rule is trickier than our other rules because it requires us to make *two* assumptions. This can make it rather complicated to keep track of all of our assumptions and premises during an $\vee E$ proof.
- Now, the official definition of $\vee E$...

Rule of \vee -Elimination: If a disjunction ' $p \vee q$ ' occurs at line g of a proof, p is assumed at line h , r is derived at line i , q is assumed at line j , and r is derived at line k , then at line m we may infer r , labeling the line ' $g, h, i, j, k \vee E$ ' and writing on its left every number on the left at line g , and at line i (except h), and at line k (except j).

a_1, \dots, a_n	(g)	$p \vee q$	
	\vdots		
	h	(h) p	Assumption
	\vdots		
b_1, \dots, b_u	(i)	r	
	\vdots		
	j	(j) q	Assumption
	\vdots		
c_1, \dots, c_w	(k)	r	
	\vdots		
\mathcal{A}	(m)	r	$g, h, i, j, k \vee E$

where \mathcal{A} is the set: $\{a_1, \dots, a_n\} \cup \{b_1, \dots, b_u\}/h \cup \{c_1, \dots, c_w\}/j$.

An Example Involving \vee E and DN

- Here's a proof of the sequent: $A \vee B, \sim B \vdash A$.

Problem is: $A \vee B, \sim B \vdash A$

1	(1) $A \vee B$	Premise
2	(2) $\sim B$	Premise
3	(3) $\sim A$	Assumption (for \sim I)
4	(4) A	Assumption (for \vee E)
3,4	(5) Δ	3,4 \sim E
6	(6) B	Assumption (for \vee E)
2,6	(7) Δ	2,6 \sim E
1,2,3	(8) Δ	1,4,5,6,7 \vee E
1,2	(9) $\sim\sim A$	3,8 \sim I
1,2	(10) A	9 DN

A Simple Example Involving \vee I and \vee E

- Here's a proof of the sequent: $A \vee B \vdash B \vee A$.

Problem is: $A \vee B \vdash B \vee A$

1	(1) $A \vee B$	Premise
2	(2) A	Assumption (\vee E)
2	(3) $B \vee A$	2 \vee I
4	(4) B	Assumption (\vee E)
4	(5) $B \vee A$	4 \vee I
1	(6) $B \vee A$	1,2,3,4,5 \vee E

Another Example Involving \vee I and Negation

- Here's a proof of the *theorem*: $\vdash A \vee \sim A$.

Problem is: $\vdash A \vee \sim A$

1	(1) $\sim(A \vee \sim A)$	Assumption (\sim I)
2	(2) A	Assumption (\sim I)
2	(3) $A \vee \sim A$	2 \vee I
1,2	(4) Δ	1,3 \sim E
1	(5) $\sim A$	2,4 \sim I
1	(6) $A \vee \sim A$	5 \vee I
1	(7) Δ	1,6 \sim E
1	(8) $\sim\sim(A \vee \sim A)$	1,7 \sim I
1	(9) $A \vee \sim A$	8 DN

A Third Example Involving \vee E

- Here's a proof of the sequent: $A \vee B, \sim B \vdash A$.

Problem is: $A \vee B, \sim B \vdash A$

1	(1) $A \vee B$	Premise
2	(2) $\sim B$	Premise
3	(3) $\sim A$	Assumption (for \sim I)
4	(4) A	Assumption (for \vee E)
3,4	(5) Δ	3,4 \sim E
6	(6) B	Assumption (for \vee E)
2,6	(7) Δ	2,6 \sim E
1,2,3	(8) Δ	1,4,5,6,7 \vee E
1,2	(9) $\sim\sim A$	3,8 \sim I
1,2	(10) A	9 DN

A Fourth Example Involving \vee I and \vee E

- Here's a proof of the sequent: $A \vee (B \& C) \vdash (A \vee B) \& (A \vee C)$.

1	(1) $A \vee (B \& C)$	Premise
2	(2) A	Assumption (\vee E)
2	(3) $A \vee B$	2 \vee I
2	(4) $A \vee C$	2 \vee I
2	(5) $(A \vee B) \& (A \vee C)$	3,4 &I
6	(6) $B \& C$	Assumption (\vee E)
6	(7) B	6 &E
6	(8) $A \vee B$	7 \vee I
6	(9) C	6 &E
6	(10) $A \vee C$	9 \vee I
6	(11) $(A \vee B) \& (A \vee C)$	8,10 &I
1	(12) $(A \vee B) \& (A \vee C)$	1,2,5,6,11 \vee E

Another Example Involving \vee

- Let's do a proof of: $(A \& B) \vee (A \& C) \vdash A \& (B \vee C)$

1	(1) $(A \& B) \vee (A \& C)$	Premise
2	(2) $A \& B$	Ass (\vee E)
2	(3) A	2 &E
4	(4) $A \& C$	Ass (\vee E)
4	(5) A	4 &E
1	(6) A	1,2,3,4,5 \vee E
2	(7) B	2 &E
2	(8) $B \vee C$	7 \vee I
4	(9) C	4 &E
4	(10) $B \vee C$	9 \vee I
1	(11) $B \vee C$	1,2,8,4,10 \vee E
1	(12) $A \& (B \vee C)$	6,11 &I

A Final Example Involving \vee and \sim

- Let's do a proof of: $\sim A \vee B \vdash A \rightarrow B$

Problem is : $\sim A \vee B \vdash A \rightarrow B$

1	(1) $\sim A \vee B$	Premise
2	(2) A	Assumption (\rightarrow I)
3	(3) $\sim A$	Assumption (\vee E)
4	(4) $\sim B$	Assumption (\sim I)
2,3	(5) Δ	3,2 \sim E
2,3	(6) $\sim \sim B$	4,5 \sim I
2,3	(7) B	6 DN
8	(8) B	Assumption (\vee E)
1,2	(9) B	1,3,7,8,8 \vee E
1	(10) $A \rightarrow B$	2,9 \rightarrow I

General Tips on Proof Strategy and Planning

- As a first line of attack, always try to prove your conclusion by using the introduction rule for its main connective as the main strategy.
- This will indicate what assumptions, if any, need to be made and what other formulae will need to be derived. This is "working backward".
- If these other formulae also contain connectives, then try to prove them by introducing their main connectives. Work backward, as far as possible.
- When this technique can no longer be applied, inspect your current stock of premises and assumptions to see if they have any *obvious* consequences.
- If your current premises and assumption contain a disjunction ' $r \vee s$ ', see if you can prove your current goal formula p from *each* of its disjuncts r and s (using your current premises and assumptions). If you think you can, then try using \vee E to prove p . If no disjunction appears anywhere in your current of premises/assumptions, then \vee E is probably not a good strategy.
- If you have tried everything you can think of to prove your current goal p , try assuming ' $\sim p$ ' and aim for ' $\sim \sim p$ ' by \sim E, \sim I; then use DN.

When to Make Assumptions, and When *Not* to

- In constructing a proof, any assumptions you make must eventually be discharged, so you should only make assumptions in connection with the three rules which discharge assumptions.
- In other words, if you make an assumption p in a proof, you *must* be able to give one of the following three reasons:
 1. p is the antecedent of a conditional ' $p \rightarrow q$ ' you are trying to derive using the \rightarrow I rule (then, try to prove q).
 2. You are trying to derive ' $\sim p$ ', so you assume p with an eye toward using the \sim I rule (then, try to prove \perp).
 3. p is one of the disjuncts of a disjunction ' $p \vee q$ ' (*somewhere in your current stock of premises and assumptions!*) to which you will be applying \vee E (then, try to prove some r from each).
- Remember, only the three rules \rightarrow I, \sim I, and \vee E involve making assumptions. *No other rules can discharge assumptions.*

10 More Examples Involving \vee I and \vee E

1. $(A \& B) \vee (A \& C) \vdash A$ [p. 111, ex. 2]
2. $(A \rightarrow \perp) \vee (B \rightarrow \perp), B \vdash \sim A$ [p. 116, §4.5, ex. 11]
3. $(A \vee B) \vee C \vdash A \vee (B \vee C)$ [p. 116, ex. 19]
4. $A \vee B \vdash (A \rightarrow B) \rightarrow B$ [p. 116, ex. 10]
5. $A \& B \vdash \sim(\sim A \vee \sim B)$ [p. 116, ex. 14 (\vdash)]
6. $A \vee B \vdash \sim(\sim A \& \sim B)$ [p. 116, ex. 13]
7. $\sim(A \& B) \vdash \sim A \vee \sim B$ [p. 116, ex. 16 (\rightarrow)]
8. $\sim C \vee (A \rightarrow B) \vdash (C \& A) \rightarrow B$ [not in text]
9. $\vdash (A \rightarrow B) \vee (B \rightarrow A)$ [not in text]
10. $\sim(A \vee B) \vdash \sim A \& \sim B$ [not in text]

Proof of Example #1

Problem is: $(A \& B) \vee (A \& C) \vdash A$

1	(1) $(A \& B) \vee (A \& C)$	Premise
2	(2) $A \& B$	Assumption (\vee E)
2	(3) A	2 &E
4	(4) $A \& C$	Assumption (\vee E)
4	(5) A	4 &E
1	(6) A	1,2,3,4,5 \vee E

Proof of Example #2

Problem is: $(A \rightarrow \Delta) \vee (B \rightarrow \Delta), B \vdash \sim A$

1	(1) $(A \rightarrow \Delta) \vee (B \rightarrow \Delta)$	Premise
2	(2) B	Premise
3	(3) A	Assumption (\sim I)
4	(4) $A \rightarrow \Delta$	Assumption (\vee E)
3,4	(5) Δ	4,3 \rightarrow E
6	(6) $B \rightarrow \Delta$	Assumption (\vee E)
2,6	(7) Δ	6,2 \rightarrow E
1,2,3	(8) Δ	1,4,5,6,7 \vee E
1,2	(9) $\sim A$	3,8 \sim I

Proof of Example #3

Problem is: $(A \vee B) \vee C \vdash A \vee (B \vee C)$

1	(1) $(A \vee B) \vee C$	Premise
2	(2) $A \vee B$	Assumption ($\vee E$)
3	(3) A	Assumption ($\vee E$)
3	(4) $A \vee (B \vee C)$	3 $\vee I$
5	(5) B	Assumption ($\vee E$)
5	(6) $B \vee C$	5 $\vee I$
5	(7) $A \vee (B \vee C)$	6 $\vee I$
2	(8) $A \vee (B \vee C)$	2,3,4,5,7 $\vee E$
9	(9) C	Assumption ($\vee E$)
9	(10) $B \vee C$	9 $\vee I$
9	(11) $A \vee (B \vee C)$	10 $\vee I$
1	(12) $A \vee (B \vee C)$	1,2,8,9,11 $\vee E$

Proof of Example #4

Problem is : $A \vee B \vdash (A \rightarrow B) \rightarrow B$

1	(1) $A \vee B$	Premise
2	(2) $A \rightarrow B$	Ass ($\rightarrow I$)
3	(3) A	Ass ($\vee E$)
2,3	(4) B	2,3 $\rightarrow E$
5	(5) B	Ass ($\vee E$)
1,2	(6) B	1,3,4,5,5 $\vee E$
1	(7) $(A \rightarrow B) \rightarrow B$	2,6 $\rightarrow I$

Proof of Example #5

Problem is: $A \& B \vdash \sim(\sim A \vee \sim B)$

1	(1) $A \& B$	Premise
2	(2) $\sim A \vee \sim B$	Assumption ($\sim I$)
3	(3) $\sim A$	Assumption ($\vee E$)
1	(4) A	1 $\&E$
1,3	(5) Δ	3,4 $\sim E$
6	(6) $\sim B$	Assumption ($\vee E$)
1	(7) B	1 $\&E$
1,6	(8) Δ	6,7 $\sim E$
1,2	(9) Δ	2,3,5,6,8 $\vee E$
1	(10) $\sim(\sim A \vee \sim B)$	2,9 $\sim I$

Proof of Example #6

Problem is : $A \vee B \vdash \sim(\sim A \& \sim B)$

1	(1) $A \vee B$	Premise
2	(2) $\sim A \& \sim B$	Ass ($\sim I$)
3	(3) A	Ass ($\vee E$)
2	(4) $\sim A$	2 $\&E$
2,3	(5) Δ	4,3 $\sim E$
6	(6) B	Ass ($\vee E$)
2	(7) $\sim B$	2 $\&E$
2,6	(8) Δ	7,6 $\sim E$
1,2	(9) Δ	1,3,5,6,8 $\vee E$
1	(10) $\sim(\sim A \& \sim B)$	2,9 $\sim I$

Proof of Example #7Problem is: $\sim(A \& B) \vdash \sim A \vee \sim B$

1	(1) $\sim(A \& B)$	Premise
2	(2) $\sim(\sim A \vee \sim B)$	Assumption ($\sim I$)
3	(3) $\sim A$	Assumption ($\sim I$)
3	(4) $\sim A \vee \sim B$	3 $\vee I$
2,3	(5) Δ	2,4 $\sim E$
2	(6) $\sim \sim A$	3,5 $\sim I$
2	(7) A	6 DN
8	(8) $\sim B$	Assumption ($\sim I$)
8	(9) $\sim A \vee \sim B$	8 $\vee I$
2,8	(10) Δ	2,9 $\sim E$
2	(11) $\sim \sim B$	8,10 $\sim I$
2	(12) B	11 DN
2	(13) $A \& B$	7,12 $\& I$
1,2	(14) Δ	1,13 $\sim E$
1	(15) $\sim \sim(\sim A \vee \sim B)$	2,14 $\sim I$
1	(16) $\sim A \vee \sim B$	15 DN

Proof of Example #8Problem is: $\sim C \vee (A \rightarrow B) \vdash (C \& A) \rightarrow B$

1	(1) $\sim C \vee (A \rightarrow B)$	Premise
2	(2) $C \& A$	Assumption ($\rightarrow I$)
3	(3) $\sim B$	Assumption ($\sim I$)
4	(4) $\sim C$	Assumption ($\vee E$)
2	(5) C	2 $\& E$
2,4	(6) Δ	4,5 $\sim E$
7	(7) $A \rightarrow B$	Assumption ($\vee E$)
2	(8) A	2 $\& E$
2,7	(9) B	7,8 $\rightarrow E$
2,3,7	(10) Δ	3,9 $\sim E$
1,2,3	(11) Δ	1,4,6,7,10 $\vee E$
1,2	(12) $\sim \sim B$	3,11 $\sim I$
1,2	(13) B	12 DN
1	(14) $(C \& A) \rightarrow B$	2,13 $\rightarrow I$

Proof of Example #9Problem is: $\vdash (A \rightarrow B) \vee (B \rightarrow A)$

1	(1) $\sim((A \rightarrow B) \vee (B \rightarrow A))$	Assumption ($\sim I$)
2	(2) B	Assumption ($\rightarrow I$)
3	(3) $\sim A$	Assumption ($\sim I$)
4	(4) A	Assumption ($\rightarrow I$)
2	(5) $A \rightarrow B$	4,2 $\rightarrow I$
2	(6) $(A \rightarrow B) \vee (B \rightarrow A)$	5 $\vee I$
1,2	(7) Δ	1,6 $\sim E$
1,2	(8) $\sim \sim A$	3,7 $\sim I$
1,2	(9) A	8 DN
1	(10) $B \rightarrow A$	2,9 $\rightarrow I$
1	(11) $(A \rightarrow B) \vee (B \rightarrow A)$	10 $\vee I$
1	(12) Δ	1,11 $\sim E$
	(13) $\sim \sim((A \rightarrow B) \vee (B \rightarrow A))$	1,12 $\sim I$
	(14) $(A \rightarrow B) \vee (B \rightarrow A)$	13 DN

Proof of Example #10Problem is: $\sim(A \vee B) \vdash \sim A \& \sim B$

1	(1) $\sim(A \vee B)$	Premise
2	(2) A	Ass ($\sim I$)
2	(3) $A \vee B$	2 $\vee I$
1,2	(4) Δ	1,3 $\sim E$
1	(5) $\sim A$	2,4 $\sim I$
6	(6) B	Ass ($\sim I$)
6	(7) $A \vee B$	6 $\vee I$
1,6	(8) Δ	1,7 $\sim E$
1	(9) $\sim B$	6,8 $\sim I$
1	(10) $\sim A \& \sim B$	5,9 $\& I$