

1. (DF3.4.4) Use Cauchy's Theorem and induction to show that a finite abelian group has a subgroup of order  $n$  for each positive divisor  $n$  of its order.

*Proof.* Let  $G$  be an arbitrary finite abelian group. We proceed by induction on the order of  $G$ ; that is, we assume that for any finite abelian group  $H$  such that  $1 \leq |H| < |G|$ , the group  $H$  has a subgroup of order  $d$  for every positive divisor  $d$  of  $|H|$ . It suffices to show that  $G$  has a subgroup of order  $n$  for every positive divisor  $n$  of  $|G|$ . Whenever  $n$  is prime, we already have by Cauchy's theorem the existence of a subgroup of prime order  $n$ . So without loss of generality, let  $n$  be composite.

Let  $p$  be any prime which divides  $n$ , so that for some positive integer  $k > 1$  (as  $n$  is composite),  $n = pk$ . By Cauchy's Theorem, it follows that  $G$  has a subgroup  $N$  of order  $p$ . Then because all subgroups of abelian groups are abelian, it follows that  $N$  is normal in  $G$  and so the quotient group  $G/N$  is well defined.

Because  $n \mid |G|$ , there exists a positive integer  $q$  such that  $|G| = nq$ . So by Lagrange's theorem, the order of the quotient group  $G/N$  is  $|G|/|N| = nq/p = pkq/p = kq$ , which because  $k > 1$ , the quotient group is not trivial. Furthermore, because  $k > 1$  and  $p$  is a prime number, it follows that  $|G/N| = kq < n$ . Then by the inductive hypothesis, it follows that  $G/N$  contains a subgroup of order  $k$ , which by the Lattice Isomorphism Theorem we may write as  $A/N$  for some  $N \trianglelefteq A \trianglelefteq G$  (all subgroups of  $G$  are normal in larger subgroups of  $G$  containing them because all subgroups of  $G$  are abelian).

Then again using Lagrange's theorem we have that  $|A/N| = |A|/|N| = |A|/p = k$ , which means  $|A| = pk = n$ , so  $A$  is a subgroup of  $G$  with order  $n$ .  $\square$

2. (DF3.4.1) Prove that if  $G$  is an abelian simple group then  $G \cong Z_p$  for some prime  $p$ . (not a homework problem, but needed for DF3.4.8, the next exercise)

*Proof.* Let  $G$  be an abelian simple group. Then there are no normal subgroups outside of the trivial subgroup and  $G$  itself. This means that for some element  $x \neq 1$  in  $G$ , the subgroup  $\langle x \rangle$  is normal in  $G$  because  $G$  is abelian, but because  $x \neq 1$ , this normal subgroup is equal to  $G$ . Hence  $G$  is cyclic.

Then suppose that  $G$  is an infinite cyclic group. Then  $\langle x \rangle = \{x, x^2, \dots\} = G$  but since  $x^2 \neq 1$  (infinite order),  $\langle x^2 \rangle = \{x^2, x^4, \dots\} \neq G$  but  $\langle x^2 \rangle$  is normal in  $G$  because  $G$  is abelian. This is in contradiction to the assumption that  $G$  was simple. So  $G$  has finite order.

Suppose that the order of  $G$  is composite. Then  $|G| = pk$  for some prime  $p$  and some positive integer  $k > 1$  ( $|G|$  is composite). But by Cauchy's theorem, we know that there is a subgroup of  $G$  of order  $p$ , which is strictly less than  $|G|$  and strictly greater than 1, which again is in contradiction to the assumption that  $G$  was simple. Hence the order of  $G$  is not composite; that is, the order of  $G$  is a prime number.

Groups with prime order are known to be cyclic due to Lagrange's theorem. Therefore  $G \cong Z_p$  for some prime  $p$ .  $\square$

3. (DF3.4.8) Let  $G$  be a finite group. Prove that the following are equivalent:

- (i)  $G$  is solvable (there is a chain of subgroups  $1 = G_0 \trianglelefteq G_1 \trianglelefteq G_2 \trianglelefteq \cdots \trianglelefteq G_s = G$  such that  $G_{i+1}/G_i$  is abelian for  $i = 0, 1, \dots, s-1$ )
- (ii)  $G$  has a chain of subgroups:  $1 = H_0 \trianglelefteq H_1 \trianglelefteq H_2 \trianglelefteq \cdots \trianglelefteq H_s = G$  such that  $H_{i+1}/H_i$  is cyclic,  $0 \leq i \leq s-1$
- (iii) all composition factors of  $G$  are of prime order
- (iv)  $G$  has a chain of subgroups:  $1 = N_0 \trianglelefteq N_1 \trianglelefteq N_2 \trianglelefteq \cdots \trianglelefteq N_t = G$  such that each  $N_i$  is a normal subgroup of  $G$  and  $N_{i+1}/N_i$  is abelian,  $0 \leq i \leq t-1$ .

*Proof.* We show that (i) is equivalent to (ii), (i) is equivalent to (iii), and that (i) is equivalent to (iv).

From (i), let  $G$  be a group with a chain of subgroups  $1 = G_0 \trianglelefteq G_1 \trianglelefteq G_2 \trianglelefteq \cdots \trianglelefteq G_s = G$  such that  $G_{i+1}/G_i$  is abelian for  $i = 0, 1, \dots, s-1$ . The factor groups  $G_{i+1}/G_i$  may not be simple, so we must insert intermediate subgroups within the chain which make simple and abelian subgroups.

If a factor group  $G_{i+1}/G_i$  is not simple, there exists a nontrivial proper normal subgroup of  $G_{i+1}/G_i$ , which because  $G_{i+1}/G_i$  is abelian, is also abelian. So by the Lattice Isomorphism Theorem, there exists a nontrivial proper normal subgroup  $A/G_i$  such that  $G_i \leq A \trianglelefteq G_{i+1}$ . Furthermore, because  $G_i$  is a normal subgroup of  $G_{i+1}$ , it follows that  $G_i$  is a normal subgroup of  $A$ .

So  $G_i \trianglelefteq A \trianglelefteq G_{i+1}$ , and  $A/G_i$  is abelian. We need to show that  $G_{i+1}/A$  is abelian. Because  $G_{i+1}/G_i$  is abelian, the multiplication of members  $xG_i, yG_i$  (with any  $x, y \in G_{i+1}$ ) commute:

$$xG_i yG_i = xyG_i = yxG_i = yG_i xG_i$$

So  $(yx)^{-1}xy \in G_i$ , and by inclusion,  $(yx)^{-1}xy \in A$ , which means that  $xAyA = xyA = yxA = yAx$ , so that the factor group  $G_{i+1}/A$  is abelian. It is possible for  $A/G_i$  or  $G_{i+1}/A$  or both to not be simple.

Therefore, it is possible to form a new chain of subgroups from the chain

$$1 = G_0 \trianglelefteq G_1 \trianglelefteq G_2 \trianglelefteq \cdots \trianglelefteq G_i \trianglelefteq G_{i+1} \trianglelefteq \cdots \trianglelefteq G_s = G$$

by replacing  $G_i \trianglelefteq G_{i+1}$  with  $G_i \trianglelefteq A \trianglelefteq G_{i+1}$  when  $G_{i+1}/G_i$  is not simple, to form

$$1 = G_0 \trianglelefteq G_1 \trianglelefteq G_2 \trianglelefteq \cdots \trianglelefteq G_i \trianglelefteq A \trianglelefteq G_{i+1} \trianglelefteq \cdots \trianglelefteq G_s = G.$$

Of course, we can keep expanding out the non-simple factors of this new series that we obtained, but because  $G$  is finite, by the Jordan-Hölder theorem, we will arrive at the composition series for  $G$ . By repeating the above procedure until all of the abelian factor groups  $N_{i+1}/N_i$  for  $0 \leq i \leq t-1$  are simple in the final chain  $1 = N_0 \trianglelefteq N_1 \trianglelefteq N_2 \trianglelefteq \cdots \trianglelefteq N_t = G$ , we arrive at the composition series for  $G$ .

Then by the previous theorem which states that abelian simple groups are isomorphic to a cyclic group of prime order, all of these factor groups  $N_{i+1}/N_i$  are cyclic groups, so (i) implies (ii). Because cyclic groups are abelian by exponent rules, (ii) implies (i). Hence (i) is equivalent to (ii).

Furthermore, by the same theorem, these factor groups (composition factors)  $N_{i+1}/N_i$  are of prime order, so (i) implies (iii). Then if all of the composition factors of  $G$  are of prime order, then necessarily they are cyclic and hence abelian. So (iii) implies (i). Hence (i) is equivalent to (iii).

What remains is to show that **(i)** is equivalent to **(iv)**. It is clear that **(iv)** implies **(i)** because **(iv)** is a stronger statement:

**(iv)**  $G$  has a chain of subgroups:  $1 = N_0 \trianglelefteq N_1 \trianglelefteq N_2 \trianglelefteq \cdots \trianglelefteq N_t = G$  such that each  $N_i$  is a normal subgroup of  $G$  and  $N_{i+1}/N_i$  is abelian,  $0 \leq i \leq t-1$ .

So not only are the subgroups  $N_i$  normal in  $N_{i+1}$  for  $0 \leq i \leq t-1$ , but they are normal in  $G$ . Conversely, assume **(i)** is true. Then let  $M$  be a minimal nontrivial normal subgroup of  $G$ , so that  $M$  does not contain any nontrivial proper subgroups which are normal in  $G$ . Let  $N \trianglelefteq M$  be a normal subgroup of  $M$  of prime index, which exists because  $G$  has a composition series and so  $M$  has one as well (subgroups of solvable groups are solvable), so by **(iii)**,  $N$  exists.

Then the quotient  $M/N$  is cyclic, so it is abelian. This means that for any  $x, y \in M$ ,  $xNyN = xyN = yxN = yNxn$ , which implies that  $(yx)^{-1}xy \in N$ . Similarly, consider  $gNg^{-1}$  for  $g \in G$ . This is a subgroup of  $G$  that is also a normal subgroup of  $M$  because for any  $m \in M$ ,  $mgNg^{-1}m^{-1} \subseteq gNg^{-1}$  is equivalent to showing that  $g^{-1}mgN(g^{-1}mg)^{-1} \subseteq N$ . But  $M \trianglelefteq G$ , so  $g^{-1}mg, (g^{-1}mg)^{-1} \in M$ , and we know that  $N \trianglelefteq M$ , so  $g^{-1}mgN(g^{-1}mg)^{-1} \subseteq N$  is true. Then  $mgNg^{-1}m^{-1} \subseteq gNg^{-1}$  is true, which means that  $gNg^{-1}$  is a normal subgroup of  $M$ .

Then the quotient  $M/gNg^{-1}$  is well defined, and because  $|gNg^{-1}| = |N|$ , this quotient has prime order as well, which means it is cyclic and hence abelian. Similarly, for any  $x, y \in M$ ,  $xgNg^{-1}ygNg^{-1} = xygNg^{-1} = yxgNg^{-1} = ygNg^{-1}xgNg^{-1}$ , so that  $(yx)^{-1}xy \in gNg^{-1}$ . Since  $g$  was arbitrary, the following is true:  $(yx)^{-1}xy \in \bigcap_{g \in G} gNg^{-1}$ .

But observe that  $\bigcap_{g \in G} gNg^{-1}$  is a subgroup of  $N$  since it contains only those elements of  $N$  which appear in each  $gNg^{-1}$ , and this set is a subgroup because the intersection of subgroups is a subgroup. Furthermore,  $\bigcap_{g \in G} gNg^{-1}$  is also a normal subgroup of  $G$ : Let  $h \in G$ . We must show that

$$h \left( \bigcap_{g \in G} gNg^{-1} \right) h^{-1} \subseteq \bigcap_{g \in G} gNg^{-1},$$

but this is clear from the fact that any of the elements on the left hand side take on the form  $g'n'(g')^{-1}$  for every  $g' \in G$  and some  $n' \in N$ . Then  $hg'n'(g')^{-1}h^{-1} = (g'h)n'(g'h)^{-1} \in \bigcap_{g \in G} gNg^{-1}$  since  $g'h$  takes on any element of  $G$ . Hence the intersection is a normal subgroup of  $G$ .

But the minimality of  $M$  suggests that this subgroup of  $N$  (and hence a proper subgroup of  $M$ ) should be the trivial subgroup, and since for every  $x, y \in M$ ,  $(yx)^{-1}xy \in \bigcap_{g \in G} gNg^{-1} = \{1\}$ , we must have that  $(yx)^{-1}xy = 1$  so that  $xy = yx$ . So  $M$  is abelian, and  $M$  was a minimal nontrivial normal subgroup of  $G$ . Furthermore, this means that  $M/1$  is abelian as well.

Suppose by induction that for all groups of a strictly lower order than  $G$  which are solvable, there exists a chain of the form given in **iv** for the lower order groups, where  $G$  is solvable. Then consider the quotient group  $G/M$ , where  $M$  was a minimal nontrivial normal subgroup of  $G$ . This group is solvable because any chain of subgroups of the form  $1 = \overline{G_0} \trianglelefteq \overline{G_1} \trianglelefteq \overline{G_2} \trianglelefteq \cdots \trianglelefteq \overline{G_r} = G/M$ , which by the Lattice Isomorphism Theorem

we can write  $\overline{G_i} = A_i/M$  for some subgroup  $A_i$  containing  $M$  ( $M \trianglelefteq A_i$ ). Then by the Third Isomorphism Theorem, we have that  $\overline{G_{i+1}}/\overline{G_i} = (A_{i+1}/M)/(A_i/M) \cong A_{i+1}/A_i$ , which are abelian since  $G$  is solvable. Hence the quotient group is solvable, furthermore, because  $M$  is nontrivial, the order of the quotient group  $G/M$  is strictly lower than the order of  $G$  and by the inductive hypothesis  $G/M$  has a series as given in **(iv)**, say  $1 = \overline{G_0} \trianglelefteq \overline{G_1} \trianglelefteq \overline{G_2} \trianglelefteq \cdots \trianglelefteq \overline{G_r} = G/M$ , with each  $\overline{G_i} \trianglelefteq G/M$  where by the Lattice Isomorphism Theorem we can write  $\overline{G_i} = G_i/M$  for some subgroup  $G_i$  containing  $M$  ( $M \trianglelefteq G_i$ ). Furthermore, by the Lattice Isomorphism Theorem, each  $G_i \trianglelefteq G$ ; and so by the Third Isomorphism Theorem it follows that  $\overline{G_{i+1}}/\overline{G_i} = (G_{i+1}/M)/(G_i/M) \cong G_{i+1}/G_i$  we can form the chain of subgroups for  $G$ :  $1 \trianglelefteq M = G_0 \trianglelefteq G_1 \trianglelefteq \cdots \trianglelefteq G_{r+1} = G$  where each  $G_i$  is normal in  $G$  as well. Each of the factors  $G_{i+1}/G_i$  (and  $M/1$ ) will be abelian as well since  $\overline{G_{i+1}}/\overline{G_i}$  is abelian by assumption, and  $M$  is abelian. Thus **(i)** implies **(iv)**, so that **(i)** is equivalent to **(iv)**. Therefore, all four statements are equivalent to each other.  $\square$

4. (DF3.5.12) Prove that  $A_n$  contains a subgroup isomorphic to  $S_{n-2}$  for each  $n \geq 3$ .

*Proof.* Let  $n \geq 3$ , since  $A_1$  and  $A_2$  are both the trivial group. View  $S_n$  as the group of permutations of  $\{1, 2, \dots, n\}$ , so that we can view  $S_{n-2}$  as the group of permutations of  $\{1, 2, \dots, n-2\}$ , which are all the permutations of  $S_n$  which fix  $n$  and  $n-1$ . Let  $\tau$  be the transposition in  $S_n$  which interchanges  $n$  and  $n-1$ ; i.e.  $\tau = ((n-1)n)$ . Note that  $\tau$  has order 2, and commutes with all elements of  $S_{n-2}$  which appear in  $S_n$ .

Then there is a homomorphism  $\varphi: S_{n-2} \rightarrow A_n$  given by

$$\varphi(\sigma) = \begin{cases} \sigma, & \text{if } \sigma \text{ is an even permutation} \\ \sigma\tau, & \text{if } \sigma \text{ is an odd permutation} \end{cases}.$$

In both cases we obtain an even permutation (we make odd permutations even by appending  $\tau$  on the right), so the image of the homomorphism is indeed a subgroup of  $A_n$ . We should check that this is a homomorphism. For permutations  $\sigma, \pi \in S_{n-2}$ , we investigate  $\varphi(\sigma\pi)$  for each of the four ways we choose the parity of  $\sigma$  and  $\pi$ , remembering that  $\tau^2 = 1$ , and that  $\tau$  commutes with all elements of  $S_{n-2}$  which appear in  $S_n$ .

$\sigma$	$\pi$	$\sigma\pi$	$\varphi(\sigma\pi) \stackrel{?}{=} \varphi(\sigma)\varphi(\pi)$
even	even	even	$\varphi(\sigma\pi) = \sigma\pi = \varphi(\sigma)\varphi(\pi)$
even	odd	odd	$\varphi(\sigma\pi) = \sigma\pi\tau = (\sigma)(\pi\tau) = \varphi(\sigma)\varphi(\pi)$
odd	even	odd	$\varphi(\sigma\pi) = \sigma\pi\tau = (\sigma\tau)(\pi) = \varphi(\sigma)\varphi(\pi)$
odd	odd	even	$\varphi(\sigma\pi) = \sigma\pi = \sigma\pi\tau^2 = (\sigma\tau)(\pi\tau) = \varphi(\sigma)\varphi(\pi)$

So  $\varphi$  is a homomorphism from  $S_{n-2}$  to  $A_n$ .

It is also true that  $\varphi$  is injective. Let  $\varphi(\sigma) = \sigma\tau^a = \pi\tau^b = \varphi(\pi)$ , for  $a, b \in \{0, 1\}$ . We reach a contradiction when  $a \neq b$ , because  $\sigma$  and  $\tau$  were permutations in  $S_{n-2}$ , so they fix  $n$  and  $n-1$ . It is not possible to write

$\sigma = \pi\tau$  or  $\pi = \sigma\tau$  as a result. Therefore,  $a = b$ , and by right cancellation  $\sigma = \pi$ . Hence  $\varphi$  is an injective homomorphism, which means  $\ker \varphi$  is the trivial subgroup. Then by the First Isomorphism Theorem,

$$\frac{S_{n-2}}{\ker \varphi} = \frac{S_{n-2}}{\{1\}} \cong S_{n-2} \cong \varphi(S_{n-2}) \leq A_n.$$

Hence for all  $n \geq 3$ ,  $A_n$  contains a subgroup isomorphic to  $S_{n-2}$ . □

5. (DF3.1.36) Prove that if  $G/Z(G)$  is cyclic then  $G$  is abelian.

*Proof.* Let  $G$  be a group such that  $G/Z(G)$  is cyclic as given. Then by definition,  $G/Z(G)$  has a generator. We may choose a representative  $x \in G$  so that  $xZ(G)$  is a generator for  $G/Z(G)$ . Then because the left cosets of  $Z(G)$  partition  $G$ , it follows that for any element  $g \in G$ ,  $g$  lies in one of the left cosets of  $Z(G)$ . In particular,  $g$  lies in  $gZ(G)$ , which is an element of  $G/Z(G)$ . Since  $xZ(G)$  generates  $G/Z(G)$ , there exists  $n \in \mathbb{Z}$  such that  $gZ(G) = (xZ(G))^n = x^nZ(G)$ . Thus there exists  $z_1, z_2 \in Z(G)$  such that  $gz_1 = x^n z_2$ , which by right multiplication we have that  $g = x^n z_2 z_1^{-1}$ . Let  $z = z_2 z_1^{-1}$ , and because  $Z(G)$  is a subgroup of  $G$ ,  $z_2 z_1^{-1} = z \in Z(G)$ .

Hence any element  $g \in G$  can be written as  $g = x^n z$ , where  $n \in \mathbb{Z}$  and  $z \in Z(G)$ . Then let  $a, b \in G$  so that  $a = x^j z_1$  and  $b = x^k z_2$  for  $j, k \in \mathbb{Z}$  and  $z_1, z_2 \in Z(G)$ . Due to exponent rules and the fact that  $z_1, z_2$  lie in the center of  $G$ ,

$$\begin{aligned} ab &= (x^j z_1)(x^k z_2) \\ &= (z_1 x^j)(x^k z_2) \\ &= z_1 (x^j x^k) z_2 \\ &= z_2 (x^k x^j) z_1 \\ &= (z_2 x^k)(x^j z_1) \\ &= (x^k z_2)(x^j z_1) = ba. \end{aligned}$$

Because  $a, b$  were arbitrary elements of  $G$  it follows that  $G$  is abelian. □