

CHAZ Florida Hurricane Hazard Map

Interactive Visualization of Tropical Cyclone Wind Hazard Data

Paul Fishwick

Claude Code

January 2026

Abstract

This report documents an interactive web-based visualization tool for exploring tropical cyclone wind hazard data for Florida. The tool displays exceedance intensity data from the Columbia HAZard (CHAZ) model, allowing users to compare historical baseline conditions (1995–2014) with mid-century (2041–2060) and late-century (2081–2100) climate projections under the SSP585 emissions scenario. Users can select from six different CMIP6 climate models or view a multi-model mean to explore uncertainty across model projections.

1 Introduction

Tropical cyclones pose significant risks to Florida’s coastal communities. Understanding how hurricane wind hazards may change under future climate scenarios is essential for long-term planning, infrastructure design, and risk assessment. This project provides an accessible, interactive tool for exploring CHAZ model output data.

The visualization is available online at: <https://metaphorx.github.io/CHAZHazard/>

2 Data Source

The underlying data comes from the CHAZ Hazard Maps dataset published on Dryad:

Meiler, S., et al. (2025). CHAZ Hazard Maps: Global coastal wind hazard maps from the CHAZ tropical cyclone model. Dryad. <https://doi.org/10.5061/dryad.qfttdz0vz>

The dataset provides exceedance intensity values—the wind speeds (in m/s) expected to be exceeded at various return periods (10, 25, 50, 100, 250, and 1000 years)—for coastal locations worldwide. For this visualization, we extracted 2,094 land-based points covering Florida.

3 The CHAZ Model

3.1 Overview

The Columbia HAZard model (CHAZ) is a statistical-dynamical downscaling model for estimating tropical cyclone hazard. Rather than predicting specific future hurricanes, CHAZ generates thousands of synthetic storms based on climate conditions from global climate models (GCMs), then aggregates statistics to estimate hazard metrics.

3.2 Model Components

CHAZ consists of three primary modules:

1. **Genesis Module:** Uses the Tropical Cyclone Genesis Index (TCGI) to estimate where and when storm precursors form based on environmental conditions.
2. **Track Module:** Employs a Beta-Advection Model (BAM) that moves storms forward using environmental steering flow from the climate model.
3. **Intensity Module:** Evolves storm intensity using an autoregressive model with deterministic forcing (based on potential intensity, wind shear, humidity) and stochastic elements.

3.3 Climate Model Input

The visualization includes data from six CMIP6 climate models, all run under the SSP585 scenario (high emissions, “business as usual”). Users can select any individual model or view a multi-model mean that averages across all six. Table 1 describes each available model.

Table 1: Climate models available in the visualization

Model	Institution	Country
CESM2	National Center for Atmospheric Research	USA
CNRM-CM6-1	Centre National de Recherches Météorologiques	France
EC-Earth3	EC-Earth Consortium	Europe
IPSL-CM6A-LR	Institut Pierre Simon Laplace	France
MIROC6	Multiple Japanese Institutions	Japan
UKESM1-0-LL	UK Met Office / NERC	UK
<i>Multi-Model Mean</i>	<i>Average of all 6 models</i>	—

3.4 About the CMIP6 Models

These models are all part of the Coupled Model Intercomparison Project Phase 6 (CMIP6), which provides the foundation for the IPCC Sixth Assessment Reports. They represent the cutting edge of climate simulation, moving beyond simple physics to include complex Earth System processes like the carbon cycle and vegetation dynamics.

CESM2 (Community Earth System Model v2) from the National Center for Atmospheric Research (NCAR), USA, is a highly flexible, community-driven model known for its advanced representation of land-ice (ice sheets) and its CAM6 atmospheric component. It is often used as a benchmark due to its extensive documentation and broad scientific community.

CNRM-CM6-1 from CNRM (Centre National de Recherches Météorologiques), France, uses the ARPEGE-Climat atmospheric model with significant improvements in cloud representation, convection, and surface-atmosphere interactions via the SURFEX system. It is highly efficient and frequently used for High Resolution Model Intercomparison (HighResMIP).

EC-Earth3 from a European consortium of 27 research institutes across 10 countries is based on the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecasting System (IFS). Uniquely, it uses the same core as the world’s leading weather prediction model, bridging short-term weather forecasting and long-term climate projection.

IPSL-CM6A-LR from Institut Pierre-Simon Laplace (IPSL), France, is a complete Earth System Model including the LMDz atmospheric model and ORCHIDEE land surface model. “LR” indicates Low Resolution, enabling very long multi-century simulations. It excels at simulating the carbon cycle and biosphere-atmosphere interactions.

MIROC6 (Model for Interdisciplinary Research on Climate) from JAMSTEC, University of Tokyo, and NIES in Japan focuses on improving internal variability—natural climate swings like El Niño. It features improved stratosphere and shallow convection representation, with particular strength in simulating the Madden-Julian Oscillation (MJO) and tropical rainfall patterns.

UKESM1-0-LL from the Met Office Hadley Centre, UK, is a full-complexity Earth System Model that takes the physical core of HadGEM3 and adds fully interactive chemistry, aerosols, and marine/terrestrial biogeochemistry. It is one of the most comprehensive models in CMIP6, used to explore tipping points and complex climate feedbacks.

Table 2 compares the core components of each model.

Table 2: Core components of the CMIP6 climate models

Model	Atmosphere	Land Surface	Ocean
CESM2	CAM6	CLM5	POP2
CNRM-CM6-1	ARPEGE-Climat v6.3	ISBA-CTrip	NEMO 3.6
EC-Earth3	IFS	HTESSEL	NEMO 3.6
IPSL-CM6A-LR	LMDz6	ORCHIDEE	NEMO 3.6
MIROC6	CCSR-NIES AGCM	MATSIRO	COCO 4.9
UKESM1-0-LL	HadGEM3-GC3.1	JULES	NEMO 3.6

3.5 Why Multiple Models?

Different climate models make different assumptions about physical processes, leading to a range of projections. By including all six models, users can:

- Assess **model uncertainty** by comparing projections across models
- Identify **robust signals** that appear consistently across models
- Use the **multi-model mean** as a consensus estimate that smooths individual model biases

4 Time Periods and Climate Projections

The visualization includes three time periods, as shown in Table 3.

Table 3: Time periods available in the visualization

Period	Years	Description
Historical (base)	1995–2014	Baseline climate conditions
Mid-Century (fut1)	2041–2060	Near-future projection
Late-Century (fut2)	2081–2100	End-of-century projection

4.1 How Future Projections Are Generated

For each time period, CHAZ:

1. Reads climate conditions (SST, winds, humidity, etc.) from the GCM
2. Calculates genesis probability using TCGI
3. Seeds thousands of synthetic storms based on genesis rates
4. Simulates each storm's track using steering winds
5. Evolves intensity based on environmental conditions
6. Aggregates statistics across 80 ensemble members (~1,600 synthetic years)

Key climate variables that change between periods include:

- **Sea Surface Temperature (SST):** Warmer oceans provide more energy for storms
- **Potential Intensity (PI):** Maximum theoretical storm strength increases
- **Vertical Wind Shear:** Affects storm formation and intensification
- **Mid-level Humidity:** Influences moisture availability
- **Atmospheric Circulation:** Alters steering currents and track patterns

5 Visualization Features

The interactive map includes the following features:

- **Climate Model Selector:** Choose from six individual CMIP6 models or the multi-model mean (CESM2 is the default)
- **Time Period Selector:** Switch between historical, mid-century, and late-century projections
- **Return Period Selector:** Choose from 10, 25, 50, 100, 250, or 1000-year return periods
- **Color-Coded Markers:** Circle markers colored by wind speed intensity
- **Hover Tooltips:** Display coordinates, wind speed (m/s, km/h, mph), hurricane category, and all return period values
- **Legend:** Wind speed scale with three unit systems

6 Results

Figure 1 shows the 250-year return period wind speeds for the historical baseline (1995–2014). Figure 2 shows the same metric for the late-century projection (2081–2100).




Figure 1: 250-year return period wind speeds for Florida under historical climate conditions (1995–2014). Colors indicate wind intensity from blue (lower) to red (higher).




Figure 2: 250-year return period wind speeds for Florida under late-century climate projections (2081–2100, SSP585). Comparison with Figure 1 reveals changes in hazard distribution.

6.1 Key Observations

Comparing the historical and future projections reveals several patterns consistent with climate change projections for tropical cyclones:

- Overall wind intensities show increases in many coastal areas
- The spatial pattern of highest hazard remains concentrated along the southeast coast and Keys
- Some inland areas show modest changes in projected hazard

7 Technical Implementation

The visualization is built using:

- **Leaflet.js**: Open-source JavaScript library for interactive maps
- **HTML/CSS/JavaScript**: Standard web technologies for broad compatibility

- **GitHub Pages:** Free hosting for static web content

Data is embedded directly in the HTML file to avoid CORS restrictions and enable offline use. The approximately 2,094 land points are rendered efficiently using Leaflet's CircleMarker class.

8 Data Processing

The original CHAZ data covers global coastal areas with over 1 million points per model. For this visualization:

1. Florida points were extracted using a bounding box (lat: 24–31°N, lon: 88–79.5°W)
2. Ocean points were filtered out using a Florida land polygon
3. Data for all six climate models and three time periods was extracted
4. The multi-model mean was computed by averaging wind speeds across all six models at each grid point
5. Wind speeds were rounded to one decimal place to reduce file size
6. All data (7 model options \times 3 time periods \times 2,094 points) was embedded in the HTML

9 Acknowledgments

This visualization was developed by Paul Fishwick with assistance from Claude Code. The underlying CHAZ model was developed by the Columbia University team: Drs. Suzana J. Camargo, Chia-Ying Lee, Michael K. Tippett, and Adam H. Sobel.

10 License

The visualization code is released under the MIT License. The underlying CHAZ hazard data is released under Creative Commons Zero (CC0).

11 References

1. Lee, C.-Y., Tippett, M.K., Sobel, A.H., and Camargo, S.J. (2018). An environmentally forced tropical cyclone hazard model. *Journal of Advances in Modeling Earth Systems*, 10, 233–241.
2. Lee, C.-Y., Camargo, S.J., Sobel, A.H., and Tippett, M.K. (2020). Statistical-dynamical downscaling projections of tropical cyclone activity in a warming climate. *Journal of Climate*, 33, 4815–4834.
3. CHAZ GitHub Repository: <https://github.com/cl3225/CHAZ>