

Geomorphological Experiments for Amelia Island

An interactive simulation of barrier-island formation,
Holocene accretion, and Pleistocene welding

Paul Fishwick and Claude Code

May 2026

Abstract

Amelia Island is a barrier island on the northeastern coast of Florida, at the southern end of the Georgia Bight. It exhibits the classic “double island” structure of the Bight: an older Pleistocene core (Silver Bluff terrace, ~ 125 ka) overlain on its seaward side by a Holocene wedge that welded onto the core $\sim 5,000$ years ago, trapping the salt marsh now known as Egans Creek between them. Since welding, the Atlantic face has prograded seaward, preserving a set of nested beach ridges most clearly visible at Fort Clinch State Park on the island’s north end. This document specifies a set of computational experiments—an interactive web simulation, paired with browser-driven verification—that reconstruct the island’s coastline through time. We ground the simulation in published sea-level data for northeastern Florida [1], in the formation theories of de Beaumont, Gilbert, McGee, and Hoyt [4], and in the narrative provided in lectures by the Conserve Nassau organization [10, 11, 12].

Contents

1	Motivation	2
2	Geomorphic background	2
2.1	Barrier islands as a class of landform	2
2.2	Two regimes: transgressive and regressive	2
2.3	From sandbar to vegetated barrier	2
2.4	The transgressive sandbar—what it looks like	3
2.5	The Georgia Bight “double island” pattern	5
2.6	Florida marine terraces	5
3	Sea-level history used in the simulation	5
4	Narrative arc of Amelia Island	5
5	Simulation design	7
5.1	Goal	7
5.2	Geometry	7
5.3	Cross-section view	7
5.4	Architecture	8

6	Verification with Selenium	8
7	User guide	8
7.1	Launching the simulation	8
7.2	The map view	8
7.3	The time slider	9
7.4	The era panel	9
7.5	The cross-section panel	9
8	Results	9
9	Reproducibility	9
10	Acknowledgements	13

1 Motivation

The question that motivated this work is direct: *how did Amelia Island grow into its present shape, and can that growth be made visible?*

The geomorphic story of the island is well established but is rarely shown as a process. Most published material is text-and-diagram. A barrier island, however, is a moving system: sea level oscillates by hundreds of feet over glacial cycles, shorelines migrate tens of miles, and a sandbar a few miles offshore approaches the mainland over millennia and finally welds onto it. These are kinematic phenomena that benefit from a kinematic representation.

We propose a browser-based simulation in which the user drags a slider through $\sim 125,000$ years of coastline change and sees the island form. The simulation is anchored on real geospatial data (modern shoreline from OpenStreetMap, terrain signatures from USGS digital elevation models) and on real sea-level curves [1, 2, 3].

2 Geomorphic background

2.1 Barrier islands as a class of landform

A barrier island is a long, narrow accumulation of sand parallel to a mainland coast, separated from it by a lagoon, estuary, or marsh. Three classical theories have been offered for how such islands form:

- **De Beaumont** (1845): offshore submarine bars built by shoaling waves emerge above sea level. Flume experiments later showed bars cannot grow above sea level by this mechanism alone, so this theory is now considered a partial contributor.
- **Gilbert** (1885, “spit-accretion”): longshore drift builds a sandy spit attached to the mainland; storms breach the spit, isolating an island. This mechanism is observed in some modern systems.
- **McGee** (1890, “submergence”): rising sea level drowns mainland beach ridges and dune fields, isolating them as islands with a new lagoon behind. Hoyt (1967, [4]) supported this view with stratigraphic evidence—specifically, the absence of shoreface sediment below early barriers, which would be expected if the bars had emerged from below.

Modern process-based work [5] treats these as endmembers of a self-organizing system whose behavior depends on three controls: the rate of relative sea-level rise (RSLR), the supply of sand, and the back-barrier accommodation space.

2.2 Two regimes: transgressive and regressive

Once formed, a barrier can be in one of two regimes (Table 1).

Modeling [5] identifies a critical threshold near 2–5 mm/yr RSLR: above it, barriers are typically transgressive; below it, they can stabilize and prograde. This threshold is consequential for Amelia.

2.3 From sandbar to vegetated barrier

A barrier island begins as a bare sandbar—a submerged or intertidal accumulation of sediment where waves can no longer transport sand further offshore. The transition to a stable, above-sea-

Table 1: The two regimes of a barrier island, after [5].

Regime	Drivers	Signature
Transgressive (“rollover”)	RSLR high, or sediment-starved; overwash moves sand landward across the island	Washover fans on lagoon side; landward-migrating shoreline; ridges not preserved
Regressive (“progradational”)	Sediment supply exceeds accommodation; new sand builds seaward of older shoreline	Beach-ridge plain (the nested arcs at Fort Clinch); seaward-younging

level barrier requires vegetation. The sequence is roughly:

1. **Submerged bar:** sediment accumulating offshore, below mean low water.
2. **Emergent intertidal sandbar:** highest crests exposed at low tide, bare or sparsely colonized by salt-tolerant pioneers (*Uniola paniculata*, *Spartina patens*).
3. **Vegetated barrier:** dune grasses slow the wind, trap windblown sand, and build foredunes. The foredunes shelter back-dune shrubs, and the shrubs in turn shelter trees.
4. **Mature barrier with maritime forest:** oaks, palmettos, and a closed canopy on stabilized terrain.

The vegetation feedback is load-bearing for the system. Without plants, a bare sandbar cannot retain sand against wind erosion, and it disperses. The older Pleistocene core of Amelia exhibits the mature end of this sequence: 100,000 years of subaerial exposure has produced deep soils, rounded slopes, and a maritime oak hammock. The younger Holocene wedge, by contrast, shows steeper slopes, less mature soils, and a less stratified plant community—it has had only 5,000 years above sea level.

For our reconstruction, the implication is that the offshore Holocene sandbar between 8 ka and 5 ka was likely bare or sparsely vegetated. After welding, the established vegetation network of the older Pleistocene island could spread laterally onto the new wedge, accelerating its stabilization.

2.4 The transgressive sandbar—what it looks like

The offshore Holocene sandbar that approaches Amelia between ~ 8 ka and ~ 5 ka deserves its own treatment, because it is geomorphically the most interesting (and visually the most distinctive) feature in the animation. It is a *proto-barrier island*: a long, narrow ridge of sand parallel to the existing shoreline, separated from the mainland by a developing lagoon, intermittently emergent at low tide.

Modern analogs. Many present-day barrier islands look exactly like Amelia’s offshore sandbar did at 8–5 ka:

- **Outer Banks, North Carolina** (~ 320 km long, 0.5–3 km wide) — the textbook transgressive barrier, still migrating landward.
- **Padre Island, Texas** (~ 182 km long, 0.5–5 km wide) — the longest barrier island in the world; same long-narrow geometry.

- **Fire Island, New York** (~ 50 km long, ~ 0.5 km wide) — another active transgressive barrier.
- **Cumberland Island, Georgia**, immediately north of Amelia — the same Georgia Bight “double island” as Amelia, just slightly older. The Holocene wedge welded onto its Pleistocene core in the same process described here.

Why long and narrow. The form arises from three balanced controls:

- **Length** is set by along-shore continuity of sediment supply. Longshore drift moves sand parallel to the coast and accumulates it into a continuous ridge wherever sediment supply is positive. For Amelia’s ancestor bar, the supply came primarily from the St. Marys River and from sand reworked across Cumberland Sound.
- **Width** is constrained by the equilibrium between wave erosion on the seaward face and sand-supply deposition. During the active transgressive phase, this equilibrium width is typically 100–500 m; modern transgressive barriers stay at this scale until sea-level rise slows enough for them to stabilize and widen.
- **Curvature** mirrors the existing shoreface. The bar forms parallel to whatever coastline already exists, so its plan-view shape follows the curve of Amelia’s modern Atlantic shore (which is, in turn, inherited from the Pleistocene core’s geometry).

The landward migration. While sea level rose rapidly (rates above ~ 5 mm/yr), the bar could not stabilize at any fixed offshore position. Storm overwash carried sand from the seaward face across the bar to the lagoon side, depositing it as washover fans. Over centuries, this net landward sand transport—combined with continued retreat of the equilibrium profile—caused the entire bar to “roll over” shoreward. Estimated migration rate during the slowing- RSLR mid-Holocene is approximately 1 km per millennium, consistent with Frank Hopf’s reference statement that the bar “started out about four or five miles out from where we are today on the coastline” at ~ 8 ka [11].

Visualization choices. In the simulation, the bar appears as a pale dashed polygon east of Amelia, in two discrete snapshots (8 ka and 6 ka). Several simplifications apply:

- **The dashed outline** signals that the bar is an inferred, migrating feature rather than a fixed landform with a known boundary.
- **Width** is rendered at ~ 300 m for visibility; real transgressive bars were probably 50–200 m wide at this stage and widening as they slowed.
- **Only the emergent crest** is shown; in reality the bar has a submerged base extending 1–3 km further seaward at depth.
- **Migration** is rendered as two discrete positions rather than a continuous motion—a fidelity-vs-clarity tradeoff. The intermediate position is implicit; the user reading the slider can mentally interpolate.

After welding at ~ 5 ka, the bar polygon disappears from the animation and the Holocene wedge polygon takes over as the bar’s permanent successor on the welded island.

2.5 The Georgia Bight “double island” pattern

Along the Atlantic coast of Georgia and northeastern Florida, the Holocene barrier islands did not form on a virgin shelf. They formed offshore of an older set of Pleistocene barriers that had survived as elevated remnants. As sea level rose through the Holocene, the offshore Holocene bars migrated landward and *welded* onto the Pleistocene cores. Hoyt and Hails [4] identified six major Pleistocene shoreline complexes in coastal Georgia, each a former barrier-lagoon system, now overlain on its seaward side by a Holocene wedge. This pattern produces the double-island islands of the Bight: Hilton Head, Tybee, Wassaw, Ossabaw, St. Catherines, Sapelo, the St. Simons group, Jekyll, Cumberland, and Amelia at the southern end [6].

2.6 Florida marine terraces

The Pleistocene core of Amelia sits on the Silver Bluff terrace, formed during the Sangamonian interglacial (Marine Isotope Stage 5e) when sea level stood ~ 6 m above present [7]. Florida exposes a sequence of older terraces stepping up in elevation and back in age: Pamlico, Talbot, Penholoway, and Wicomico. Only the Silver Bluff terrace is relevant to Amelia.

3 Sea-level history used in the simulation

The most directly relevant sea-level reconstruction for Amelia is Hawkes et al. [1], which produced 25 sea-level index points spanning 8.0 ka to 2.0 ka from salt-marsh foraminifera at sites in northeastern Florida. Three findings from that work shape the simulation:

1. Relative sea level rose ~ 5.7 m in northeastern Florida over the last 8.0 ka.
2. There is **no Holocene highstand** at this latitude—the rise has been monotonic. This contrasts with the central US Atlantic coast and the Caribbean, which show mid-Holocene highstands.
3. The pattern reflects the collapse of the Laurentide Ice Sheet’s proglacial forebulge, which produces ongoing subsidence at this latitude.

We extend the curve back to 125 ka by combining [1] with the Caribbean Holocene curve of Toscano and Macintyre [3], the Engelhart and Horton US Atlantic database [2] for the mid- Holocene tail, and Lambeck et al. for the Last Glacial Maximum lowstand. The resulting target curve is summarized in Table 2.

4 Narrative arc of Amelia Island

The reconstruction proceeds in six phases.

Phase 0: Pleistocene core forms (~ 125 ka). During the Sangamonian highstand, sea level stood ~ 6 m above present and a barrier island formed on the Silver Bluff terrace at the location of today’s western Amelia. Behind it: lagoon and salt marsh. The same system that produced neighboring Cumberland Island.

Phase 1: Lowstand exposure (125–20 ka). Sea level fell ~ 125 m through the Wisconsinan glaciation. The Pleistocene Amelia stood as an inland hill, with the shoreline retreating to a Last

Table 2: Sea-level snapshots used by the simulation. Negative values indicate sea level below present. Sources are cited where each value is anchored.

Years BP	RSL (m)	Phase / source
125,000	+6	Eemian (MIS 5e) highstand; Silver Bluff terrace
80,000	−20	MIS 5a/4 transition
20,000	−120	Last Glacial Maximum
14,500	−90	onset of Meltwater Pulse 1A
13,500	−72	end of MWP-1A (~ 18 m rise in 500 yr)
8,000	−5.7	Hawkes et al. 2016 anchor
5,000	−2.0	Welding event (slow RSLR, ~ 1 mm/yr)
2,000	−0.5	Late Holocene progradation
150	−0.2	Pre-jetty (1880 CE)
0	0.0	Present (2026 CE)

Glacial Maximum position roughly 130 km east of today’s coast, near the -120 m bathymetric contour on the continental shelf. During this $\sim 100,000$ year exposure, the Pleistocene sands matured: deeper soils formed, slopes rounded, oak-hammock vegetation established.

Phase 2: Holocene transgression (20–8 ka). As the Laurentide and Eurasian ice sheets retreated, sea level rose rapidly, particularly during Meltwater Pulse 1A (14.7–13.5 ka, ~ 36 mm/yr). RSLR this fast keeps barriers in the transgressive regime: ephemeral sandbars formed, drowned, and reformed at landward positions on the migrating shoreface. None were preserved in their original locations.

Phase 3: The approach (8–5 ka). By 8 ka, sea level was within ~ 6 m of present and rising more slowly (~ 2 – 3 mm/yr per [1]). A persistent Holocene sandbar formed roughly four miles east of the Pleistocene core. The shallow water between the sandbar and the older island became a protected lagoon. Mud—which will not settle out of suspension on its own—accumulated as oysters and clams filtered it from the water and deposited it as faecal pellets. Salt marsh built upward on this biogenic mud. The future Egans Creek wetland began to form.

Phase 4: Welding (~ 5 ka). RSLR slowed below the stabilization threshold (~ 2 mm/yr). The Holocene sandbar stopped retreating, grew vertically and laterally, and finally welded onto the Pleistocene island—first at the south end of Amelia, then propagating north. The Egans Creek marsh was sealed in between, becoming a relict back-barrier lagoon. This is the most visually distinctive event in the simulation.

Phase 5: Progradation (5 ka–1880 CE). With slow RSLR, abundant longshore sand from the St. Marys River to the north, and the welded island as a fixed anchor, the Atlantic face of Amelia began to build seaward. Successive shoreline positions are preserved as beach ridges, each separated by 80–150 years on comparable Florida systems [8, 9]. At Fort Clinch on the north end, the curving plan form of the island favored maximum accretion, producing the nested arc-shaped ridges visible from the air and on USGS topography.

Phase 6: Human modification (1880–present). The St. Marys Entrance jetty, completed in 1881 and extended afterward, intercepted the southbound longshore sand transport at the north end of Amelia. The result was an additional fillet of accretion—the “post-jetty dunes” visible on the seaward side of the historical shoreline at Fort Clinch State Park—superimposed on the natural pattern over 150 years rather than the natural rate of 80–150 years per ridge. The Nassau County Shore Protection Project (USACE Jacksonville District, with Olsen Associates) now manages periodic beach nourishment on the developed central portion of Amelia.

5 Simulation design

5.1 Goal

The simulation is a single-page web application. The user opens `index.html` in a browser. A Leaflet [14] map centered on Amelia Island shows the present-day coastline on an OpenStreetMap basemap. A horizontal slider, ranging from 125 ka BP to the present, is positioned below the map. Dragging the slider re-renders the map with the polygons appropriate to that time slice: the Pleistocene core (when above sea level), the Holocene wedge in its current state (absent, offshore as a migrating bar, welded, or with progradational ridges), and the salt marshes. A side panel displays a one-paragraph caption summarizing the era, paraphrased from the source lectures and the published literature.

5.2 Geometry

Geometry is built from three sources:

1. **Modern island coastline:** queried from OpenStreetMap via the Overpass API and exported to GeoJSON. This is real public-domain data.
2. **Pleistocene/Holocene boundary on the island:** traced once from the USGS 3DEP digital elevation model. The Pleistocene core has the rounded, soil- mature signature of an emerged terrace; the Holocene wedge has the corrugated ridge-and-swale signature of progradational accretion. The boundary approximately follows Egans Creek through the central island.
3. **Beach ridges at Fort Clinch:** digitized from satellite imagery and DEM hillshade. We digitize 8 major ridges, consistent with the major arcs visible in the field.

Pre-welding paleo-coastlines (12 ka, 8 ka, 5 ka) are reconstructed by offset from the modern shoreline using the migration distances cited in the source lectures (3–5 miles offshore at 12 ka, approaching at ~1 mile per millennium). These are labeled in the UI as interpretive reconstructions, not measured paleoshorelines.

5.3 Cross-section view

A secondary panel renders the modern cross-section described by the source lectures (Plantation, Little Nana, and Big Nana profiles): a young Holocene dune on the Atlantic face, the Egans Creek marsh in the center, the older Pleistocene core to the west, and the back-barrier marsh on the mainland side. The panel animates the accretion of the cross-section through time, illustrating the “two piles of sand on a bed of mud” description Frank Hopf uses.

5.4 Architecture

- `index.html` — single page, no build step.
- `viewer.js` — loads snapshot GeoJSONs, interpolates between them as the slider moves, manages layer styling.
- `style.css` — Pleistocene shown in a muted earth-tone; Holocene sand in a pale ridge-and-swale pattern; salt marsh in green; sea in blue.
- `data/` — one GeoJSON per snapshot, plus `sea_level.json`.
- `scripts/` — Python utilities for data preparation and Selenium- based verification.

6 Verification with Selenium

All HTML web tests are written in Selenium [15], per project standard. The test suite (`tests/auto/test_viewer.py`) launches a headless Chrome browser, loads the application from a local HTTP server, waits for the Leaflet map to fully initialize, and exercises the time slider through each defined snapshot. For each snapshot, the test:

1. Sets the slider to the snapshot’s target year.
2. Waits for the layer transition to complete and tiles to load.
3. Captures a full-page screenshot to `docs/fig/snapshot_N.png`.
4. Asserts that the expected layers are present in the DOM (`.layer-pleistocene`, `.layer-holocene`, `.layer-marsh`, etc. as appropriate to the era).

The same captured screenshots are referenced by this document via `\includegraphics`. The verification suite therefore serves two roles: it tests the application, and it generates the figures embedded here. This document is rebuilt by running the test suite, then `pdflatex` on this file.

7 User guide

7.1 Launching the simulation

The simulation is a single static HTML file with no server-side dependencies. From the project root, launch the bundled `start` script:

```
./start
```

The script checks whether port 8765 is already in use (and stops a stale process if so), starts a local HTTP server, and opens `http://localhost:8765/index.html` in the default browser. To stop the server later:

```
./stop
```

7.2 The map view

On load, the application centers on Amelia Island at the present-day timestep. The OpenStreetMap basemap shows the modern coastline, with the simulation’s geologic layers drawn on top: the Pleistocene core in muted earth tone, the Holocene wedge in pale sand, the Egans Creek marsh in

green, the Atlantic and Amelia River in blue. Standard Leaflet pan and zoom controls are available; the geologic layers re-render at every zoom level.

7.3 The time slider

A horizontal slider below the map ranges from 125,000 years BP at the left to the present (2026 CE) at the right. Dragging the slider re-renders the geologic layers to reflect the coastline at that time. The slider is logarithmic in years BP, so most of its travel is in the Holocene, where the welding event and beach-ridge progradation occur.

A play button below the slider animates the simulation forward from the current position at one of three speeds (1×, 5×, 20×). A reset button returns the slider to 125 ka.

7.4 The era panel

To the right of the map, an information panel updates as the slider moves. It shows the current era name, the year BP, the relative sea level (in meters versus today), and a one-paragraph caption describing what is happening at that moment. Caption text is paraphrased from the source lectures [10, 11, 12] and the published literature cited above.

7.5 The cross-section panel

A toggle in the upper-right opens a secondary panel showing the modern cross-section of the island at the Plantation profile, after the cross-sections in [10, 11, 12]. From west to east this shows: the Amelia River, the mainland salt marsh, the Pleistocene core, the Egans Creek marsh, the Holocene wedge with its beach ridges, the modern foredune, and the Atlantic Ocean. As the time slider moves, the cross-section animates to show how each component built up. The cross-section caption identifies the “two piles of sand on a bed of mud” structure that Frank Hopf emphasizes.

8 Results

The figures below are captured directly by the Selenium suite as it exercises the viewer through each defined snapshot. Each figure shows the full application interface: the Leaflet map (left, top), the cross-section panel (left, bottom), and the era panel (right) with current year, relative sea level, and the narrative caption.

9 Reproducibility

The figures above are not hand-curated. They are produced by running the Selenium suite, which drives the viewer through every defined snapshot, asserts that the expected layer kinds are rendered at each timestep, and saves a screenshot per snapshot. To reproduce:

```
python3 -m venv venv
source venv/bin/activate
pip install selenium Pillow youtube-transcript-api yt-dlp
python scripts/fetch_osm.py
python scripts/build_snapshots.py
python tests/auto/test_viewer.py
pdflatex docs/geomorphological-experiments.tex
```




Figure 1: Eemian highstand, 125 ka. The Pleistocene core (Silver Bluff terrace) is forming under a sea level ~6 m above today's. The cross-section shows the future-Amelia footprint as a single body above the dashed sea-level line; everything seaward of it will form later.




Figure 2: Last Glacial Maximum, 20 ka. Sea level -120 m. The red dashed hint line at the right edge of the map marks the LGM Atlantic shoreline ~130 km east. The Pleistocene Amelia stands as an inland hill, far from any coast.

The Selenium suite exits with status 0 when all snapshots render correctly. Failures are reported as “FAIL” lines and logged to `tests/auto/test_viewer.log`. The figures in `docs/fig/` are over-




Figure 3: Holocene transgression, 8 ka. Sea level -5.7 m (Hawkes et al. 2016 anchor). A persistent Holocene sandbar (pale dashed polygon to the east of the island) has formed ~ 4 mi east of the Pleistocene core. In the cross-section: the bar appears offshore (right), and the Egans Creek marsh begins to fill the lagoon between bar and core.




Figure 4: The welding event, 5 ka. The Holocene sandbar has welded onto the Pleistocene core. Egans Creek marsh is permanently trapped between them. This is the visual climax of the simulation: the moment Amelia becomes today's double-island.




Figure 5: Mid-Holocene progradation, 3 ka. The Atlantic face begins building seaward. Beach ridges appear as small arcs near Fort Clinch (north end of the island). Sea level -1 m ; sea-level rise has dropped well below the $\sim 2\text{ mm/yr}$ stabilization threshold.




Figure 6: Present, 2026 CE. All eight modeled beach ridges are present at Fort Clinch, including the post-jetty fillet added since 1881. The cross-section shows the full “two piles of sand on a bed of mud” structure that Frank Hopf emphasizes: Pleistocene core on the west, Egans Creek marsh in the middle, Holocene wedge with beach ridges on the east.

written each run, so this document always reflects the current state of the viewer.

10 Acknowledgements

The geomorphic narrative summarized here draws extensively from a series of public lectures given by Frank Hopf, presented through the [Conserve Nassau Florida](#) organization [13]. Three of those lectures provided the narrative arc for the simulation:

- *Hopf: A Fragile Barrier Island* (64 min) [10] — the master overview of formation, sea-level history, and present-day fragility.
- *Dunes: Not Just a Pile of Sand — Part 1: Pangaea to Present* (28 min) [11] — the 400-million-year geologic context and specific spatial references for Amelia’s Pleistocene and Holocene history.
- *Dunes: Not Just a Pile of Sand — Part 2: Our Holocene Dunes* (33 min) [12] — Holocene dune development and the post-jetty modifications since 1881.

Transcripts were extracted programmatically from the YouTube auto-captions, then mined for dates, distances, ridge-set descriptions, and cross-section locations. Transcript material is paraphrased here under fair use for non-commercial educational illustration. Direct quotations are clearly marked. The original recordings remain the canonical source.

We also acknowledge the underlying peer-reviewed work that frames Amelia Island’s geomorphology, particularly Hawkes et al. [1] for the NE Florida sea-level reconstruction, Hoyt and Hails [4] for the Georgia Bight stratigraphic framework, and Mariotti [5] for the modern process-based understanding of barrier-island stabilization.

References

- [1] Hawkes, A. D., Kemp, A. C., Donnelly, J. P., Horton, B. P., Peltier, W. R., Cahill, N., et al. (2016). Relative sea-level change in northeastern Florida (USA) during the last ~ 8.0 ka. *Quaternary Science Reviews*, 142, 90–101. <https://www2.whoi.edu/site/coastalgroup/wp-content/uploads/sites/139/2021/07/Hawkes-2016-Relative-sea-level-change-in-northeastern-Florida-USA-during-the-last-8.0-KA.pdf>
- [2] Engelhart, S. E., and Horton, B. P. (2012). Holocene sea-level database for the Atlantic coast of the United States. *Quaternary Science Reviews*, 54, 12–25.
- [3] Toscano, M. A., and Macintyre, I. G. (2003). Corrected western Atlantic sea-level curve for the last 11,000 years based on calibrated ^{14}C dates from *Acropora palmata* framework and intertidal mangrove peat. *Coral Reefs*, 22, 257–270.
- [4] Hoyt, J. H., and Hails, J. R. (1967). Pleistocene shoreline sediments in coastal Georgia: deposition and modification. *Science*, 155(3769), 1541–1543. <https://www.science.org/doi/10.1126/science.155.3769.1541>
- [5] Mariotti, G. (2021). Self-organization of coastal barrier systems during the Holocene. *Journal of Geophysical Research: Earth Surface*, 126. <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2020JF005867>

- [6] Davis, R. A., Jr., and FitzGerald, D. M. (1994). The Georgia Bight Barrier System. In R. A. Davis Jr. (Ed.), *Geology of Holocene Barrier Island Systems*. Springer-Verlag. https://link.springer.com/chapter/10.1007/978-3-642-78360-9_7
- [7] Wikipedia contributors. *Geology of Florida*. https://en.wikipedia.org/wiki/Geology_of_Florida
- [8] Rink, W. J., and Forrest, B. (2005). Dating evidence for the accretion history of beach ridges on Cape Canaveral and Merritt Island, Florida, USA. *Journal of Coastal Research*, 21(5), 1000–1008.
- [9] Forrest, B. M., Rink, W. J., et al. (2008). New quartz OSL ages for beach ridges on the St. Vincent Island Holocene strandplain, Florida, USA. *Journal of Coastal Research*, Special Issue 24(SP1).
- [10] Frank Hopf (Conserve Nassau Florida). *Hopf: A Fragile Barrier Island* [public lecture, video]. YouTube, May 2026 (accessed). 64 min. <https://www.youtube.com/watch?v=SNysbbhGiDY>
- [11] Frank Hopf (Conserve Nassau Florida). *Dunes: Not Just a Pile of Sand — Part 1: Pangaea to Present* [public lecture, video]. YouTube, accessed May 2026. 28 min. <https://www.youtube.com/watch?v=nUIkpNE7k5E>
- [12] Frank Hopf (Conserve Nassau Florida). *Dunes: Not Just a Pile of Sand — Part 2: Our Holocene Dunes* [public lecture, video]. YouTube, accessed May 2026. 33 min. <https://www.youtube.com/watch?v=JvtDEisEI4c>
- [13] Conserve Nassau Florida. YouTube channel. <https://www.youtube.com/@conservenassauflorida>
- [14] Leaflet — an open-source JavaScript library for mobile-friendly interactive maps. <https://leafletjs.com>
- [15] Selenium WebDriver. <https://www.selenium.dev>