Methods Camp

UT Austin, Department of Government

Andrés Cruz and Matt Martin

2023-08-10

Table of contents

CI	ass so	hedule	6
	Desc	ription	6
	Cou	ese outline	6
	Con	act info	8
Se	etup		9
	Insta	ulling R and RStudio	9
	Setti	ng up for Methods Camp	11
1	Intro	o to R	1 2
	1.1	Objects	12
	1.2	Vectors and functions	13
	1.3	Data frames and lists	16
	1.4	Packages	18
2	Tidy	data analysis I	20
	2.1	•	20
	2.2	9	22
		• •	22
		9	26
			27
			28
		_	30
			33
		9	34
	2.3		34
3	Mat	rices	10
	3.1	Introduction	40
		3.1.1 Scalars	40
		3.1.2 Vectors	40
	3.2	Operators	41
		3.2.1 Summation	41
		3.2.2 Product	12
		3.2.3 Basics	43
		3.2.4 Structure	1 /

	3.3	Matrix operations
		3.3.1 Addition and subtraction
		3.3.2 Scalar multiplication
		3.3.3 Matrix multiplication
		3.3.4 Multiplication steps
	3.4	Properties
	3.5	Special matrices
	3.6	Transpose
	3.7	Inverse
4	Tidy	data analysis II 57
_		ctions and loops 58
5		•
	5.1	Basics
		5.1.1 What is a function?
		5.1.2 Function machine
	- 0	5.1.3 Visualization
	5.2	Types of functions
		5.2.1 Linear functions
		5.2.2 Quadratic
		5.2.3 Cubic
		5.2.4 Polynomial
		5.2.5 Exponential
		5.2.6 Trigonometric functions
	5.3	Logarithms and exponents
		5.3.1 Logarithms
		5.3.2 Relationships
		5.3.3 Basic rules
		5.3.4 Natural logarithms
		5.3.5 Definition of e
	5.4	Functions of functions
		5.4.1 Basics
		5.4.2 PMF, PDF, and CDF
6	Calc	ulus 65
U	6.1	Theory
	0.1	6.1.1 Derviative
		6.1.2 Calculating derivatives
		1
	6.0	
	6.2	Advanced rules
		n / L Product rule 60

		6.2.2	Quotient rule	70
		6.2.3	Chain rule	70
		6.2.4	Second derivative	71
	6.3	Differe	ntiable and Continuous Functions	72
		6.3.1	When is f not differentiable?	73
	6.4	Extren	na and optimization	73
		6.4.1	Extrema	73
		6.4.2	Identifying extrema	74
		6.4.3	Minimum or maximum?	74
		6.4.4	Second derivatives	75
		6.4.5	Local vs. Global Extrema	75
	6.5	Partial	derivatives	76
		6.5.1	Application	77
	6.6	Integra	als	78
		6.6.1	Area under a curve	78
		6.6.2	Integrals as summation	78
		6.6.3	Definite integrals	78
		6.6.4	Indefinite integrals	79
		6.6.5	Solving definite integrals	80
		6.6.6	Rules of integration	80
		6.6.7	More rules	80
		6.6.8	Solving the problem	81
		6.6.9	Integration by parts	82
7		pability		83
	7.1		is probability?	83
	7.2		gorov's axioms	83
	7.3		definitions	84
	7.4		te probability	84
		7.4.1	Probability Mass Function (PMF)	85
		7.4.2	Discrete distribution	85
	7.5		uous probability	86
		7.5.1	Basics	86
			Probability Density Function (PDF)	86
	7.6		ative Density Function (CDF)	86
		7.6.1	Discrete	86
		7.6.2	Continuous	87
	7.7	Statist		87
		7.7.1	Introduction	87
		7.7.2	Univariate statistics	87
		7.7.3	Examples of univariate statistics	87
		7.7.3 7.7.4	Examples of univariate statistics	87 88

		7.7.6	Variance	88
		7.7.7	Standard deviation	88
	7.8	Bivaria	ate statistics	89
		7.8.1	Covariance	89
		7.8.2	Correlation	89
	7.9	Regres	ssion	89
		7.9.1	Ordinary least squares	89
		7.9.2	Residuals	90
		7.9.3	Finding the right line	90
8		ulations		91
9	lext	analys	Sis	92
	V ra		SiS	92 93
	Wra	p up	ods at UT	_
	Wra	p up Metho		93
	Wra	p up Metho 10.1.1	ods at UT	93
	Wra	p up Metho 10.1.1 10.1.2	ods at UT	93 93 93
	Wra	p up Metho 10.1.1 10.1.2 10.1.3	ods at UT	93 93 93 93
	Wra	p up Metho 10.1.1 10.1.2 10.1.3 10.1.4	ods at UT	93 93 93 93 93

Class schedule

Date	Time	Location
Thurs, Aug. 10	9:00 AM - 4:00 PM	RLP 1.302E
Fri, Aug. 11	9:00 AM - 4:00 PM	RLP $1.302E$
Sat, Aug. 12	No class	-
Sun, Aug. 13	No class	-
Mon, Aug. 14	9:00 AM - 4:00 PM	RLP 1.302E
Tues, Aug. 15	9:00 AM - 4:00 PM	RLP 1.302E
Weds, Aug. 16	9:00 AM - 4:00 PM	RLP 1.302E

On class days, we will have a lunch break from 12:00-1:00 PM. We'll also take short breaks periodically during the morning and afternoon sessions as needed.

Description

Welcome to Introduction to Methods for Political Science, aka "Methods Camp"! In the past our incoming students have told us their math skills are rusty and they would like to be better prepared for UT's methods courses. Methods Camp is designed to give everyone a chance to brush up on some skills in preparation for the Stats I and Formal Theory I courses. The other goal of Methods Camp is to allow you to get to know your cohort. We hope that struggling with matrix algebra and the dreaded chain rule will still prove to be a good bonding exercise.

As you can see from the above schedule, we'll be meeting on Thursday, August 10th and Friday, August 11th as well as from Monday, August 14th through Wednesday, August 16th. Classes at UT begin the start of the following week on Monday, August 22nd. Below is a tentaive schedule outlining what will be covered in the class, although we may rearrange things a bit if we find we're going too slowly or too quickly through any of the material.

Course outline

1 Thursday morning: R and RStudio

- Introductions
- RStudio (materials are on the website as zipped RStudio projects)
- Objects (vectors, matrices, data frames)
- Basic functions (mean(), length(), etc.)

2 Thursday afternoon: tidyverse basics I

- Packages: installation and loading (including the tidyverse)
- Data wrangling with dplyr (basic verbs, including the new .by = syntax)
- Data visualization basics with ggplot2
- Data loading (.csv, .rds, .dta, .xlsx)
- Quarto fundamentals

3 Friday morning: Matrices

- Matrices
- Systems of linear equations
- Matrix operations (multiplication, transpose, inverse, determinant).
- Solving systems of linear equations in matrix form (and why that's cool)
- Introduction to OLS

4 Friday afternoon: tidyverse basics II

- Data merging and pivoting (*_join(), pivot_*())
- Value recoding (if_else(), case_when())
- Missing values
- Data visualization extensions: facets, text annotations

5 Monday morning: Functions and loops

- Functions
- For-loops and lapply()
- Finding R help (help files, effective Googling, ChatGPT)

6 Monday afternoon: Calculus

- Limits (not sure how to teach this in an R-centric way yet, but there must be a way)
- Derivatives (symbolic, numerical, automatic)
- Integrals

7 Tuesday morning: Probability

- Concepts: probability, random variables, etc.
- PMF, PDF, CDF, etc.
- Distributions (binomial, normal; different functions in R and how to use them)
- Expectation and variance

8 Tuesday afternoon: Simulations

- Simulations (ideas, seed setting, etc.)
- Sampling
- Bootstrapping

9 Wednesday morning: Text analysis

- String manipulation with stringr
- Simple text analysis (counts, tf-idf, etc.) with tidytext and visualization

10 Wednesday afternoon: Wrap-up

- Project management fundamentals (RStudio projects, keeping raw data, etc.)
- Self-study resources and materials
- Other software (Overleaf, Zotero, etc.)
- Methods at UT

Contact info

If you have any questions during or outside of methods camp, you can contact Andrés at andres.cruz@utexas.edu and Matt at mjmartin@utexas.edu.

Setup

Installing R and RStudio

R is a programming language optimized for statistics and data analysis. Most people use R from RStudio, a graphical user interface (GUI) that includes a file pane, a graphics pane, and other goodies. Both R and RStudio are open source, i.e., free as in beer and free as in freedom!

Your first steps should be to install R and RStudio, in that order (if you have installed these programs before, make sure that your versions are up-to-date—if they are not, follow the instructions below):

- 1. Download and install R from the official website, CRAN. Click on "Download R for <Windows/Mac>" and follow the instructions. If you have a Mac, make sure to select the version appropriate for your system (Apple Silicon for newer M1/M2 Macs and Intel for older Macs).
- 2. Download and install RStudio from the official website. Scroll down and select the installer for your operating system.

After these two steps, you can open RStudio in your system, as you would with any program. You should see something like this:

That's it for the installation! We also *strongly* recommend that you change a couple of RStudio's default settings.¹ You can change settings by clicking on Tools > Global Options in the menubar. Here are our recommendations:

- General > Uncheck "Restore .RData into workspace at startup"
- General > Save workspace to .RData on Exit > Select "Never"
- Code > Check "Use native pipe operator"
- Tools > Global Options > Appearance to change to a dark theme, if you want! Pros: better for night sessions, hacker vibes...

¹The idea behind these settings (or at least the first two) is to force R to start from scratch with each new session. No lingering objects from previous coding sessions avoids misunderstandings and helps with reproducibility!



Figure 1: How RStudio looks after a clean installation.

Setting up for Methods Camp

All materials for Methods Camp are both on this website and an RStudio project. An RStudio project is simply a folder where one keeps scripts, datasets, and other files needed for a data analysis project.

You can download our RStudio project here, as a .zip compressed file. On MacOS, the file will be uncompressed automatically. On Windows, you should do Right click > Extract all.



Make sure to properly unzip the materials. Double-clicking the .zip file on most Windows systems will not unzip the folder—you must do Right click > Extract all.

You should now have a folder called methodscamp/ on your computer. Navigate to the methodscamp.Rproj file within it and open it. RStudio should open the project right away. You should see methodscamp on the top-right of RStudio—this indicates that you are working in our RStudio project.



Figure 2: How the bottom-right corner of RStudio looks after opening our project.

That's all for setup! We can now start coding. After opening our RStudio project, we'll begin by opening the O1_r_intro.qmd file from the "Files" panel, in the bottom-right portion of RStudio. This is a Quarto document,² which contains both code and explanations (you can also read in the next chapter of this website).

²Perhaps you have used R Markdown before. Quarto is the next iteration of R Markdown, and is both more flexible and more powerful!

1 Intro to R

In Quarto documents like this one, we can write comments by just using plain text. In contrast, code needs to be within *code blocks*, like the one below. To execute a code block, you can click on the little "Play" button or press Cmd/Ctrl + Shift + Enter when your keyboard is hovering the code block.

```
2 + 2
```

[1] 4

That was our first R command, a simple math operation. Of course, we can also do more complex arithmetic:

```
12345 ^{\circ} 2 / (200 + 25 - 6 * 2) # this is an inline comment, see the leading "#"
```

[1] 715488.4

In order to create a code block, you can press Cmd/Ctrl + Alt + i or click on the little green "+C" icon on top of the script.

i Exercise

Create your own code block below and run a math operation.

1.1 Objects

A huge part of R is working with *objects*. Let's see how they work:

```
my_object <- 10 # opt/alt + minus sign will make the arrow
my_object # to print the value of an object, just call its name</pre>
```

```
[1] 10
```

We can now use this object in our operations:

```
2 ^ my_object
```

[1] 1024

Or even create another object out of it:

```
my_object2 <- my_object * 2
my_object2</pre>
```

[1] 20

You can delete objects with the rm() function (for "remove"):

```
rm(my_object2)
```

1.2 Vectors and functions

Objects can be of different types. One of the most useful ones is the vector, which holds a series of values. To create one manually, we can use the c() function (for "combine"):

```
my_vector <- c(6, -11, my_object, 0, 20)
my_vector</pre>
```

[1] 6 -11 10 0 20

We can use square brackets to retrieve parts of vectors:

```
my_vector[4] # fourth element
```

[1] 0

```
my_vector[1:2] # first two elements
[1]
      6 -11
Let's check out some basic functions we can use with numbers and numeric vectors:
  sqrt(my_object) # squared root
[1] 3.162278
  log(my_object) # logarithm (natural by default)
[1] 2.302585
  abs(-5) # absolute value
[1] 5
  mean(my_vector)
[1] 5
  median(my_vector)
[1] 6
  sd(my_vector) # standard deviation
[1] 11.53256
  sum(my_vector)
[1] 25
```

```
min(my_vector) # minimum value

[1] -11

max(my_vector) # maximum value

[1] 20

length(my_vector) # length (number of elements)
```

[1] 5

Notice that if we wanted to save any of these results for later, we would need to assign them:

```
my_mean <- mean(my_vector)
my_mean</pre>
```

[1] 5

Γ1] -11

These functions are quite simple: they take one object and do one operation. A lot of functions are a bit more complex—they take multiple objects or take options. For example, see the sort() function, which by default sorts a vector *increasingly*:

```
sort(my_vector)
```

If we instead want to sort our vector *decreasingly*, we can use the decreasing = TRUE argument (T also works as an abbreviation for TRUE).

```
sort(my_vector, decreasing = TRUE)
```

6 10

[1] 20 10 6 0 -11



If you use the argument values in order, you can avoid writing the argument names (see below). This is sometimes useful, but can also lead to confusing code—use it with caution.

```
sort(my_vector, T)
[1] 20 10 6 0 -11
```

To check the arguments of a function, you can examine its help file: look the function up on the "Help" panel on RStudio or use a command like the following: ?sort.

i Exercise

Examine the help file of the log() function. How can we compute the base-10 logarithm of my_object? Your code:

Other than numeric vectors, character vectors are also useful:

```
my_character_vector <- c("Apple", "Orange", "Watermelon", "Banana")

my_character_vector[3]

[1] "Watermelon"

nchar(my_character_vector) # count number of characters

[1] 5 6 10 6</pre>
```

1.3 Data frames and lists

Another useful object type is the *data frame*. Data frames can store multiple vectors in a tabular format. We can manually create one with the data.frame() function:

my_data_frame

```
fruit calories_per_100g water_per_100g
1 Apple 52 85.6
2 Orange 47 86.8
3 Watermelon 30 91.4
4 Banana 89 74.9
```

Now we have a little 4x3 data frame of fruits with their calorie counts and water composition. We gathered the nutritional information from the USDA (2019).

We can use the data_frame\$column construct to access the vectors within the data frame:

```
mean(my_data_frame$calories_per_100g)
```

[1] 54.5

i Exercise

Obtain the maximum value of water content per 100g in the data. Your code:

We will learn much more about data frames in our next module on data analysis.

After talking about vectors and data frames, the last object type that we will cover is the *list*. Lists are super flexible objects that can contain just about anything:

```
my_list <- list(my_object, my_vector, my_data_frame)</pre>
  my_list
[[1]]
[1] 10
[[2]]
[1]
      6 -11 10
                      20
[[3]]
       fruit calories_per_100g water_per_100g
       Apple
                              52
                                            85.6
1
2
      Orange
                              47
                                            86.8
```

```
3 Watermelon 30 91.4
4 Banana 89 74.9
```

To retrieve the elements of a list, we need to use double square brackets:

```
my_list[[1]]
```

[1] 10

Lists are sometimes useful due to their flexibility, but are much less common in routine data analysis compared to vectors or data frames.

1.4 Packages

The R community has developed thousands of *packages*, which are specialized collections of functions, datasets, and other resources. To install one, you should use the <code>install.packages()</code> command. Below we will install the <code>tidyverse</code> package, a suite for data analysis that we will use in the next modules. You just need to install packages once, and then they will be available system-wide.

```
install.packages("tidyverse") # this can take a couple of minutes
```

If you want to use an installed package in your script, you must load it with the library() function. Some packages, as shown below, will print descriptive messages once loaded.

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
v dplyr
           1.1.2
                      v readr
                                  2.1.4
v forcats
           1.0.0
                                  1.5.0
                     v stringr
v ggplot2
           3.4.2
                     v tibble
                                  3.2.1
v lubridate 1.9.2
                     v tidyr
                                  1.3.0
v purrr
            1.0.1
-- Conflicts ----- tidyverse conflicts() --
x dplyr::filter() masks stats::filter()
                 masks stats::lag()
x dplyr::lag()
i Use the conflicted package (<a href="http://conflicted.r-lib.org/">http://conflicted.r-lib.org/</a>) to force all conflicts to become
```

⚠ Warning

Remember that install.packages("package") needs to be executed just once, while library(package) needs to be in each script in which you plan to use the package. In general, never include install.packages("package") as part of your scripts or Quarto documents!

2 Tidy data analysis I

The tidyverse is a suite of packages that streamline data analysis in R. After installing the tidyverse with install.packages("tidyverse") (see the previous module), you can load it with:

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
v dplyr
           1.1.2
                     v readr
                                 2.1.4
                     v stringr
v forcats
           1.0.0
                                 1.5.0
v ggplot2 3.4.2
                     v tibble
                                 3.2.1
v lubridate 1.9.2
                     v tidyr
                                 1.3.0
           1.0.1
v purrr
-- Conflicts ----- tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()
                 masks stats::lag()
i Use the conflicted package (<a href="http://conflicted.r-lib.org/">http://conflicted.r-lib.org/</a>) to force all conflicts to become
```



Upon loading, the tidyverse prints a message like the one above. Notice that multiple packages (the constituent elements of the "suite") are actually loaded. For instance, dplyr and tidyr help with data transformation and wrangling, while ggplot2 allows us to draw plots. In most cases, one just loads the tidyverse and forgets about these details, as the constituent packages work together nicely.

Throughout this module, we will use tidyverse functions to load, wrangle, and visualize real data.

2.1 Loading data

Throughout this module we will work with a dataset of senators during the Trump presidency, which was adapted from FiveThirtyEight (2021).

We have stored the dataset in .csv format under the data/ subfolder. Loading it into R is simple (notice that we need to assign it to an object):

```
trump_scores <- read_csv("data/trump_scores_538.csv")</pre>
```

Rows: 122 Columns: 8

-- Column specification ------

Delimiter: ","

chr (4): bioguide, last_name, state, party

dbl (4): num_votes, agree, agree_pred, margin_trump

- i Use `spec()` to retrieve the full column specification for this data.
- i Specify the column types or set `show_col_types = FALSE` to quiet this message.

trump_scores

# 1	# A tibble: 122 x 8							
	bioguide	last_name	state	party	${\tt num_votes}$	agree	agree_pred	margin_trump
	<chr></chr>	<chr></chr>	<chr></chr>	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	A000360	Alexander	TN	R	118	0.890	0.856	26.0
2	B000575	Blunt	MO	R	128	0.906	0.787	18.6
3	B000944	Brown	OH	D	128	0.258	0.642	8.13
4	B001135	Burr	NC	R	121	0.893	0.560	3.66
5	B001230	Baldwin	WI	D	128	0.227	0.510	0.764
6	B001236	Boozman	AR	R	129	0.915	0.851	26.9
7	B001243	Blackburn	TN	R	131	0.885	0.889	26.0
8	B001261	Barrasso	WY	R	129	0.891	0.895	46.3
9	B001267	Bennet	CO	D	121	0.273	0.417	-4.91
10	B001277	${\tt Blumenthal}$	CT	D	128	0.203	0.294	-13.6
# :	# i 112 more rows							

Let's review the dataset's columns:

- bioguide: A unique ID for each politician, from the Congress Bioguide.
- last_name
- state
- party
- num_votes: Number of votes for which data was available.
- agree: Proportion (0-1) of votes in which the senator voted in agreement with Trump.
- agree_pred: Predicted proportion of vote agreement, calculated using Trump's margin (see next variable).

• margin_trump: Margin of victory (percentage points) of Trump in the senator's state.

We can inspect our data by using the interface above. An alternative is to run the command View(trump_scores) or click on the object in RStudio's environment panel (in the top-right section).

Do you have any questions about the data?

2.2 Wrangling data with dplyr

We often need to modify data to conduct our analyses, e.g., creating columns, filtering rows, etc. In the tidyverse, these operations are conducted with multiple *verbs*, which we will review now.

2.2.1 Selecting columns

We can select specific columns in our dataset with the select() function. All dplyr wrangling verbs take a data frame as their first argument—in this case, the columns we want to select are the other arguments.

```
select(trump_scores, last_name, party)
# A tibble: 122 x 2
   last_name
              party
   <chr>
              <chr>
1 Alexander
              R
2 Blunt
              R
3 Brown
              D
4 Burr
              R
5 Baldwin
              D
6 Boozman
              R
7 Blackburn
              R
8 Barrasso
              R.
9 Bennet
              D
10 Blumenthal D
# i 112 more rows
```

This is a good moment to talk about "pipes." Notice how the code below produces the same output as the one above, but with a slightly different syntax. Pipes (|>) "kick" the object on the left of the pipe to the first argument of the function on the right. One can read pipes

as "then," so the code below can be read as "take trump_scores, then select the columns last_name and party." Pipes are very useful to *chain multiple operations*, as we will see in a moment.

```
trump_scores |>
    select(last_name, party)
# A tibble: 122 x 2
   last_name party
   <chr>
              <chr>
 1 Alexander
2 Blunt
              R.
3 Brown
              D
4 Burr
              R
5 Baldwin
              D
6 Boozman
              R
7 Blackburn R
8 Barrasso
9 Bennet
              D
10 Blumenthal D
# i 112 more rows
```

? Tip

You can insert a pipe with the Cmd/Ctrl + Shift + M shortcut. If you have not changed the default RStudio settings, an "old" pipe (%>%) might appear. While most of the functionality is the same, the |> "new" pipes are more readable. You can change this RStudio option in Tools > Global Options > Code > Use native pipe operator. Make sure to check the other suggested settings in our Setup module!

Going back to selecting columns, you can select ranges:

```
trump_scores |>
    select(bioguide:party)
# A tibble: 122 x 4
  bioguide last_name state party
           <chr>
                      <chr> <chr>
  <chr>
1 A000360 Alexander
                     TN
                            R
2 B000575 Blunt
                      MO
                            R
3 B000944 Brown
                      OH
                            D
```

```
4 B001135 Burr
                      NC
                            R
5 B001230 Baldwin
                      WI
                            D
6 B001236 Boozman
                            R
                      AR
7 B001243 Blackburn TN
                            R
8 B001261 Barrasso
                      WY
                            R
9 B001267 Bennet
                      CO
                            D
10 B001277 Blumenthal CT
                            D
# i 112 more rows
```

And use a few helper functions, like matches():

```
trump_scores |>
   select(last_name, matches("agree"))
```

```
# A tibble: 122 x 3
```

	last_name	agree	agree_pred
	<chr></chr>	<dbl></dbl>	<dbl></dbl>
1	Alexander	0.890	0.856
2	Blunt	0.906	0.787
3	Brown	0.258	0.642
4	Burr	0.893	0.560
5	Baldwin	0.227	0.510
6	Boozman	0.915	0.851
7	Blackburn	0.885	0.889
8	Barrasso	0.891	0.895
9	Bennet	0.273	0.417
10	${\tt Blumenthal}$	0.203	0.294

Or everything(), which we usually use to reorder columns:

```
trump_scores |>
   select(last_name, everything())
```

A tibble: 122 x 8

i 112 more rows

	last_name	bioguide	state	party	${\tt num_votes}$	agree	agree_pred	${\tt margin_trump}$
	<chr></chr>	<chr></chr>	<chr></chr>	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	Alexander	A000360	TN	R	118	0.890	0.856	26.0
2	Blunt	B000575	MO	R	128	0.906	0.787	18.6
3	Brown	B000944	OH	D	128	0.258	0.642	8.13

4	Burr	B001135	NC	R	121	0.893	0.560	3.66
5	Baldwin	B001230	WI	D	128	0.227	0.510	0.764
6	Boozman	B001236	AR	R	129	0.915	0.851	26.9
7	Blackburn	B001243	TN	R	131	0.885	0.889	26.0
8	Barrasso	B001261	WY	R	129	0.891	0.895	46.3
9	Bennet	B001267	CO	D	121	0.273	0.417	-4.91
10	${\tt Blumenthal}$	B001277	CT	D	128	0.203	0.294	-13.6

? Tip

i 112 more rows

Notice that all these commands have not edited our existent objects—they have just printed the requested outputs to the screen. In order to modify objects, you need to use the assignment operator (<-). For example:

```
trump_scores_reduced <- trump_scores |>
   select(last_name, matches("agree"))

trump_scores_reduced
```

A tibble: 122 x 3

	last_name	agree	agree_pred
	<chr></chr>	<dbl></dbl>	<dbl></dbl>
1	Alexander	0.890	0.856
2	Blunt	0.906	0.787
3	Brown	0.258	0.642
4	Burr	0.893	0.560
5	Baldwin	0.227	0.510
6	Boozman	0.915	0.851
7	Blackburn	0.885	0.889
8	Barrasso	0.891	0.895
9	Bennet	0.273	0.417
10	Blumenthal	0.203	0.294
# :	i 112 more :	rows	

i Exercise

Select the variables $last_name$, party, num_votes , and agree from the data frame. Your code:

2.2.2 Renaming columns

We can use the rename() function to rename columns, with the syntax new_name = old_name. For example:

```
trump_scores |>
    rename(prop_agree = agree, prop_agree_pred = agree_pred)
# A tibble: 122 x 8
  bioguide last_name
                      state party num_votes prop_agree prop_agree_pred
           <chr>
  <chr>
                       <chr> <chr>
                                       <dbl>
                                                  <dbl>
                                                                   <dbl>
1 A000360 Alexander TN
                                                  0.890
                                                                   0.856
                             R
                                         118
2 B000575 Blunt
                                                                   0.787
                       MO
                             R
                                         128
                                                  0.906
3 B000944 Brown
                       OH
                                                                   0.642
                             D
                                         128
                                                  0.258
4 B001135 Burr
                       NC
                             R
                                         121
                                                  0.893
                                                                   0.560
5 B001230 Baldwin
                                         128
                                                  0.227
                                                                   0.510
                       WΙ
                             D
6 B001236 Boozman
                       AR
                             R
                                         129
                                                  0.915
                                                                   0.851
7 B001243 Blackburn
                       TN
                             R
                                         131
                                                  0.885
                                                                   0.889
8 B001261 Barrasso
                       WY
                             R
                                         129
                                                  0.891
                                                                   0.895
9 B001267 Bennet
                       CO
                             D
                                         121
                                                  0.273
                                                                   0.417
10 B001277 Blumenthal CT
                             D
                                         128
                                                  0.203
                                                                   0.294
# i 112 more rows
# i 1 more variable: margin_trump <dbl>
```

This is a good occasion to show how pipes allow us to chain operations. How do we read the following code out loud? (Remember that pipes are read as "then").

```
trump_scores |>
  select(last_name, matches("agree")) |>
  rename(prop_agree = agree, prop_agree_pred = agree_pred)
```

A tibble: 122 x 3

	last_name	<pre>prop_agree</pre>	<pre>prop_agree_pred</pre>
	<chr></chr>	<dbl></dbl>	<dbl></dbl>
1	Alexander	0.890	0.856
2	Blunt	0.906	0.787
3	Brown	0.258	0.642
4	Burr	0.893	0.560
5	Baldwin	0.227	0.510
6	Boozman	0.915	0.851
7	Blackburn	0.885	0.889

8 Barrasso	0.891	0.895
9 Bennet	0.273	0.417
10 Blumenthal	0.203	0.294
# i 112 more rows		

2.2.3 Creating columns

It is common to want to create columns, based on existing ones. We can use mutate() to do so. For example, we could want our main variables of interest in terms of percentages instead of proportions:

A tibble: 122 x 5

	last_name	agree	$agree_pred$	<pre>pct_agree</pre>	<pre>pct_agree_pred</pre>		
	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>		
1	Alexander	0.890	0.856	89.0	85.6		
2	Blunt	0.906	0.787	90.6	78.7		
3	Brown	0.258	0.642	25.8	64.2		
4	Burr	0.893	0.560	89.3	56.0		
5	Baldwin	0.227	0.510	22.7	51.0		
6	Boozman	0.915	0.851	91.5	85.1		
7	Blackburn	0.885	0.889	88.5	88.9		
8	Barrasso	0.891	0.895	89.1	89.5		
9	Bennet	0.273	0.417	27.3	41.7		
10	${\tt Blumenthal}$	0.203	0.294	20.3	29.4		
# :	# i 112 more rows						

We can also use multiple columns for creating a new one. For example, let's retrieve the total *number* of votes in which the senator agreed with Trump:

```
trump_scores |>
  select(last_name, num_votes, agree) |> # select just for clarity
  mutate(num_votes_agree = num_votes * agree)
```

```
# A tibble: 122 x 4
    last_name    num_votes agree num_votes_agree
```

<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1 Alexander	118	0.890	105
2 Blunt	128	0.906	116
3 Brown	128	0.258	33
4 Burr	121	0.893	108
5 Baldwin	128	0.227	29
6 Boozman	129	0.915	118
7 Blackburn	131	0.885	116
8 Barrasso	129	0.891	115
9 Bennet	121	0.273	33.0
10 Blumenthal	128	0.203	26
# i 112 more rows			

2.2.4 Filtering rows

Another common operation is to filter rows based on logical conditions. We can do so with the filter() function. For example, we can filter to only get Democrats:

```
trump_scores |>
  filter(party == "D")
```

# .	A tibble:	55 x 8						
	bioguide	last_name	state	party	${\tt num_votes}$	agree	agree_pred	margin_trump
	<chr></chr>	<chr></chr>	<chr></chr>	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	B000944	Brown	OH	D	128	0.258	0.642	8.13
2	B001230	Baldwin	WI	D	128	0.227	0.510	0.764
3	B001267	Bennet	CO	D	121	0.273	0.417	-4.91
4	B001277	${\tt Blumenthal}$	CT	D	128	0.203	0.294	-13.6
5	B001288	Booker	NJ	D	119	0.160	0.290	-14.1
6	C000127	Cantwell	WA	D	128	0.242	0.276	-15.5
7	C000141	Cardin	MD	D	128	0.25	0.209	-26.4
8	C000174	Carper	DE	D	129	0.295	0.318	-11.4
9	C001070	Casey	PA	D	129	0.287	0.508	0.724
10	C001088	Coons	DE	D	128	0.289	0.319	-11.4
# :	i 45 more	rows						

Notice that == here is a *logical operator*, read as "is equal to." So our full chain of operations says the following: take trump_scores, then filter it to get rows where party is equal to "D".

There are other logical operators:

Logical operator	Meaning
==	"is equal to"
!=	"is not equal to"
>	"is greater than"
<	"is less than"
>=	"is greater than or equal to"
<=	"is less than or equal to"
%in%	"is contained in"
&	"and" (intersection)
1	"or" (union)

Let's see a couple of other examples.

```
trump_scores |>
  filter(agree > 0.5)
```

A tibble: 69 x 8

```
bioguide last_name state party num_votes agree agree_pred margin_trump
  <chr>
            <chr>
                      <chr> <chr>
                                       <dbl> <dbl>
                                                        <dbl>
                                                                      <dbl>
1 A000360 Alexander TN
                                         118 0.890
                                                        0.856
                                                                      26.0
                            R
2 B000575 Blunt
                      MO
                                         128 0.906
                                                                      18.6
                                                        0.787
                      NC
3 B001135
            Burr
                            R
                                         121 0.893
                                                        0.560
                                                                      3.66
4 B001236 Boozman
                      AR.
                            R
                                         129 0.915
                                                        0.851
                                                                      26.9
5 B001243 Blackburn TN
                                         131 0.885
                                                        0.889
                                                                      26.0
                            R
6 B001261 Barrasso
                      WY
                            R
                                         129 0.891
                                                        0.895
                                                                      46.3
7 B001310 Braun
                      IN
                            R
                                         44 0.909
                                                        0.713
                                                                      19.2
                                                                      17.8
8 C000567
            Cochran
                      MS
                                          68 0.971
                                                        0.830
                            R
9 C000880
            Crapo
                      ID
                            R
                                         125 0.904
                                                        0.870
                                                                     31.8
10 C001035 Collins
                      ME
                                         129 0.651
                                                        0.441
                                                                      -2.96
```

i 59 more rows

```
trump_scores |>
  filter(state %in% c("CA", "TX"))
```

A tibble: 4 x 8

	ртовитае	rast_name	state	party	num_votes	agree	agree_pred	margin_trump
	<chr></chr>	<chr></chr>	<chr>></chr>	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	C001056	Cornyn	TX	R	129	0.922	0.659	9.00
2	C001098	Cruz	TX	R	126	0.921	0.663	9.00

```
3 F000062 Feinstein CA D 128 0.242 0.201 -30.1
4 H001075 Harris CA D 116 0.164 0.209 -30.1
```

```
trump_scores |>
  filter(state == "WV" & party == "D")
```

A tibble: 1 x 8

i Exercise

- 1. Add a new column to the data frame, called diff_agree, which subtracts agree and agree_pred. How would you create abs_diff_agree, defined as the absolute value of diff_agree?
- 2. Filter the data frame to only get senators for which we have information on fewer than (or equal to) five votes. Your code:
- 3. Filter the data frame to only get Democrats who agreed with Trump in at least 30% of votes. Your code:

2.2.5 Ordering rows

The arrange() function allows us to order rows according to values. For example, let's order based on the agree variable:

```
trump_scores |>
  arrange(agree)
```

A tibble: 122 x 8

	bioguide	last_name	state	party	num_votes	agree	agree_pred	margin_trump
	<chr></chr>	<chr></chr>	<chr></chr>	<chr></chr>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	H000273	Hickenlooper	CO	D	2	0	0.0302	-4.91
2	H000601	Hagerty	TN	R	2	0	0.115	26.0
3	L000570	Luján	NM	D	186	0.124	0.243	-8.21
4	G000555	Gillibrand	NY	D	121	0.124	0.242	-22.5
5	M001176	Merkley	OR	D	129	0.155	0.323	-11.0
6	W000817	Warren	MA	D	116	0.155	0.216	-27.2

7 B001288	Booker	NJ	D	119 0.160	0.290	-14.1
8 S000033	Sanders	VT	D	112 0.161	0.221	-26.4
9 H001075	Harris	CA	D	116 0.164	0.209	-30.1
10 M000133	Markey	MA	D	127 0.165	0.213	-27.2
# i 112 more rows						

Maybe we only want senators with more than a few data points. Remember that we can chain operations:

```
trump_scores |>
  filter(num_votes >= 10) |>
  arrange(agree)
```

```
# A tibble: 115 x 8
   bioguide last_name
                        state party num_votes agree agree_pred margin_trump
   <chr>>
            <chr>
                        <chr> <chr>
                                         <dbl> <dbl>
                                                           <dbl>
                                                                         <dbl>
 1 L000570
            Luján
                        NM
                              D
                                           186 0.124
                                                           0.243
                                                                         -8.21
2 G000555
            Gillibrand NY
                              D
                                           121 0.124
                                                           0.242
                                                                        -22.5
3 M001176
            Merkley
                        OR
                              D
                                           129 0.155
                                                           0.323
                                                                        -11.0
4 W000817
            Warren
                              D
                                           116 0.155
                                                           0.216
                                                                        -27.2
                        MA
5 B001288 Booker
                        NJ
                              D
                                           119 0.160
                                                           0.290
                                                                        -14.1
6 S000033
           Sanders
                        VT
                              D
                                           112 0.161
                                                           0.221
                                                                        -26.4
7 H001075 Harris
                        CA
                              D
                                           116 0.164
                                                           0.209
                                                                        -30.1
8 M000133 Markey
                        MA
                              D
                                           127 0.165
                                                           0.213
                                                                        -27.2
9 W000779
            Wyden
                        \mathsf{OR}
                              D
                                           129 0.186
                                                           0.323
                                                                        -11.0
            Blumenthal CT
10 B001277
                              D
                                           128 0.203
                                                           0.294
                                                                        -13.6
# i 105 more rows
```

By default, arrange() uses increasing order (like sort()). To use decreasing order, add a minus sign:

```
trump_scores |>
  filter(num_votes >= 10) |>
  arrange(-agree)
```

A tibble: 115 x 8

bioguide last_name state party num_votes agree agree_pred margin_trump <chr> <chr> <chr> <chr> <dbl> <dbl> <dbl> <dbl> 1 M001198 Marshall KS R 183 0.973 0.933 20.6 2 C000567 Cochran MS R 68 0.971 0.830 17.8

3 H000338	Hatch	UT	R	84	0.964	0.825	18.1
4 M001197	McSally	ΑZ	R	136	0.949	0.562	3.55
5 P000612	Perdue	GA	R	119	0.941	0.606	5.16
6 C001096	Cramer	ND	R	135	0.941	0.908	35.7
7 R000307	Roberts	KS	R	127	0.937	0.818	20.6
8 C001056	Cornyn	TX	R	129	0.922	0.659	9.00
9 H001061	Hoeven	ND	R	129	0.922	0.883	35.7
10 C001047	Capito	WV	R	127	0.921	0.896	42.2
# i 105 mor	e rows						

You can also order rows by more than one variable. What this does is to order by the first variable, and resolve any ties by ordering by the second variable (and so forth if you have more than two ordering variables). For example, let's first order our data frame by party, and then within party order by agreement with Trump:

```
trump_scores |>
  filter(num_votes >= 10) |>
  arrange(party, agree)
```

A tibble: 115 x 8

	${\tt bioguide}$	last_name	state	party	num	_votes	agree	agree_pred	margin_trump
	<chr></chr>	<chr></chr>	<chr></chr>	<chr></chr>		<dbl></dbl>	<dbl></dbl>	<dbl></dbl>	<dbl></dbl>
1	L000570	Luján	NM	D		186	0.124	0.243	-8.21
2	G000555	${\tt Gillibrand}$	NY	D		121	0.124	0.242	-22.5
3	M001176	Merkley	OR	D		129	0.155	0.323	-11.0
4	W000817	Warren	MA	D		116	0.155	0.216	-27.2
5	B001288	Booker	NJ	D		119	0.160	0.290	-14.1
6	S000033	Sanders	VT	D		112	0.161	0.221	-26.4
7	H001075	Harris	CA	D		116	0.164	0.209	-30.1
8	M000133	Markey	MA	D		127	0.165	0.213	-27.2
9	W000779	Wyden	OR	D		129	0.186	0.323	-11.0
10	B001277	${\tt Blumenthal}$	CT	D		128	0.203	0.294	-13.6

[#] i 105 more rows

i Exercise

Arrange the data by diff_pred, the difference between agreement and predicted agreement with Trump. (You should have code on how to create this variable from the last exercise).

2.2.6 Summarizing data

dplyr makes summarizing data a breeze using the summarize() function:

To make summaries, we can use any function that takes a vector and returns one value. Another example:

Grouped summaries allow us to disaggregate summaries according to other variables (usually categorical):

```
trump_scores |>
    filter(num_votes >= 5) |> # to filter out senators with few data points
    summarize(mean_agree = mean(agree),
             max_agree = max(agree),
             min agree = min(agree),
              .by = party) # to group by party
# A tibble: 2 x 4
 party mean_agree max_agree min_agree
 <chr>
            <dbl>
                     <dbl>
                                <dbl>
            0.876
1 R
                                0.651
                      1
            0.272 0.548
2 D
                                0.124
```

i Exercise

Obtain the maximum absolute difference in agreement with Trump (the abs_diff_agree variable from before) for each party.

2.2.7 Overview

Function	Purpose
select()	Select columns
rename()	Rename columns
<pre>mutate()</pre>	Creating columns
filter()	Filtering rows
arrange()	Ordering rows
<pre>summarize()</pre>	Summarizing data
<pre>summarize(, .by =)</pre>	Summarizing data (by group)

2.3 Visualizing data with ggplot2

- ggplot2 is an R package dedicated to data visualization.
- We can build charts through layers using geom_ functions. There are more than 20 options available such as the following:


```
- geom_bar (bar plots)
- geom_boxplot (box plots)
- geom_hist (histograms)
```

• The ggtitle() function allows to add a title to the chart.

(TODO)


```
trump_scores_ss <- trump_scores |>
  filter(num_votes >= 10)

ggplot(trump_scores_ss, aes(x = party)) +
  geom_bar()
```




```
ggplot(trump_scores_ss, aes(x = agree)) +
  geom_histogram()
```

`stat_bin()` using `bins = 30`. Pick better value with `binwidth`.




```
ggplot(trump_scores_ss, aes(x = agree, fill = party)) +
   geom_histogram()
```


`stat_bin()` using `bins = 30`. Pick better value with `binwidth`.




ggplot(trump_scores_ss, aes(x = margin_trump, y = agree)) +
 geom_point()



```
ggplot(trump_scores_ss, aes(x = margin_trump, y = agree, color = party)) +
  geom_point()
```



Relationship between Trump margins and senators' votes



3 Matrices

3.1 Introduction

3.1.1 Scalars

- One number (for example, 12) is referred to as a scalar.
- Each scalar in a matrix is an *element* of that matrix.

[12]

Note

This is also called a 1×1 ("one by one") matrix.

3.1.2 Vectors

- We can put several scalars together to make a vector.
- Here is an example:

$$\begin{bmatrix} 12\\14\\15 \end{bmatrix} = b$$

- Since this is a column of numbers, we cleverly refer to it as a column vector.
- Here is another example of a vector:

$$\begin{bmatrix} 12 & 14 & 15 \end{bmatrix} = d$$

• This, in contrast, is called a row vector.

3.1.2.1 Vector construction in RStudio

- Vectors are very easy to construct in RStudio.
- Here are two examples:

```
# Making a simple column vector
col_vector <- matrix(1:12)
col_vector</pre>
```

```
[,1]
 [1,]
 [2,]
         2
 [3,]
         3
 [4,]
         4
 [5,]
        5
 [6,]
         6
 [7,]
         7
 [8,]
        8
 [9,]
        9
[10,]
        10
[11,]
        11
[12,]
        12
```

```
# Making a simple row vector
row_vector <- 15:23
row_vector</pre>
```

[1] 15 16 17 18 19 20 21 22 23

3.2 Operators

3.2.1 Summation

• The summation operator \sum lets us perform an operation on a sequence of numbers, which is often but not always a vector.

$$x = \begin{bmatrix} 12 & 7 & -2 & 0 & 1 \end{bmatrix}$$

• We can then calculate the sum of the first three elements of the vector, which is expressed as follows:

$$\sum_{i=1}^{3} x_i$$

• Then, we do the following math:

$$12 + 7 + (-2) = 17$$

3.2.2 Product

• The product operator \prod can also perform operations over a sequence of elements in a vector.

$$z = \begin{bmatrix} 5 & -3 & 5 & 1 \end{bmatrix}$$

• We can then calculate the calculate the product of the four elements in the vector, which is expressed as follows:

$$\prod_{i=1}^4 z_i$$

• Then, we do the following math:

$$5\times -3\times 5\times 1=-75$$

Matrices

3.2.2.1 Operators in RStudio

```
# Summation
mat <- c(5,-3,5,1)
sum(mat)
```

[1] 8

```
# Product
prod(mat)
```

[1] -75

3.2.3 Basics

• We can append vectors together to form a matrix:

$$\begin{bmatrix} 12 & 14 & 15 \\ 115 & 22 & 127 \\ 193 & 29 & 219 \end{bmatrix} = A$$

- The number of rows and columns of a matrix constitute the dimensions of the matrix.
- The first number is the number of rows ("r") and the second number is the number of columns ("c") in the matrix.

Important

Find a way to remember "r x c" permanently. The order of the dimensions never changes.

- Matrix A above, for example, is a 3x3 matrix.
- We often use capital letters (sometimes **bold-faced**) to represent matrices.

3.2.3.1 Appending in RStudio

- We use rbind() to create a matrix, in which each vector will be a row.
- If we use cbind(), then each vector will be a column.

```
# Create some vectors
  vec_1 <- 1:4
  vec_2 <- 5:8
  vec_3 <- 9:12
  vec_4 <- 13:16
  # rbind
  rbind_mat <- rbind(vec_1, vec_2, vec_3, vec_4)</pre>
  rbind_mat
      [,1] [,2] [,3] [,4]
vec_1
         1
                    7
vec_2
         5
              6
                         8
         9
             10
vec_3
                   11
                        12
vec_4 13
             14
                        16
```

```
# cbind
  cbind_mat <- cbind(vec_1, vec_2, vec_3, vec_4)</pre>
  cbind_mat
     vec_1 vec_2 vec_3 vec_4
[1,]
         1
[2,]
         2
                      10
         3
                7
[3,]
                      11
                             15
[4,]
                             16
```

3.2.4 Structure

- How do we refer to specific elements of the matrix?
- Matrix A is an $m \times n$ matrix where m = n = 3
- More generally, matrix B is an $m \times n$ matrix where the elements look like this:

$$B = \begin{bmatrix} b_{11} & b_{12} & b_{13} & \dots & b_{1n} \\ b_{21} & b_{22} & b_{23} & \dots & b_{2n} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ b_{m1} & b_{m2} & b_{m3} & \dots & b_{mn} \end{bmatrix}$$

• Thus b_{23} refers to the second unit down and third across.



When trying to identify a specific element, the first subscript is the element's row and the second subscript is the element's column (always in that order).

3.2.4.1 Constructing matrices in RStudio

- Matrices are very easy to construct in RStudio.
- The basic syntax is matrix(data, nrow, ncol, byrow, dimnames).
 - data is the input vector which becomes the data elements of the matrix.
 - nrow is the number of rows to be created.
 - ncol is the number of columns to be created.
 - byrow is a logical clue. If TRUE then the input vector elements are arranged by row.
 - dimname is the names assigned to the rows and columns.
- Here are some examples:

```
# Elements are arranged sequentially by row.
  M \leftarrow matrix(c(1:12), nrow = 4, byrow = TRUE)
  print(M)
     [,1] [,2] [,3]
[1,]
       1
             2
[2,]
       4
             5
                  6
[3,]
       7
             8
                 9
[4,]
     10
           11
                 12
  # Elements are arranged sequentially by column.
  N \leftarrow matrix(c(1:12), nrow = 4, byrow = FALSE)
  print(N)
     [,1] [,2] [,3]
[1,]
            5
[2,]
        2
             6
                 10
[3,]
     3
             7
               11
[4,]
             8
                 12
  # Define the column and row names.
  rownames = c("row1", "row2", "row3", "row4")
  colnames = c("col1", "col2", "col3")
  P <- matrix(c(1:12), nrow = 4, byrow = TRUE, dimnames = list(rownames, colnames))
  print(P)
     col1 col2 col3
             2
row1
row2
             5
                  6
row3
        7
             8
                 9
       10
                 12
row4
            11
  # Access the element at 1st row and 3rd column.
  print(P[1,3])
[1] 3
```

```
# Access the element at 4th row and 2nd column.
print(P[4,2])
```

[1] 11

3.3 Matrix operations

3.3.1 Addition and subtraction

- Addition and subtraction are straightforward operations.
- Matrices must have *exactly* the same dimensions for both of these operations.
- We add or subtract each element with the corresponding element from the other matrix.
- This is expressed as follows:

$$A \pm B = C$$

$$c_{ij} = a_{ij} \pm b_{ij} \ \forall i, j$$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \pm \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix} = \begin{bmatrix} a_{11} \pm b_{11} & a_{12} \pm b_{12} & a_{13} \pm b_{13} \\ a_{21} \pm b_{21} & a_{22} \pm b_{22} & a_{23} \pm b_{23} \\ a_{31} \pm b_{31} & a_{32} \pm b_{32} & a_{33} \pm b_{33} \end{bmatrix}$$

3.3.1.1 Add and subtract in RStudio

```
matrix2 \leftarrow matrix(c(5, 2, 0, 9, 3, 4), nrow = 2)
  print(matrix2)
   [,1] [,2] [,3]
[1,] 5 0
[2,] 2
          9 4
  # Add the matrices.
  result1 <- matrix1 + matrix2</pre>
  cat("Result of addition","\n")
Result of addition
  print(result1)
   [,1] [,2] [,3]
[1,] 8 -1 5
[2,]
           13
      11
                10
  # Subtract the matrices
  result2 <- matrix1 - matrix2</pre>
  cat("Result of subtraction","\n")
Result of subtraction
  print(result2)
   [,1] [,2] [,3]
[1,] -2 -1 -1
[2,] 7 -5 2
```

Practice

$$A = \begin{bmatrix} 1 & 4 & 2 \\ -2 & -1 & 0 \\ 0 & -1 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 5 & 1 & 0 \\ 2 & -1 & 0 \\ 7 & 1 & 2 \end{bmatrix}$$

Calculate A + B

Practice

$$A = \begin{bmatrix} 6 & -2 & 8 & 12 \\ 4 & 42 & 8 & -6 \\ -14 & 5 & 0 & 0 \end{bmatrix}$$

$$B = \begin{bmatrix} 18 & 42 & 3 & 7 \\ 0 & -42 & 15 & 4 \\ -7 & 0 & 21 & -18 \end{bmatrix}$$

Calculate A - B

3.3.2 Scalar multiplication

- Scalar multiplication is very intuitive.
- As we know, a scalar is a single number, or a 1 x 1 matrix.
- We multiply each value in the matrix by the scalar to perform this operation.
- This is expressed as follows:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$cA = \begin{bmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \\ ca_{31} & ca_{32} & ca_{33} \end{bmatrix}$$

3.3.2.1 Scalar multiplication in RStudio

• All we need to do is take an established matrix and multiply it by some scalar.

matrix1

matrix1*3

Practice

$$A = \begin{bmatrix} 1 & 4 & 2 \\ 8 & -1 & 3 \\ 0 & -2 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} -15 & 1 & 5 \\ 2 & -42 & 0 \\ 7 & 1 & 6 \end{bmatrix}$$

Calculate $2 \times A$ and $-3 \times B$

3.3.3 Matrix multiplication

- Two matrices must be *conformable* for them to be multiplied together.
- This means that the number of columns in the first matrix equals the number of rows in the second.
- When multiplying $A \times B$, if A is $m \times n$, B must have n rows.

Important

The conformability requirement never changes. Before multiplying anything, check to make sure the matrices are indeed conformable.

- The resulting matrix will have the same number of rows as the first matrix and the number of columns in the second.
- For example, if A is $i \times k$ and B is $k \times j$, then $A \times B$ will be $i \times j$.

Which of the following can we multiply? What will be the dimensions of the resulting matrix?

$$B = \begin{bmatrix} 2 \\ 3 \\ 4 \\ 1 \end{bmatrix} M = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 4 \\ 2 & 3 & 2 \end{bmatrix} L = \begin{bmatrix} 6 & 5 & -1 \\ 1 & 4 & 3 \end{bmatrix}$$

Warning

When multiplying matrices, order matters.

• Why can't we multiply in the opposite order?

3.3.4 Multiplication steps

• Multiply each row by each column, summing up each pair of multiplied terms.

Note

This is sometimes to referred to as the "dot product," where we multiply matching members, then sum up.

• The element in position ij is the sum of the products of elements in the ith row of the first matrix (A) and the corresponding elements in the jth column of the second matrix (B).

$$c_{ij} = \sum_{k=1}^{n} a_{ik} b_{kj}$$

3.3.4.1 Example

- Suppose a company manufactures two kinds of furniture: chairs and sofas.
 - A chair costs \$100 for wood, \$270 for cloth, and \$130 for feathers.
 - Each sofa costs \$150 for wood, \$420 for cloth, and \$195 for feathers.

	Chair	Sofa
Wood	100	150
Cloth	270	420
Feathers	130	195

• The same information about unit cost (C) can be presented as a matrix.

$$C = \begin{bmatrix} 100 & 150 \\ 270 & 420 \\ 130 & 195 \end{bmatrix}$$

i Note

Note that each of the three rows of this 3×2 matrix represents a material (wood, cloth, or feathers), and each of the two columns represents a product (chair or coach). The elements are the unit cost (in USD).

- Now, suppose that the company will produce 45 chairs and 30 sofas this month.
- This production quantity can be represented in the following table, and also as a 2 x 1 matrix (Q).

Product	Quantity
Chair	45
Sofa	30

$$Q = \begin{bmatrix} 45 \\ 30 \end{bmatrix}$$

- The "total expenditure" is equal to the "unit cost" times the "production quantity" (the number of units).
- The total expenditure (E) for each material this month is calculated by multiplying these two matrices.

$$E = CQ = \begin{bmatrix} 100 & 150 \\ 270 & 420 \\ 130 & 195 \end{bmatrix} \begin{bmatrix} 45 \\ 30 \end{bmatrix} = \begin{bmatrix} (100)(45) + (150)(30) \\ (270)(45) + (420)(30) \\ (130)(45) + (195)(30) \end{bmatrix} = \begin{bmatrix} 9,000 \\ 24,750 \\ 11,700 \end{bmatrix}$$

- Multiplying the 3 x 2 Cost matrix (C) times the 2 x 1 Quantity matrix (Q) yields the 3 x 1 Expenditure matrix (E).
- As a result of this matrix multiplication, we determine that this month the company will incur expenditures of:
 - \$9,000 for wood
 - \$24,750 for cloth
 - \$11,700 for feathers.

3.3.4.2 Matrix multiplication in RStudio

- We must make sure the matrices are conformable, as we do for our manual calculations.
- Then, we can multiply them together without issue.
- When multiplying together matrices in RStudio, remember to use the **% operator, otherwise you will receive an error message.



If you have a missing value or NA in one of the matrices you are trying to multiply, you will have NAs in your resulting matrix.

```
# Creating matrices
m <- matrix(1:8, nrow=2) #
n <- matrix(8:15, nrow=4)
m</pre>
```

n

Multiplying matrices using operator $\mathtt{m}\%*\%\mathtt{n}$

3.4 Properties

- Addition and subtraction:
 - Associative: $(A \pm B) \pm C = A \pm (B \pm C)$
 - Communicative: $A \pm B = B \pm A$
- Multiplication:

$$-AB \neq BA$$

$$-A(BC) = (AB)C$$

$$-A(B+C) = AB + AC$$

$$- (A+B)C = AC + BC$$

3.5 Special matrices

Square matrix:

• The **diagonal** of a square matrix is a set of numbers consisting of the elements on the line from the upper-left-hand to the lower-right-hand corner of the matrix. Only a square matrix has a diagonal.

• The **trace** of a matrix is simply the sum of the diagonal elements of the matrix. So, then, a matrix must be square to have a trace.

Diagonal matrix:

• In a **diagonal matrix**, all of the elements of the matrix that are not on the diagonal are equal to zero.

Scalar matrix:

• A scalar matrix is a diagonal matrix where the diagonal elements are all equal to each other. In other words, we're really only concerned with one scalar (or element) held in the diagonal.

Identity matrix:

- The **identity matrix** is a scalar matrix with all of the diagonal elements equal to one.
- All of the off-diagonal elements are equal to zero.
- The capital letter I is reserved for the identity matrix.

3.6 Transpose

- The transpose is the original matrix with the rows and the columns interchanged.
- The notation is either J' ("J prime") or J^T ("J transpose").

$$J = \begin{bmatrix} 4 & 5 \\ 3 & 0 \\ 7 & -2 \end{bmatrix}$$

$$J' = J^T = \begin{bmatrix} 4 & 3 & 7 \\ 5 & 0 & -2 \end{bmatrix}$$

3.6.0.1 Transpose in RStudio

• We use t() to get the transpose.

```
# Construct our matrix
  J \leftarrow matrix(c(4,5,3,0,7,-2), nrow = 3, byrow = TRUE,)
     [,1] [,2]
[1,]
        4
[2,]
        3
              0
[3,]
        7
             -2
  t(J)
     [,1] [,2] [,3]
[1,]
[2,]
        5
```

3.7 Inverse

- Just like a number has a reciprocal, a matrix has an inverse.
- When we multiply a matrix by its inverse we get the identity matrix (which is like "1" for matrices).

$$A \times A^{-1} = I$$

• The inverse of A is A-1 only when:

$$AA^{-1} = A^{-1}A = I$$

• Sometimes there is no inverse at all.

Note

For now, don't worry about calculating the inverse of a matrix manually. This is the type of task we use RStudio for.

3.7.0.1 Examples

• We use the solve() function to calculate the inverse of a matrix.

```
# Create 3 different vectors
  # using combine method.
  a1 <- c(3, 2, 5)
  a2 < c(2, 3, 2)
  a3 <- c(5, 2, 4)
  # bind the three vectors into a matrix
  # using rbind() which is basically
  # row-wise binding.
  A <- rbind(a1, a2, a3)
  # print the original matrix
  print(A)
  [,1] [,2] [,3]
a1
a2
     2
          3
              2
a3
     5
          2
  # Use the solve() function to calculate the inverse.
  T1 <- solve(A)
  # print the inverse of the matrix
  print(T1)
             a1
                        a2
                                  a3
[1,] -0.29629630 -0.07407407 0.4074074
[3,] 0.40740741 -0.14814815 -0.1851852
```

4 Tidy data analysis II

5 Functions and loops

5.1 Basics

5.1.1 What is a function?

- Anything that takes input(s) and gives one defined output.
- They assign a unique value in its range (y values) for each value in its domain (x values).
- In math, this usually looks something like f(x) = 3x + 4.
 - -x is the argument that the function takes.
 - For any x, multiply x by 3 and then add 4
 - Alternative but equivalent notation: y = 3x + 4
 - -y is "a function of" x, so y = f(x)
- We describe functions with both equations and graphs.

5.1.2 Function machine




Figure 5.1: Function machine

5.1.3 Visualization

When graphed, we can't draw vertical line through a function. Why not?

Vertical Line Test - Functions



5.2 Types of functions

5.2.1 Linear functions

• We can easily make a function that describes a line.

$$y = mx + b$$


- m is the slope (for every one unit increase in x, y increases m units).
 - b is the y-intercept: the value of y when x = 0.
 - More generally, y = a + bx a is the intercept and b is the slope.

5.2.2 Quadratic

• These lines have one curve.

$$y = ax^2 + bx + c$$

- a, b, and c don't have well-defined meanings here.
- If a is negative, the function opens downward; if a is positive, it opens upward.
- Note that x^2 always returns positive values.



5.2.3 Cubic

- $\bullet\,$ These lines (generally) have two curves (inflection points).
- $y = ax^3 + bx^2 + cx + d$
- a, b, c, and d don't have well-defined meanings here.



5.2.4 Polynomial

$$y = ax^n + bx^{n-1} + \dots + c$$

- These functions have (maximum) n-1 changes in direction (turning points). - They also have (maximum) n x-intercepts. - They can be made arbitrarily precise.

5.2.5 Exponential

$$y = ab^x$$

or

$$f(x) = ab^x$$

• Here our independent variable, or input (x), is the exponent.

5.2.6 Trigonometric functions

- These functions include sine, cosine, and tangent.
- They are interesting (to some), but not usually useful for social science.

5.3 Logarithms and exponents

5.3.1 Logarithms

- Logarithms are basically the opposite (inverse) of exponents.
- They ask how many times you must raise the base to get x.
- $log_a(b) = x$ is asking "a raised to what power x gives b?
- $\log_3(81) = 4$ because $3^4 = 81$
- Logarithms can be undefined.
- The base cannot be 0, 1, or negative.

5.3.2 Relationships

If,

$$log_a x = b$$

then,

$$a^{log_a x} = a^b$$

and

$$x = a^b$$

5.3.3 Basic rules

$$\frac{\log_x n}{\log_x m} = \log_m n$$

$$\log_x(ab) = \log_x a + \log_x b$$

$$\log_x \left(\frac{a}{b}\right) = \log_x a - \log_x b$$

$$\log_x a^b = b \log_x a$$

$$\log_x 1 = 0$$

$$log_x x = 1$$

$$m^{\log_m(a)} = a$$

5.3.4 Natural logarithms

- We most often use natural logarithms.
- This means $\log_e(x)$, often written $\ln(x)$.
- $e \approx 2.7183$.
- $\ln(x)$ and its exponent opposite, e^x , have nice properties when we hit calculus.

5.3.5 Definition of e

• Imagine you invest \$1 in a bank and receive 100% interest for one year, and the bank pays you back once a year:

$$(1+1)^1 = 2$$

• When it pays you twice a year with compound interest:

$$(1+1/2)^2 = 2.25$$

• If it pays you three times a year:

$$(1+1/3)^3 = 2.37...$$

• What will happen when the bank pays you once a month? Once a day?

$$(1+\frac{1}{n})^n$$

• However, there is limit to what you can get

$$\lim_{n\to\infty}(1+\frac{1}{n})^n=2.7183...=e$$

• For any interest rate k and number of times the bank pays you t:

$$\lim_{n\to\infty}(1+\frac{k}{n})^{nt}=e^{kt}$$

• e is important for defining exponential growth. Since $ln(e^x) = x$, the natural logarithm helps us turn exponential functions into linear ones.

Practice

Solve the problems below, simplifying as much as you can.

$$log_{10}(1000)$$

$$log_{2}(\frac{8}{32})$$

$$10^{log_{10}(300)}$$

$$ln(1)$$

$$ln(e^{2})$$

$$ln(5e)$$

5.4 Functions of functions

5.4.1 Basics

- Functions can take other functions as arguments.
- This means that outside function takes output of inside function as its input.
- This is typically written as f(g(x)).
- Say we have the exterior function $f(x)=x^2$ and the interior function g(x)=x-3.
- Then if we want f(g(x)), we would subtract 3 from any input, and then square the result.
- We write this $(x-3)^2$, NOT x^2-3 .

5.4.2 PMF, PDF, and CDF

- PMF probability mass function
 - This gives the probability that a discrete random variable is exactly equal to some value.
- PDF probability density function
 - This gives the probability that a continuous random variable falls within a particular range of values.
- CDF cumulative distribution function
 - This gives the probability that a random variable X takes a value less than or equal to x.

6 Calculus

6.1 Theory

- Calculus is about dealing with infinitesimal values.
- We are going to focus on two big ideas:
 - Derivatives
 - Integrals

6.1.1 Derviative

- \bullet "Derivative" is just a fancy term for slope.
- Slope is the rate of change $\frac{\delta y}{\delta x}$ or $\frac{dy}{dx}$.
- Specifically, the derivative is the *instantaneous* rate of change.
- We need slope for our statistics, which are all about fitting lines.
- We also need slope for taking maxima and minima.
- The equation for a line is y = mx + b. What is its slope?

6.1.2 Calculating derivatives

- Slope is rise over run, which is $\frac{f(x+\Delta x)-f(x)}{\Delta x}$ To see why, consider the slope of a line connecting two points:

$$m=\frac{f(x_2)-f(x_1)}{x_2-x_1}$$

- We can define $x_2 = x_1 + \Delta x$ (or equivalently $\Delta x = x_2 - x_1)$

$$m = \frac{f(x_1 + \Delta x) - f(x_1)}{\Delta x}$$

• As we've seen, for a curve, we need to be infinitely close for our line's defining points, yielding

$$\lim_{\Delta x \to 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

• This gives us this instantaneous slope (rate of change) of a function at every point on its domain. The above equation is the definition of the derivative.

6.1.3 Notation

- $\frac{d}{dx}f(x)$ is read "The derivative of f of x with respect to x." You can also say "The instantaneous rate of change in f of x with respect to x."
- If y = f(x), $\frac{dy}{dx}$ is "The derivative of y with respect to x". Warning: Do not try to cancel out the d's, no matter how tempting it is.
- There is the advantage of always specifying the variable with respect to which we're differentiating (it's the one in the denominator).

Lagrange's prime notation: - f'(x) (read: "f prime x") is the derivative of f(x). - This is useful when it is clear which variable were are referring to (e.g., when there's only one).

- What is $\frac{d(x^2)}{dx}$? x^2 $2x^{2-1}$ -2x
- What is $\frac{d(4x^3)}{dx}$?
 - $-4x^{3}$
 - $-4*3x^{3-1}$
 - $-12x^{2}$

Practice

Take the derivative of each of these.

$$x^3$$

$$3x^2$$

$$60x^{11}$$

$$\begin{array}{c}
 x \\
 \frac{4}{x^2} \\
 9\sqrt{x} \\
 6x^{5/2} \\
 11,596,232
 \end{array}$$

Evaluate the derivatives at x = 2 and x = -1

$$x^{3}$$

$$3x^{2}$$

$$60x^{11}$$

$$x$$

$$\frac{4}{x^{2}}$$

$$9\sqrt{x}$$

$$6x^{5/2}$$

$$11,596,232$$

Practice

Take the derivative of each of these.

$$x^{3}$$

$$3x^{2}$$

$$60x^{11}$$

$$x$$

$$\frac{4}{x^{2}}$$

$$9\sqrt{x}$$

$$6x^{5/2}$$

$$11,596,232$$

Evaluate the derivatives at x = 2 and x = -1

$$x^{3}$$

$$3x^{2}$$

$$60x^{11}$$

$$x$$

$$\frac{4}{x^{2}}$$

$$9\sqrt{x}$$

$$6x^{5/2}$$

$$11,596,232$$

6.1.4 Special functions

A few functions have particular rules:

•
$$\frac{d(ln(x))}{dx} = \frac{1}{x}$$

$$\bullet \ \frac{d(log_b(x))}{dx} = \frac{1}{x*ln(b)}$$

$$\bullet \quad \frac{d(e^x)}{dx} = e^x$$

•
$$\frac{d(a^x)}{dx} = a^x ln(a)$$

•
$$\frac{dy}{dx}c = 0$$

•
$$\frac{d(x^x)}{dx} = x^x (1 + \ln(x))$$

6.1.5 Derivatives with addition and substraction

• Easiest rule to remember:

$$\frac{d(f(x) \pm g(x))}{dx} = f'(x) \pm g'(x)$$

Practice

Take the derivative of each of these

$$x^{2} + x + 5$$

$$x^{4} - 4x^{3} + 5x^{2} + 8x - 6$$

$$3x^{5} - 6x^{2}$$

$$5x^{2} + 8\sqrt{x} - \frac{1}{x}$$

$$ln(x) + 5e^{x} - 4x^{3}$$

6.2 Advanced rules

6.2.1 Product rule

• A little more complicated:

$$\frac{d(f(x) \times g(x))}{dx} = f'(x)g(x) + g'(x)f(x)$$

• Example: $2x \times 3x$

Practice

Take the derivative of each of these:

$$x^{3} * x$$

$$e^{x} * x^{2}$$

$$ln(x) * x^{-3}$$

Remember, $\frac{d(f(x)*g(x))}{dx} = f'(x)g(x) + g'(x)f(x)$.

6.2.2 Quotient rule

$$\frac{d\frac{f(x)}{g(x)}}{dx} = \frac{f'(x)g(x) - g'(x)f(x)}{[g(x)]^2}$$

If you're having trouble with this, just apply the product rule to:

$$\frac{d[f(x)*g^{-1}(x)]}{dx}$$

Remember, $\frac{d\frac{f(x)}{g(x)}}{dx} = \frac{f'(x)g(x) - g'(x)f(x)}{[g(x)]^2}.$

6.2.3 Chain rule

$$\frac{d[f(g(x))]}{dx} = f'(g(x)) * g'(x)$$

Let's take the derivative of a function of a function:

$$\frac{d[ln(x^2)]}{dx}$$

$$f(x) = ln(x), g(x) = \frac{x^2}{2}$$

$$f'(x) = \frac{1}{x}, g'(x) = 2x$$

$$\frac{1}{x^2} * 2x = \frac{2}{x}$$

Practice

Take the derivative of each of these:

$$(3x^4 - 8)^2$$
 e^{x^2}

$$e^{x^2}$$

Remember, $\frac{d(f(g(x)))}{dx} = f'(g(x)) * g'(x)$.

6.2.4 Second derivative

- Same process as taking single derivative, except input for second derivative is output from first.
- Second derivative tells us whether the slope of a function is increasing, decreasing, or staying the same at any point x on the function's domain.
- Example: driving a car.
 - -f(x) =distance traveled at time x
 - -f'(x) =speed at time x
 - -f''(x) = acceleration at time x

Graph $f(x) = x^2$, f'(x), and f''(x).

$$\frac{d^2(x^4)}{dx^2} = f''(x^4)$$

- First, we take the first derivative:

$$f'(x^4) = 4x^3$$

- Then we use that output to take the second derivative:

$$f''(x^4) = f'(4x^3) = 12x^2$$

Practice

Take the second derivative of the following functions:

 x^5

 $6x^2$

4ln(x)

3x

 $4x^{3/2}$

6.3 Differentiable and Continuous Functions

- Informally: A function is continuous at a point if its graph has no holes or breaks at that point
- Formally: A function is continuous at a point a if:

$$\lim_{x \to a} f(x) = f(a)$$

Continuity requires 3 conditions to hold:

- f(a) is defined (a is in the domain of f)
- $\lim_{x \to a} f(x)$ exists
- $\lim_{x\to a} f(x) = f(a)$ (the value of f equals the limit of f at a)

Differentiable:

- If f'(x) exists, f is differentiable at x.
- If f is differentiable at every point of an open interval I, f is differentiable on I.
- Graph must have a (non-vertical) tangent line at each point, be relatively smooth, and not contain any breaks, bends, or cusps.
- If a function is differentiable at a point, it is also continuous at that point.
- If a function is continuous at a point, it is *not* necessarily differentiable at that point.

6.3.1 When is f not differentiable?

When does f'(x) not exist?

- When the function is discontinuous at that point.
 - Jump or break in the graph.
- There are different slopes approaching the point from the left and from the right.
 - Corner point
- When the graph of the function has a vertical tangent line at that point.
 - Cusp
 - Vertical inflection point

6.4 Extrema and optimization

Optimization lets us find the minimum or maximum value a function takes.

- Formal theory
 - Utility maximization, continuous choices
- Ordinary Least Squares (OLS)
 - Focuses on minimizing the squared errors between observed data and values predicted by a regression
- Maximum Likelihood Estimation (MLE)
 - Focuses on maximizing a likelihood function, given observed values

6.4.1 Extrema

Informally, a maximum is just the highest value a function takes, and a minimum is the lowest value.

- Easy to identify extrema (maxima or minima) intuitively by looking at a graph of the function.
 - Maxima are high points ("peaks")
 - Minima are low points ("valleys")
- Extrema can be local or global.

6.4.2 Identifying extrema

The derivative of a function gives the rate of change. - When the derivative is zero (or fails to exist), the function has usually reached a (local) maximum or minimum.

• Why?

At a maximum, the function must be increasing before the point and decreasing after it.

At a minimum, the function must be decreasing before the point and increasing after it.

So we'll start by identifying points where this is the case ("critical points" or "stationary points").

A technical note:

A point where f'(x) = 0 or f'(x) does not exist is called a *critical point* (or *stationary point*). Local extrema occur at critical points, but not all critical points are extrema. For instance, sometimes the graph is changing between concave and convex ("inflection points"). Sometimes the function is not differentiable at that point for other reasons, as discussed earlier.

So we can find the local maxima and/or minima of a function by taking the derivative, setting it equal to zero, and solving for x (or whatever).

$$f'(x) = 0$$

This gives us the first-order condition (FOC).

6.4.3 Minimum or maximum?

BUT we don't know if we've found a maximum or minimum, or even if we've found an extremum or just an inflection point.

6.4.4 Second derivatives

The second derivative gives us the rate of change of the rate of change of the original function. So it tells us whether the slope is getting larger or smaller.

$$f(x) = x^{2}$$
$$f'(x) = 2x$$
$$f''(x) = 2$$

Second Derivative Test - Start by identifying f''(x)

- Substitute in the stationary points (x^*) identified from the FOC
- $f''(x^*) > 0$ we have a local minimum
- $f''(x^*) < 0$ we have a local maximum
- $f''(x^*) = 0$ we (may) have an inflection point need to calculate higher-order derivatives (don't worry about this now)

Collectively these give use the Second-Order Condition (SOC).

6.4.5 Local vs. Global Extrema

To find the minimum/maximum on some interval, compare the local min/max to the value of the function at the interval's endpoints.

- To find the global minimum/maximum, check the function's limits as it approaches $+\infty$ and $-\infty$.
- Extreme value theorem: if a real-valued function f is continuous on the closed interval [a,b], then f must attain a (global) maximum and a (global) minimum.

6.5 Partial derivatives

- Can take derivative with respect to different variables
- Notation: For a function fy = (x, z) = xz, we might want to know how the function changes with x:

$$\frac{\partial}{\partial_x} f(x,y) = \frac{\partial_y}{\partial_x} = \partial_x f$$

• Treat all other variables as constants and take derivative with respect to the variable of interest (here x).

How do we take a partial derivative?

Treat all other variables as constants and take derivative with respect to the variable of interest.

From our earlier example:

$$y = f(x, z) = xz$$
$$\frac{\partial_y}{\partial_x} = ?$$

$$y = f(x, z) = xz$$
$$\frac{\partial_y}{\partial_x} = z$$

Why? Because the partial derivative of xz with respect to x treats z as a constant.

What is $\frac{\partial_y}{\partial_z}$?

6.5.1 Application

$$\bullet \quad \frac{\partial (x^2y{+}xy^2{-}x)}{\partial x}$$

• We apply the addition rule to take the derivative of each term with respect to x.

$$\bullet \quad \frac{\partial (x^2y)}{\partial x} + \frac{\partial (xy^2)}{\partial x} + \frac{\partial (-x)}{\partial x}$$

•
$$2xy + y^2 - 1$$

$$\bullet \quad \frac{\partial (x^2y{+}xy^2{-}x)}{\partial y}$$

• We apply the addition rule to take the derivative of each term with respect to y

$$\bullet \quad \frac{\partial (x^2y)}{\partial y} + \frac{\partial (xy^2)}{\partial y} + \frac{\partial (-x)}{\partial y}$$

•
$$x^2 + 2xy$$

Practice

Take the partial derivative with respect to x and to y of the following functions. What would the notation for each look like?

$$3xy - x$$

$$ln(xy)$$

$$x^3 + y^3 + x^4y^4$$

$$e^{xy}$$

6.6 Integrals

6.6.1 Area under a curve

Often we want to find the area under a curve. - Net effect of change - Cumulative density functions (CDFs) - Expected values and utilities

Sometimes this is easy. What's the area under the curve between x = -1 and x = 1 for this function?

$$f(x) = \begin{cases} \frac{1}{3} & \text{for } x \in [0, 3] \\ 0 & \text{otherwise} \end{cases}$$

Hint: We can draw this and look at the graph. Remember $Area = \ell * w$

Sometimes (usually) finding the area under a curve is harder. But this is basically the question behind integration.

6.6.2 Integrals as summation

You're familiar with summation notation.

$$\sum_{i=1}^{n} i$$

But this only works when we have discrete values to add. When we need to add continuously, we have to use something else. Specifically, integrals.

6.6.3 Definite integrals

Let's say we have a function

$$y = x^2$$

And we want to find the area under the curve from x = 0 to x = 1. To find the area we're interested in here, we can use the definite integral.

Generally speaking, the notation looks like this:

$$\int_{x=a}^{b} f(x), dx$$

Here a is the lower limit of integration, b is the upper limit of integration, our function f(x) is our integrand, and x is our variable of integration.

For our question, we're looking for

$$\int_{x=0}^{1} f(x)dx$$

Which will give us a real number denoting the area under the curve of our function $(y = x^2)$ between x = 0 and x = 1.

If f is continuous on [a, b] or bounded on [a, b] with a finite number of discontinuities, then f is integrable on [a, b].

6.6.4 Indefinite integrals

The indefinite integral, or anti-derivative, F(x) is the inverse of the function f'(x).

$$F(x) = \int f(x) \ dx$$

This means if you take the derivative of F(x), you wind up back at f(x).

$$F' = f$$
 or $\frac{dF(x)}{dx} = f(x)$

This process is called anti-differentiation, or indefinite integration.

While the definite integral gives us a real number (the total area under a curve), the indefinite integral gives us a function.

We need the concept of indefinite integrals to help us solve definite integrals.

6.6.5 Solving definite integrals

$$\int_{a}^{b} f(x) \ dx = F(b) - F(a) = F(x) \bigg|_{a}^{b}$$

Constant of integraton

A quick note:

C in the following slides is the called the "constant of integration." We need to add it when we define all antiderivatives (integrals) of a function because the anti-derivative "undoes" the derivative.

Remember that the derivative of any constant is zero. So if we find an integral F(x) whose derivative is f(x), adding (or subtracting) any constant will give us another integral F(x) + C whose derivative is also f(x).

6.6.6 Rules of integration

$$\int_a^a f(x)\ dx = 0$$

$$\int_a^b f(x)\ dx = -\int_b^a f(x)dx$$

$$\int a\ dx = ax + C \text{ where } a \text{ is a constant}$$

$$\int af(x)dx = a\int f(x)\ dx \text{ where } a \text{ is a constant}$$

6.6.7 More rules

$$\int (f(x) + g(x)) dx = \int f(x)dx + \int g(x)dx$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \qquad \forall n \neq -1$$

$$\int x^{-1} dx = \ln|x| + C$$

6.6.8 Solving the problem

Remember our function $y = x^2$ and our goal of finding the area under the curve from x = 0 to x = 1.

• Find the indefinite integral, F(x)

$$-\int x^2 dx$$
$$-\frac{x^3}{3} + C$$

• Evaluate at our lowest and highest points, F(0) and F(1).

$$- F(0) = 0 - F(1) = \frac{1}{3}$$

- Technically 0+C and $\frac{1}{3}+C$, but the C's will fall out in the next step
- Calculate F(1) F(0)

$$- \ \frac{1}{3} - 0 = \frac{1}{3}$$

Practice — indefinite integrals

$$\int x^2 dx$$

$$\int 3x^2 dx$$

$$\int x dx$$

$$\int 3x^2 + 2x - 7 dx$$

$$\int \frac{2}{x} dx$$

Practice — definite integrals

$$\int_{1}^{7} x^{2} dx$$

$$\int_{1}^{10} 3x^{2} dx$$

$$\int_{7}^{7} x dx$$

$$\int_{1}^{5} 3x^{2} + 2x - 7 dx$$

$$\int_{1}^{e} \frac{2}{x} dx$$

6.6.9 Integration by parts

- What if we want to integrate the product of two functions? How to evaluate $\int [f(x)g(x)]dx$?
- There's a formula for that: integration by parts.
- We can derive the formula from the product rule for derivatives, which you already know.

$$\frac{d(f(x)g(x))}{d(x)} = f^{'}(x)g(x) + g^{'}(x)f(x)$$

$$\int \frac{d(f(x)g(x))}{d(x)} dx = \int [f^{'}(x)g(x) + g^{'}(x)f(x)] dx$$

$$f(x)g(x) = \int f^{'}(x)g(x) dx + \int g^{'}(x)f(x) dx$$

- Note that we can't just plug in our two original functions into this formula. We have some work to do first.
- Board example:

$$\int x\sqrt{x}dx$$

7 Probability

7.1 What is probability?

- Frequency with which an event occurs.
 - Typically:

$$Pr(A) = P(A) = \pi(A) = \frac{\text{Number of ways an event can occur}}{\text{Total number of possible outcomes}}$$

- Probability predicts real-world events using theoretical quantities.
 - Formally, it assigns a likelihood of occurrence to each event in sample space
 - We use the probability space triplet (Ω, S, P) , which are the sample space, event space, and probability mapping, respectively.
- We can consider probability as a function that maps $\Omega \to \mathbb{R}$.
- We can conceive it in terms of relative frequency or subjective belief.

7.2 Kolmogorov's axioms

- $Pr(S_i) \in \mathbb{R}, \ 1 \ge Pr(S_i) \ge 0 \quad \forall S_i \in S$
 - Where S is the event space, S_i are events.
 - Probabilities must be non-negative.
- $Pr(\Omega) = 1$
 - Where Ω is the sample space.
 - Something has to happen.
 - Probabilities sum/integrate to 1.

•
$$Pr\left(\bigcup_{i=1}^{\infty}S_i\right) = \sum_{i=1}^{\infty}Pr(S_i) \iff Pr(S_i\cap S_j) = 0 \ \forall i \neq j$$

 The probability of disjoint (mutually exclusive) sets is equal to the sum of their individual probabilities.

7.3 Some definitions

- Random variable: a variable whose value is determined by the outcome of a random process.
 - Sometimes also called a *stochastic variable*.
 - May be discrete or continuous.
- **Distribution** (of a random variable): the set of values the variable might take.
 - Probability mass function / probability density function defines the probability with which each value occurs.
 - Always sums / integrates to 1.
- Realization (of a random variable): a particular value taken by the variable.
- **Population**: the entire set of objects (people, cases, etc.) in which we are interested.
 - Often denoted N.
- **Sample**: a subset of the population we can observe, from which we try to make generalizations about the population.
 - Often denoted n.
- Frequency distribution: a count of how often a variable takes each of its possible values.
 - The number of members of a sample that take each value of a variable.
- **Independent random variables**: two variables are statistically independent if the value of one does not affect the value of the other.
 - Formally, $Pr(A \cap B) = Pr(A)Pr(B)$

7.4 Discrete probability

- A sample space in which there are a (finite or infinite) countable number of outcomes
- Each realization of random process has a discrete probability of occurring.
 - $-f(X=x_i)=P(X=x_i)$ is the probability the variable takes the value x_i .

7.4.1 Probability Mass Function (PMF)

Probability of each occurrence encoded in probability mass function (PMF)

- $0 \le f(x_i) \le 1$
 - Probability of any value occurring must be between 0 and 1.
- $\bullet \ \sum_x f(x_i) = 1$
 - Probabilities of all values must sum to 1.

7.4.2 Discrete distribution

• What's the probability that we'll roll a 3 on one die roll:

$$Pr(y=3) = \frac{1}{6}$$

- If one roll of the die is an "experiment."
- We can think of a 3 as a "success."
- $Y \sim Bernoulli\left(\frac{1}{6}\right)$
- Fair coins are $\sim Bernoulli(.5)$, for example.
- More generally, $Bernoulli(\pi)$.
 - $-\pi$ represents the probability of success.
- Drawing a specific card from a deck:

$$Pr(y = \text{ace of spades}) = \frac{1}{52}$$

• Drawing any card with a specific value from a deck:

$$Pr(y=ace) = \frac{4}{52}$$

• Getting a specific value on two dice rolls:

$$Pr(y=8) = \frac{5}{36}$$

• We can express the probability mass function in tabular format or in a graph.

85

7.5 Continuous probability

- What happens when our outcome is continuous?
- There are an infinite number of outcomes.
- This makes the denominator of our fraction difficult to work with.
- The probability of the whole space must equal 1.
- Even if all events are equally likely, $\frac{1}{\infty} = 0$

7.5.1 Basics

- The domain may not span $-\infty$ to ∞ .
 - Even space between 0 and 1 is infinite.
- The domain is defined as the area under the probability density function.
- Two common examples are the uniform and bell curves.

7.5.2 Probability Density Function (PDF)

- Similar to PMF from before, but for continuous variables.
- Gives the probability a value falls within a particular interval

$$-P[a \le X \le b] = \int_a^b f(x) \, dx$$

- Total area under the curve is 1.
- -P(a < X < b) is the area under the curve between a and b (where b > a).

7.6 Cumulative Density Function (CDF)

7.6.1 Discrete

- Cumulatve density function is probability X will take a value of x or lower.
- PDF is written f(x), and CDF is written F'(x).

$$F_X(x) = Pr(X \le x)$$

- For discrete CDFs, that means summing up over all values.
- What is the probability of rolling a 6 or lower with two dice? F(6) = ?

7.6.2 Continuous

- We can't sum probabilities for continuous distributions (remember the 0 problem).
- Solution: integration

$$F_Y(y) = \int_{-\infty}^y f(y)dy$$

• Examples of uniform distribution.

7.7 Statistics

7.7.1 Introduction

- While probability allows us to make predictions about events using distributions, statistics uses events to make estimates about distributions and variables.
- It is the process of learning from data.
- A statistic is a summary of data, capturing some theoretically-relevant quantity.
- Broad categories of numerical and categorical.

7.7.2 Univariate statistics

- These measure a single variable.
- Readily expressed in graphical form.
- Common examples:
 - Central tendency (mean, median, and mode)
 - Variance

7.7.3 Examples of univariate statistics

• The mean (\bar{x}) is calculated by summing the data, then dividing by the number of observations:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^{n} x_i$$

- The median is found by ordering the observations from highest to lowest and finding the one in the middle.
- The mode is the most common number.

$$x = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 6 & 7 & 8 & 9 \end{bmatrix}$$

• What are the mean, median, and mode of x?

7.7.4 Measures of central tendency

- Mean balances values on either side.
- Median balances observations on either side.
- Mode finds the most typical observation.
- Which is the best? Like most of what you'll learn in statistics, it depends.

7.7.5 Deviations from central tendency

• Consider two data sets:

$$x = \begin{bmatrix} 1 & 1.5 & 2 & 2.5 & 5.5 & 8.5 & 9 & 9.5 & 10 \end{bmatrix}$$

 $y = \begin{bmatrix} 4.5 & 4.8 & 5 & 5.3 & 5.5 & 5.7 & 6 & 6.2 & 6.5 \end{bmatrix}$

- What is the mean of each?
- What is the median of each?
- Are they similar distributions?

7.7.6 Variance

- We use variance to measure the spread of a single variable.
- Formally defined as the squared deviation from the mean (μ) .
- For discrete random variables, it is written $Var(x) = \sigma^2 = \frac{1}{n} \sum_{i=1}^{n} (x_i \mu)^2$
- For continuous random variables, it is written $Var(x) = \sigma^2 = \int (x \mu)^2 f(x) \ dx$

7.7.7 Standard deviation

- Sometimes variance doesn't make sense, either mathematically or conceptually.
 - Not always clear how to interpret "squared deviation from the mean."
- Instead, will frequently see standard deviation, which is square root of variance.
- It is written σ .

7.8 Bivariate statistics

7.8.1 Covariance

- While measures of central tendency and variance/standard deviation provide useful summaries of a single variable, they don't provide insights into relationships between variables
- For that, we need bivariate statistics.
- Most common and straightforward is covariance.
- Colloquially, can think of covariance as measure of linear deviation from mean.
- When values from one variable are above their mean, are values from the other above or below their mean?
- Put another way, if I told you the value of x was high, would you expect values of y to be high or low?
- Formally, it is written as:

$$cov(X,Y) = E(X - E(X))(Y - E(Y)) = E(XY) - E(X)E(Y)$$

• It is important to note that the magnitude is meaningless; only the direction is interpretable.

7.8.2 Correlation

- Correlation is a normalized measure of covariance
- It is calculated as:

$$\rho_{X,Y} = \frac{cov(X,Y)}{\sigma_X \sigma_Y}$$

- It varies between -1 and 1.
- What is correlation of two independent variables?

7.9 Regression

7.9.1 Ordinary least squares

- Ordinary least squares regression (OLS) is probably the most widely-used model in political science.
- It is all about drawing a line through data.

- This allows us to evaluate the relationship (the association) between x on y.
- The dependent variable, y, must be continuous, generally speaking.
- The main question is which line to draw.

Line and equation $(\hat{y_i} = \hat{\beta_0} + \hat{\beta_1} x_i)$ on board

7.9.2 Residuals

- In basically any set of data, no line can pass through every point (observation).
- We will always have make some error in predicting values.
- The error between the line and some point is referred to as the residual.
- If we refer to our predicted value as \hat{y} , then we can calculate the residual for each observation with the following equation:

$$e_i = y_i - \hat{y}_i$$

7.9.3 Finding the right line

- OLS determines the "best" line by minimizing the sum of squared residuals.
- Plug in all the values for the slope amd intercept and calculate the sum of squared residuals for these infinity combinations.
- That is a lot of work.
- The best solution turns out to be calculus.
- We want to minimize the sum of squared residuals with respect to our β 's.

8 Simulations

9 Text analysis

10 Wrap up

10.1 Methods at UT

10.1.1 Required methods courses

- Scope and Methods of Political Science
 - Statistics I (Statistics/linear regression)
- Statistics II (Linear regression and more)
- Statistics III (Maximum likelihood estimation)
 - Only required if your major field is methods

10.1.2 Other methods courses

- Statistics/econometrics:
 - Bayesian Statistics
 - Causal Inference
 - Math Methods for Political Analysis
 - Time Series and Panel Data
 - Panel and Multilevel Analysis

10.1.3 More courses

- Formal Theory
 - Intro to Formal Political Analysis
 - Formal Political Analysis II
 - Formal Theories of International Relations
- Everything else
 - Conceptualization and Measurement
 - Experimental Methods in Political Science
 - Qualitative Methods

- Network Analysis
- Seminar in Field Experiments

10.1.4 Other departments at UT

You can also take courses through the Economics, Mathematics, or Statistics (Statistics and Data Science) departments.

• M.S. in Statistics

Software and Topic Short Courses - R, Python, Stata, etc.

• More info here.

10.1.5 Other resources

Summer programs at UT:

- Short courses in statistics
 - Department sometimes offers scholarships to cover part of the cost.

Summer programs outside UT:

- ICPSR (Inter-university Consortium for Political and Social Research)
 - Ann Arbor, Michigan
- EITM (Empirical Implications of Theoretical Models)
 - Houston and other locations (Michigan, Duke, Berkeley, Emory)
- IQMR (Institute for Qualitative and Multi-Method Research)
 - Syracuse, NY

References