



21563R

I put my root beer in a square cup...  
now it's just beer.

Unit circle: radius = 1





sohcahtoa

$$\sin(\emptyset) = \text{opp} / \text{hyp}$$

$$\sin(\emptyset) = y / 1$$

$$y = \sin(\emptyset)$$

$$\cos(\emptyset) = \text{adj} / \text{hyp}$$

$$\cos(\emptyset) = x / 1$$


$$x = \cos(\emptyset)$$

$$\tan(\emptyset) = \text{opp} / \text{adj}$$

$$\tan(\emptyset) = y / x$$

$$\emptyset = \text{atan}(y / x)$$

if we have an angle and radius,  
we can calculate an x and y position on a circle




```
angle = radians(45);  
x = radius * cos(angle);  
y = radius * sin(angle);
```

Define a radius for a circle

Loop from 0 to 360 (use a for loop) and increment by 10 ( $i+=10$ )

Calculate the angle using  $i$  (remember to convert to radians)

Draw an ellipse at each position using sin and cos




```
x = radius * cos(angle)
y = radius * sin(angle)
```

if we have an x and y, we can get the angle

$$\theta = \text{atan}(y/x)$$

but because arc tangent uses division,  
it would treat  $y=3, x=2$  the same as  $y = -3, x = -2$   
so most languages have a function **atan2** that takes this  
into account

$$\text{angle} = \text{atan2}(\textcolor{brown}{y}, x)$$



```
float angle = atan2(targY-y,targX-x);
```