

POSITIVE MATCHING DECOMPOSITION OF GRAPHS

M. FARROKHI D. G.

JOINT WORK WITH

S. GHARAKHLOO AND A. A. YAZDAN POUR

IASBS

DEPARTMENT OF MATHEMATICS

SAMS 2021

NOVEMBER 29, 2021

PRELIMINARIES

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

- $\{X_1, \dots, X_c\}$ a proper coloring of $\bar{\Gamma}$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ a simple graph
- $\bar{\Gamma}$ the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

- $\{X_1, \dots, X_c\}$ a proper coloring of $\bar{\Gamma}$
- Each vertex $x \in X_i \mapsto e_i \in \mathbb{R}^c$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

- A set of vectors X in \mathbb{R}^d is in *general position* if
any d -subset of X is *linearly independent*.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ a simple graph

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ a simple graph
- \mathbb{K} a field

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ a simple graph
- \mathbb{K} a field
- $S = \mathbb{K}[x_{i,k} : i = 1, \dots, n, k = 1, \dots, d]$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ a simple graph
- \mathbb{K} a field
- $S = \mathbb{K}[x_{i,k} : i = 1, \dots, n, k = 1, \dots, d]$
- $L_{\bar{\Gamma}}^{\mathbb{K}}(d) = (x_{i,1}x_{j,1} + \dots + x_{i,d}x_{j,d} : \{i, j\} \in E(\bar{\Gamma}))$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ a simple graph
- \mathbb{K} a field
- $S = \mathbb{K}[x_{i,k} : i = 1, \dots, n, k = 1, \dots, d]$
- $L_{\bar{\Gamma}}^{\mathbb{K}}(d) = (x_{i,1}x_{j,1} + \dots + x_{i,d}x_{j,d} : \{i, j\} \in E(\bar{\Gamma}))$
- $V(L_{\bar{\Gamma}}^{\mathbb{K}}(d)) = \{\text{orthogonal representations of } \Gamma\}$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

POSITIVE MATCHING DECOMPOSITIONS (PMD)

Definition (**Positive matchings**¹ of graphs)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Definition (Positive matchings¹ of graphs)

- $\Gamma = (V, E)$ a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Definition (**Positive matchings**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- M a matching in Γ

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHINGS

Definition (Positive matchings¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- M a matching in Γ

M is **positive** if $M = \{e \in E: \omega(e) > 0\}$
for some weight function $\omega: V \rightarrow \mathbb{R}$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHINGS

Definition (Positive matchings¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- M a matching in Γ

M is **positive** if $M = \{e \in E : \omega(e) > 0\}$
for some weight function $\omega: V \rightarrow \mathbb{R}$

Remark

\mathbb{R} can be replaced with \mathbb{Z}

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- $\{E_1, \dots, E_p\}$ an edge partition of E into matchings

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- $\{E_1, \dots, E_p\}$ an edge partition of E into matchings
- $E_0 = \emptyset$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- $\{E_1, \dots, E_p\}$ an edge partition of E into matchings
- $E_0 = \emptyset$

(E_1, \dots, E_p) is a **pmd** of Γ if

E_i is a **positive matching** in $\Gamma \setminus E_1 \cup \dots \cup E_{i-1}$,
for all i with $1 \leq i \leq p$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ a simple graph
- $\{E_1, \dots, E_p\}$ an edge partition of E into matchings
- $E_0 = \emptyset$

(E_1, \dots, E_p) is a **pmd** of Γ if

E_i is a **positive matching** in $\Gamma \setminus E_1 \cup \dots \cup E_{i-1}$,
for all i with $1 \leq i \leq p$

$$\text{pmd}(\Gamma) = \min(p: (E_1, \dots, E_p) \text{ a pmd of } \Gamma)$$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

- Γ a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

- Γ a simple graph
- $d \geq \text{pmd}(\Gamma)$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019¹)

- Γ a simple graph
- $d \geq \text{pmd}(\Gamma)$

$L_{\Gamma}^{\mathbb{K}}(d)$ is a *radical complete intersection*

$L_{\Gamma}^{\mathbb{K}}(d)$ is *prime* if $d \geq \text{pmd}(\Gamma) + 1$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

$$(1) \text{ pmd}(\Gamma) \leq 2n - 3,$$

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

Corollary (A. Conca and V. Welker, 2019)

If Γ is a forest with maximum degree $\Delta(\Gamma)$, then

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

Corollary (A. Conca and V. Welker, 2019)

If Γ is a forest with maximum degree $\Delta(\Gamma)$, then

- (1) $L_{\Gamma}^{\mathbb{K}}(d)$ is *radical* for all d ,

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

Corollary (A. Conca and V. Welker, 2019)

If Γ is a forest with maximum degree $\Delta(\Gamma)$, then

- (1) $L_{\Gamma}^{\mathbb{K}}(d)$ is *radical* for all d ,
- (2) $L_{\Gamma}^{\mathbb{K}}(d)$ is a *complete intersection* iff $d \geq \Delta(\Gamma)$,

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

Corollary (A. Conca and V. Welker, 2019)

If Γ is a forest with maximum degree $\Delta(\Gamma)$, then

- (1) $L_{\Gamma}^{\mathbb{K}}(d)$ is *radical* for all d ,
- (2) $L_{\Gamma}^{\mathbb{K}}(d)$ is a *complete intersection* iff $d \geq \Delta(\Gamma)$,
- (3) $L_{\Gamma}^{\mathbb{K}}(d)$ is *prime* iff $d \geq \Delta(\Gamma) + 1$.

POSITIVE MATCHING DECOMPOSITIONS

Definition

Definition

- Γ a simple graph

Definition

- Γ a simple graph
- M a matching of Γ

Definition

- Γ a simple graph
- M a matching of Γ
- $M^c := E(\Gamma) \setminus M$

Definition

- Γ a simple graph
- M a matching of Γ
- $M^c := E(\Gamma) \setminus M$
- W a walk in Γ

POSITIVE MATCHING DECOMPOSITIONS

Definition

- Γ a simple graph
- M a matching of Γ
- $M^c := E(\Gamma) \setminus M$
- W a walk in Γ

W is **alternating** w.r.t. M if

the edges of W **alternate** between M and M^c

Theorem

*The following conditions are equivalent for a **matching** M of a **graph** Γ :*

Theorem

*The following conditions are equivalent for a **matching** M of a **graph** Γ :*

- (1) M is **positive matching**,

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is *pendant* in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :


- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is *pendant* in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is *pendant* in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.



PMD OF GRAPHS

Example

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}.$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$
- $\text{pmd}(C_n) = 3$ for all $n \geq 3$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$
- $\text{pmd}(C_n) = 3$ for all $n \geq 3$

Question

Any characterization of graphs Γ with $\text{pmd}(\Gamma) = 3$?

Remark

$\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ for any graph Γ .

Remark

$\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ for any graph Γ .

Question

Any classification of graphs Γ with $\text{pmd}(\Gamma) = \Delta(\Gamma)$?

GRAPHS ATTAINING THE LOWER BOUND FOR PMD

Remark

$\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ for any graph Γ .

Question

Any classification of graphs Γ with $\text{pmd}(\Gamma) = \Delta(\Gamma)$?

Theorem

Every graph Γ with maximum valency at least three has a *subdivision* Γ' satisfying $\text{pmd}(\Gamma') = \Delta(\Gamma')$.

Definition

Definition

- Γ a simple graph

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or

a corona of corona of ... corona of Γ .

Theorem

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

- Γ a graph

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

- Γ a graph
- Γ' an antler graph of Γ

Definition

- Γ a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or

a corona of corona of ... corona of Γ .

Theorem

- Γ a graph
- Γ' an antler graph of Γ

$$\text{pmd}(\Gamma') = \max\{\text{pmd}(\Gamma), \Delta(\Gamma')\}.$$

Definition (Cactus graph)

A connected graph with any two cycles having at most one vertex in common.

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ a cactus

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ a cactus

$$\Delta(\Gamma) \leq \text{pmd}(\Gamma) \leq \Delta(\Gamma) + 1.$$

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ a cactus


$$\Delta(\Gamma) \leq \text{pmd}(\Gamma) \leq \Delta(\Gamma) + 1.$$

$\text{pmd}(\Gamma) = \Delta(\Gamma)$ if Γ is *triangle-free* and *non-cycle*

CACTUS GRAPHS

Problem

Any characterization of cacti with given pmd ?



A cactus with $\text{pmd} = 4$

COMPLETE MULTIPARTITE GRAPHS

Theorem

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

$$\Delta(\Gamma) = n \text{ and } \text{pmd}(\Gamma) - \Delta(\Gamma) = m.$$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

$$\Delta(\Gamma) = n \text{ and } \text{pmd}(\Gamma) - \Delta(\Gamma) = m.$$

Conjecture

$\text{pmd}(\Gamma) \leq 2\Delta(\Gamma) - 1$ for any graph Γ .

Theorem

Theorem

- $\Gamma = K_{n_1, \dots, n_m} (n_1 \leq \dots \leq n_m)$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)
- $\text{pmd}(K_{2,2,2}) = 6$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)
- $\text{pmd}(K_{2,2,2}) = 6$
- $\text{pmd}(K_{2,2,3}) = 7$

Theorem

Theorem

- Γ a bipartite graph with $\Delta(\Gamma) \geq 3$

Theorem

- Γ a bipartite graph with $\Delta(\Gamma) \geq 3$
- M a perfect matching of Γ

Theorem

- Γ a bipartite graph with $\Delta(\Gamma) \geq 3$
- M a perfect matching of Γ

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma - M) + \Delta(\Gamma) - 1.$$

Corollary

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ *with Γ_1, Γ_2 spanning subgraphs*

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r -regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r -regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- Γ an *r-regular* bipartite graph

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- Γ an *r-regular* bipartite graph

$$\text{pmd}(\Gamma) \leq \binom{r}{2} + 2.$$

Definition (Cayley Graphs)

Definition (Cayley Graphs)

- G a group

Definition (Cayley Graphs)


- G a group
- C an inversed closed subset of G

Definition (Cayley Graphs)

- G a group
- C an inversed closed subset of G

$$\text{Cay}(G, C) = (G, \{\text{edges } \{g, gc\}\})$$

CAYLEY GRAPHS: $\text{Cay}(A_4, \{\alpha^{\pm 1}, \beta\})$



$$\alpha = (1\ 2\ 3) \text{ and } \beta = (1\ 2)(3\ 4)$$

$$a = (), b = (1\ 4)(2\ 3), c = (1\ 2)(3\ 4), \text{ and } d = (1\ 3)(2\ 4)$$

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

(1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

- (1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.
- (2) $H \cdot e$ is a *p.m.* iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

(1) $H \cdot e$ is a **matching** iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.

(2) $H \cdot e$ is a **p.m.** iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

(3) $H \cdot e$ is a **p.m.** iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

(1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.

(2) $H \cdot e$ is a *p.m.* iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

(3) $H \cdot e$ is a *p.m.* iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

(4) $\Gamma[H \cdot e]$ is a *union of edges* iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{and} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

(1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

- (1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.
- (2) the edge sets $H_c \cdot \{x, xc\}$ give rise to a *pmd* of Γ .

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

- (1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.
- (2) the edge sets $H_c \cdot \{x, xc\}$ give rise to a *pmd* of Γ .
- (3) if $I(C) = \text{Involutions}(C)$ and $H \in \{H_c\}_{c \in C}$ has *min. order*, then

$$\text{pmd}(\Gamma) \leq \left(|C| - \frac{1}{2} |I(C)| \right) [G : H].$$

Theorem

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
- $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
 - $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$
- $$\text{pmd}(Q_n) \leq 2n.$$

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
- $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$

$$\text{pmd}(Q_n) \leq 2n.$$

- Merging the last two (suitably chosen) matchings yields

$$\text{pmd}(Q_n) \leq 2n - 1.$$

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
- $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$

$$\text{pmd}(Q_n) \leq 2n.$$

- Merging the last two (suitably chosen) matchings yields

$$\text{pmd}(Q_n) \leq 2n - 1.$$

Conjecture

$$\text{pmd}(Q_n) = 2n - 1$$

THANKS!