

On the center of automorphism group of a group

M. Farrokhi D. G. and M. R. R. Moghaddam

Ferdowsi University of Mashhad, Iran

March 10-12, 2010

Let G be a group.

Let G be a group.

It is clear that $Z(\text{Aut}(G))$ is trivial if $Z(G)$ is trivial.

Let G be a group.

It is clear that $Z(\text{Aut}(G))$ is trivial if $Z(G)$ is trivial.

Thus it is natural to ask what happens when $Z(G)$ is not trivial, i.e., how the structure of $Z(\text{Aut}(G))$ depends on the structure of $Z(G)$?

Let G be a group.

It is clear that $Z(\text{Aut}(G))$ is trivial if $Z(G)$ is trivial.

Thus it is natural to ask what happens when $Z(G)$ is not trivial, i.e., how the structure of $Z(\text{Aut}(G))$ depends on the structure of $Z(G)$?

In this talk we assume that $Z(G)$ is cyclic and we obtain the structure of $Z(\text{Aut}(G))$.

Let G be a group.

It is clear that $Z(\text{Aut}(G))$ is trivial if $Z(G)$ is trivial.

Thus it is natural to ask what happens when $Z(G)$ is not trivial, i.e., how the structure of $Z(\text{Aut}(G))$ depends on the structure of $Z(G)$?

In this talk we assume that $Z(G)$ is cyclic and we obtain the structure of $Z(\text{Aut}(G))$.

As a result we will show that $\exp(Z(\text{Aut}(G))) \leq \exp(Z(G))$ while $|Z(\text{Aut}(G))|$ might be greater than $|Z(G)|$. (exp. QD_{16})

Let $Aut_c(G) = C_{Aut(G)}(Inn(G))$ be the group of central automorphisms of G . Then $g^{-1}\theta(g) \in Z(G)$ for each $\theta \in Aut_c(G)$ and $g \in G$. In particular $Z(Aut(G)) \subseteq Aut_c(G)$.

Definitions

Let $Aut_c(G) = C_{Aut(G)}(Inn(G))$ be the group of central automorphisms of G . Then $g^{-1}\theta(g) \in Z(G)$ for each $\theta \in Aut_c(G)$ and $g \in G$. In particular $Z(Aut(G)) \subseteq Aut_c(G)$.

Definition

Let G be a group with cyclic center $Z(G) = \langle z \rangle$ and $\theta \in Aut_c(G)$. Then $\bar{\theta}$ is the homomorphism from G to $Z(G)$, which sends g to $g^{-1}\theta(g)$ for each $g \in G$. Also α_θ is the smallest nonnegative integer k such that $\bar{\theta}(z) = z^k$.

Definitions

Let $Aut_c(G) = C_{Aut(G)}(Inn(G))$ be the group of central automorphisms of G . Then $g^{-1}\theta(g) \in Z(G)$ for each $\theta \in Aut_c(G)$ and $g \in G$. In particular $Z(Aut(G)) \subseteq Aut_c(G)$.

Definition

Let G be a group with cyclic center $Z(G) = \langle z \rangle$ and $\theta \in Aut_c(G)$. Then $\bar{\theta}$ is the homomorphism from G to $Z(G)$, which sends g to $g^{-1}\theta(g)$ for each $g \in G$. Also α_θ is the smallest nonnegative integer k such that $\bar{\theta}(z) = z^k$.

Fact

In the above definition α_θ is independent of the choice of z .

An elementary result

Utilizing the definition of α_θ we obtain:

An elementary result

Utilizing the definition of α_θ we obtain:

Lemma

Let G be a group with cyclic center. If $\theta \in \text{Aut}_c(G)$ and $g \in G$, then

$$(a) \quad \bar{\theta}^k(g) = \bar{\theta}(g)^{\alpha_\theta^{k-1}};$$

for each nonnegative integer k .

An elementary result

Utilizing the definition of α_θ we obtain:

Lemma

Let G be a group with cyclic center. If $\theta \in \text{Aut}_c(G)$ and $g \in G$, then

- (a) $\bar{\theta}^k(g) = \bar{\theta}(g)^{\alpha_\theta^{k-1}};$
- (b) $\theta^k(g) = g^{\binom{k}{0}} \bar{\theta}(g)^{\binom{k}{1}} \dots \bar{\theta}^{k-1}(g)^{\binom{k}{k-1}} \bar{\theta}^k(g)^{\binom{k}{k}},$ and

for each nonnegative integer k .

An elementary result

Utilizing the definition of α_θ we obtain:

Lemma

Let G be a group with cyclic center. If $\theta \in \text{Aut}_c(G)$ and $g \in G$, then

- (a) $\bar{\theta}^k(g) = \bar{\theta}(g)^{\alpha_\theta^{k-1}};$
- (b) $\theta^k(g) = g^{\binom{k}{0}} \bar{\theta}(g)^{\binom{k}{1}} \dots \bar{\theta}^{k-1}(g)^{\binom{k}{k-1}} \bar{\theta}^k(g)^{\binom{k}{k}},$ and
- (c) $\theta^k(g) = g \bar{\theta}(g)^{\frac{1}{\alpha_\theta}((1+\alpha_\theta)^k - 1)},$

for each nonnegative integer k .

Lemma

Let G be a group with cyclic center of order n . Then for all $\varphi, \psi \in Z(\text{Aut}(G))$,

(a) $\alpha_{\varphi\psi} + 1 \overset{n}{\equiv} (\alpha_{\varphi} + 1)(\alpha_{\psi} + 1)$, and

Lemma

Let G be a group with cyclic center of order n . Then for all $\varphi, \psi \in Z(\text{Aut}(G))$,

- (a) $\alpha_{\varphi\psi} + 1 \overset{n}{\equiv} (\alpha_{\varphi} + 1)(\alpha_{\psi} + 1)$, and*
- (b) The map $\alpha^* : Z(\text{Aut}(G)) \longrightarrow \text{Aut}(Z(G)) \cong U(\mathbb{Z}_n)$ given by $\alpha^*(\varphi) = \alpha_{\varphi} + 1$ is a homomorphism, where $\alpha_{\varphi} + 1$ is identified with the automorphism, which sends z to $z^{\alpha_{\varphi}+1}$.*

Lemma

Let G be a group with cyclic center of finite order $n = p_1^{a_1} \cdots p_m^{a_m}$ and $\varphi \in Z(\text{Aut}(G))$. Then

$$|\varphi| \mid \text{lcm}(d_1, \dots, d_m),$$

where $d_i = p_i^{a_i}$, when $p_i \mid \alpha_\varphi$ and $d_i = p_i^{a_i-1}(p_i - 1)$, when $p_i \nmid \alpha_\varphi$. In particular $\exp(Z(\text{Aut}(G))) \leq \exp(Z(G))$.

Main theorem: Groups with finite cyclic center

Theorem

Let G be a group with cyclic center of finite order $n = p_1^{a_1} \cdots p_m^{a_m}$. Then $Z(\text{Aut}(G)) \cong A_1 \times \cdots \times A_m$, where the subgroup A_i is isomorphic to

(a) the trivial group;

for $i = 1, 2, \dots, n$.

Main theorem: Groups with finite cyclic center

Theorem

Let G be a group with cyclic center of finite order $n = p_1^{a_1} \cdots p_m^{a_m}$. Then $Z(\text{Aut}(G)) \cong A_1 \times \cdots \times A_m$, where the subgroup A_i is isomorphic to

- (a) the trivial group;*
- (b) an abelian p_i -group with $\exp(A_i) \mid p_i^{a_i}$, or*

for $i = 1, 2, \dots, m$.

Main theorem: Groups with finite cyclic center

Theorem

Let G be a group with cyclic center of finite order $n = p_1^{a_1} \cdots p_m^{a_m}$. Then $Z(\text{Aut}(G)) \cong A_1 \times \cdots \times A_m$, where the subgroup A_i is isomorphic to

- (a) the trivial group;*
 - (b) an abelian p_i -group with $\exp(A_i) \mid p_i^{a_i}$, or*
 - (c) the cyclic group of order $p_i^{a_i-1}(p_i - 1)$,*
- for $i = 1, 2, \dots, m$.*

Proof: Groups with finite cyclic center

Let $Z(G) = P_1 \times \cdots \times P_m$, $\varphi \in Z(\text{Aut}(G))$ and $g \in G$.

Proof: Groups with finite cyclic center

Let $Z(G) = P_1 \times \cdots \times P_m$, $\varphi \in Z(\text{Aut}(G))$ and $g \in G$.

Then $\bar{\varphi}(g) = \bar{\varphi}_1(g) \cdots \bar{\varphi}_m(g)$, where $\bar{\varphi}_i(g) \in P_i$.

Proof: Groups with finite cyclic center

Let $Z(G) = P_1 \times \cdots \times P_m$, $\varphi \in Z(\text{Aut}(G))$ and $g \in G$.

Then $\bar{\varphi}(g) = \bar{\varphi}_1(g) \cdots \bar{\varphi}_m(g)$, where $\bar{\varphi}_i(g) \in P_i$.

Let φ_i be the following map

$$\begin{aligned}\varphi_i : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}_i(g).\end{aligned}$$

Proof: Groups with finite cyclic center

Let $Z(G) = P_1 \times \cdots \times P_m$, $\varphi \in Z(\text{Aut}(G))$ and $g \in G$.

Then $\bar{\varphi}(g) = \bar{\varphi}_1(g) \cdots \bar{\varphi}_m(g)$, where $\bar{\varphi}_i(g) \in P_i$.

Let φ_i be the following map

$$\begin{aligned}\varphi_i : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}_i(g).\end{aligned}$$

Then $\varphi_i \in Z(\text{Aut}(G))$ so that

$$Z(\text{Aut}(G)) = A_1 \times \cdots \times A_m,$$

where $\bar{\psi}(g) \in P_i$, for each $\psi \in A_i$.

Proof: Groups with finite cyclic center

If $\alpha_i^* = \alpha^*|_{A_i}$, then

$$\frac{A_i}{\text{Ker}\alpha_i^*} \cong B_i \leq U(P_i) \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Proof: Groups with finite cyclic center

If $\alpha_i^* = \alpha^*|_{A_i}$, then

$$\frac{A_i}{\text{Ker}\alpha_i^*} \cong B_i \leq U(P_i) \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Now if $\exp(A_i) \nmid p_i^{a_i}$, then $p_i > 2$ and $\exp(A_i) | p_i^{a_i-1}(p_i-1)$ so that there exists an automorphism $\varphi \in A_i \setminus \{I\}$ such that $|\varphi| | p_i - 1$.

Proof: Groups with finite cyclic center

If $\alpha_i^* = \alpha^*|_{A_i}$, then

$$\frac{A_i}{\text{Ker}\alpha_i^*} \cong B_i \leq U(P_i) \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Now if $\exp(A_i) \nmid p_i^{a_i}$, then $p_i > 2$ and $\exp(A_i) | p_i^{a_i-1}(p_i-1)$ so that there exists an automorphism $\varphi \in A_i \setminus \{I\}$ such that $|\varphi| | p_i - 1$.

Let $n_i = n/p_i^{a_i}$, $z_i = z^{n_i}$ and $\alpha_\varphi^i = \alpha_\varphi/n_i$.

Proof: Groups with finite cyclic center

If $\alpha_i^* = \alpha^*|_{A_i}$, then

$$\frac{A_i}{\text{Ker}\alpha_i^*} \cong B_i \leq U(P_i) \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Now if $\exp(A_i) \nmid p_i^{a_i}$, then $p_i > 2$ and $\exp(A_i) | p_i^{a_i-1}(p_i-1)$ so that there exists an automorphism $\varphi \in A_i \setminus \{I\}$ such that $|\varphi| | p_i - 1$.

Let $n_i = n/p_i^{a_i}$, $z_i = z^{n_i}$ and $\alpha_\varphi^i = \alpha_\varphi/n_i$.

Then $P_i = \langle z_i \rangle$, $n_i | \alpha_\varphi$ and $p_i \nmid \alpha_\varphi$.

Proof: Groups with finite cyclic center

If $\alpha_i^* = \alpha^*|_{A_i}$, then

$$\frac{A_i}{\text{Ker}\alpha_i^*} \cong B_i \leq U(P_i) \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Now if $\exp(A_i) \nmid p_i^{a_i}$, then $p_i > 2$ and $\exp(A_i) | p_i^{a_i-1}(p_i-1)$ so that there exists an automorphism $\varphi \in A_i \setminus \{I\}$ such that $|\varphi| | p_i - 1$.

Let $n_i = n/p_i^{a_i}$, $z_i = z^{n_i}$ and $\alpha_\varphi^i = \alpha_\varphi/n_i$.

Then $P_i = \langle z_i \rangle$, $n_i | \alpha_\varphi$ and $p_i \nmid \alpha_\varphi$.

Hence α_φ^i is an integer coprime to p_i and there exists an integer β such that $1 + \alpha_\varphi^i \beta$ is a primitive root modulo $p_i^{a_i}$.

Proof: Groups with finite cyclic center

Let

$$\begin{aligned}\varphi_\beta : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}(g)^\beta.\end{aligned}$$

Proof: Groups with finite cyclic center

Let

$$\begin{aligned}\varphi_\beta : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}(g)^\beta.\end{aligned}$$

It is straitforward to show that $\varphi_\beta \in Z(\text{Aut}(G))$ is an automorphism of order $p_i^{a_i-1}(p_i - 1)$.

Proof: Groups with finite cyclic center

Let

$$\begin{aligned}\varphi_\beta : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}(g)^\beta.\end{aligned}$$

It is straitforward to show that $\varphi_\beta \in Z(\text{Aut}(G))$ is an automorphism of order $p_i^{a_i-1}(p_i - 1)$.

In particular $P_i \cap \text{Ker}\varphi_\beta = \langle 1 \rangle$ and $G = P_i \text{Ker}\varphi_\beta \cong P_i \times \text{Ker}\varphi_\beta$.

Proof: Groups with finite cyclic center

Let

$$\begin{aligned}\varphi_\beta : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}(g)^\beta.\end{aligned}$$

It is straightforward to show that $\varphi_\beta \in Z(\text{Aut}(G))$ is an automorphism of order $p_i^{a_i-1}(p_i - 1)$.

In particular $P_i \cap \text{Ker}\varphi_\beta = \langle 1 \rangle$ and $G = P_i \text{Ker}\varphi_\beta \cong P_i \times \text{Ker}\varphi_\beta$.

Clearly $\psi \in A_i$ is of order 2 if and only if $\alpha_\psi^i \equiv -2 \pmod{p_i^{a_i}}$.

Proof: Groups with finite cyclic center

Let

$$\begin{aligned}\varphi_\beta : G &\longrightarrow G \\ g &\longmapsto g\bar{\varphi}(g)^\beta.\end{aligned}$$

It is straitforward to show that $\varphi_\beta \in Z(\text{Aut}(G))$ is an automorphism of order $p_i^{a_i-1}(p_i - 1)$.

In particular $P_i \cap \text{Ker}\varphi_\beta = \langle 1 \rangle$ and $G = P_i \text{Ker}\varphi_\beta \cong P_i \times \text{Ker}\varphi_\beta$.

Clearly $\psi \in A_i$ is of order 2 if and only if $\alpha_\psi^i \equiv -2 \pmod{p_i^{a_i}}$.

If $\psi \in A_i$ is an automorphism of order 2 and $\theta \in \text{Ker}\alpha_i^*$, then $\alpha_{\psi\theta}^i \equiv -2 \pmod{p_i^{a_i}}$.

Proof: Groups with finite cyclic center

On the other hand θ is a p_i -element and so $|\psi||\theta| = |\psi\theta| = 2$.

Proof: Groups with finite cyclic center

On the other hand θ is a p_i -element and so $|\psi||\theta| = |\psi\theta| = 2$.

Hence $\theta = I$ and consequently $\text{Ker}\alpha_i^* = \langle I \rangle$.

Proof: Groups with finite cyclic center

On the other hand θ is a p_i -element and so $|\psi||\theta| = |\psi\theta| = 2$.

Hence $\theta = I$ and consequently $\text{Ker}\alpha_i^* = \langle I \rangle$.

Therefore

$$A_i \cong \mathbb{Z}_{p_i^{a_i-1}(p_i-1)}.$$

Examples: Groups with finite cyclic center

Example

Let $G = \langle a, b : a^p = b^p = [a, b]^p = 1, [a, b]^a = [a, b]^b = [a, b] \rangle$ be the p -group of order p^3 and exponent p , where p is a prime. Then $Z(\text{Aut}(G)) = \langle I \rangle$.

Examples: Groups with finite cyclic center

Example

Let $G = \langle a, b : a^p = b^p = [a, b]^p = 1, [a, b]^a = [a, b]^b = [a, b] \rangle$ be the p -group of order p^3 and exponent p , where p is a prime. Then $Z(\text{Aut}(G)) = \langle I \rangle$.

Example

Let $G = \langle a, b : a^{p^2} = b^p = 1, a^b = a^{p+1} \rangle$ be the p -group of order p^3 and exponent p^2 , where p is a prime. Then $Z(\text{Aut}(G)) \cong \mathbb{Z}_p$.

Examples: Groups with finite cyclic center

Example

Let $G = \langle a, b : a^p = b^p = [a, b]^p = 1, [a, b]^a = [a, b]^b = [a, b] \rangle$ be the p -group of order p^3 and exponent p , where p is a prime. Then $Z(\text{Aut}(G)) = \langle I \rangle$.

Example

Let $G = \langle a, b : a^{p^2} = b^p = 1, a^b = a^{p+1} \rangle$ be the p -group of order p^3 and exponent p^2 , where p is a prime. Then $Z(\text{Aut}(G)) \cong \mathbb{Z}_p$.

Example

If $G = \mathbb{Z}_{p^n}$, then $Z(\text{Aut}(G)) \cong \mathbb{Z}_{p^{n-1}(p-1)}$.

Corollary: Nilpotent groups with finite cyclic center

Corollary

Let G be a finite nilpotent group with cyclic center of order $n = p_1^{a_1} \cdots p_m^{a_m}$. Then either the Sylow p_i -subgroup of G is cyclic or the subgroup A_i in the previous theorem is isomorphic to

(a) the trivial group, or

for $i = 1, 2, \dots, n$.

Corollary: Nilpotent groups with finite cyclic center

Corollary

Let G be a finite nilpotent group with cyclic center of order $n = p_1^{a_1} \cdots p_m^{a_m}$. Then either the Sylow p_i -subgroup of G is cyclic or the subgroup A_i in the previous theorem is isomorphic to

- (a) the trivial group, or*
 - (b) an abelian p_i -group with $\exp(A_i) \mid p_i^{a_i}$,*
- for $i = 1, 2, \dots, n$.*

Main theorem: Groups with infinite cyclic center

Theorem

Let G be a group with infinite cyclic center. Then $Z(\text{Aut}(G))$ is isomorphic to

- (a) the trivial group;*

Main theorem: Groups with infinite cyclic center

Theorem

Let G be a group with infinite cyclic center. Then $Z(\text{Aut}(G))$ is isomorphic to

- (a) the trivial group;*
- (b) the cyclic group of order 2, or*

Main theorem: Groups with infinite cyclic center

Theorem

Let G be a group with infinite cyclic center. Then $Z(\text{Aut}(G))$ is isomorphic to

- (a) the trivial group;*
- (b) the cyclic group of order 2, or*
- (c) a nontrivial torsion-free abelian group.*

Examples: Groups with infinite cyclic center

Example

Let $G = \langle a, b, c : [a, c] = [b, c] = 1 \rangle$ be a group with infinite cyclic center. Then $Z(\text{Aut}(G)) = \langle I \rangle$.

Examples: Groups with infinite cyclic center

Example

Let $G = \langle a, b, c : [a, c] = [b, c] = 1 \rangle$ be a group with infinite cyclic center. Then $Z(\text{Aut}(G)) = \langle I \rangle$.

Example

If $G = \mathbb{Z}$, then $Z(\text{Aut}(G)) \cong \mathbb{Z}_2$.

Problem

Is there a group G with infinite cyclic center such that $Z(\text{Aut}(G))$ is a nontrivial torsion-free abelian group?

Problem

Is there a group G with infinite cyclic center such that $Z(\text{Aut}(G))$ is a nontrivial torsion-free abelian group?

Problem

Is it true that $\exp(Z(\text{Aut}(G))) \leq \exp(Z(G))$ for any group G ?

Problem

Is there a group G with infinite cyclic center such that $Z(\text{Aut}(G))$ is a nontrivial torsion-free abelian group?

Problem

Is it true that $\exp(Z(\text{Aut}(G))) \leq \exp(Z(G))$ for any group G ?

Problem

Suppose that $\gcd(p_i - 1, |A_i|) \neq 1$. Is it true that $p_i - 1 \parallel |A_i|$ or even A_i has an element of order $p_i - 1$?

Thank You!