

POSITIVE MATCHING DECOMPOSITION OF GRAPHS

M. FARROKHI D. G.

IASBS

DEPARTMENT OF MATHEMATICS

AS-SMS

JANUARY 07, 2026

PRELIMINARIES

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

- $\{X_1, \dots, X_c\}$ a proper coloring of $\bar{\Gamma}$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

- $[n] = \{1, \dots, n\}$
- $\Gamma = ([n], E)$ is a simple graph
- $\bar{\Gamma}$ is the complement of Γ

Definition (Orthogonal representation¹ of Γ in \mathbb{R}^d)

- Each vertex $i \in [n] \mapsto$ a vector $u_i \in \mathbb{R}^d$
- $u_i^T u_j = 0$ whenever $\{i, j\} \in E(\bar{\Gamma})$

Example

- $\{X_1, \dots, X_c\}$ a proper coloring of $\bar{\Gamma}$
- Each vertex $x \in X_i \mapsto e_i \in \mathbb{R}^c$

¹L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* **25** (1979), 1–7.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

Theorem (Lovász, Saks, and Schrijver, 1989¹, 2000²)

The followings are equivalent:

- (1) Γ is $(n - d)$ -connected,
- (2) Γ has an *orthogonal representation* in \mathbb{R}^d in general position.

- A set of vectors X in \mathbb{R}^d is in *general position* if
any d -subset of X is *linearly independent*.

¹L. Lovász, M. Saks, and A. Schrijver, Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **114/115** (1989), 439–454.

²L. Lovász, M. Saks, and A. Schrijver, A correction: Orthogonal representations and connectivity of graphs, *Linear Algebra Appl.* **313** (2000), 101–105.

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ is a simple graph

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ is a simple graph
- \mathbb{K} is a field

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ is a simple graph
- \mathbb{K} is a field
- $S = \mathbb{K}[x_{i,k} : i = 1, \dots, n, k = 1, \dots, d]$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ is a simple graph
- \mathbb{K} is a field
- $S = \mathbb{K}[x_{i,k}: i = 1, \dots, n, k = 1, \dots, d]$
- $L_{\bar{\Gamma}}^{\mathbb{K}}(d) = (x_{i,1}x_{j,1} + \dots + x_{i,d}x_{j,d}: \{i, j\} \in E(\bar{\Gamma}))$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

ORTHOGONAL REPRESENTATIONS

- $S = \mathbb{R}[x_1, \dots, x_d]$
- $V(\text{ideal } I) = \{\text{zeros of } I\} \subseteq \mathbb{R}^d$ is the variety of I

Definition (**Lovász-Saks-Schrijver ideal** of Γ in $\mathbb{K}[x_1, \dots, x_d]$)

- $\Gamma = (V, E)$ is a simple graph
- \mathbb{K} is a field
- $S = \mathbb{K}[x_{i,k} : i = 1, \dots, n, k = 1, \dots, d]$
- $L_{\bar{\Gamma}}^{\mathbb{K}}(d) = (x_{i,1}x_{j,1} + \dots + x_{i,d}x_{j,d} : \{i, j\} \in E(\bar{\Gamma}))$
- $V(L_{\bar{\Gamma}}^{\mathbb{K}}(d)) = \{\text{orthogonal representations of } \Gamma\}$

¹L. Lovász, *Graphs and Geometry*, American Mathematical Society Colloquium Publications, **65**. American Mathematical Society, Providence, RI, 2019. (**Chapter 10**)

POSITIVE MATCHING DECOMPOSITIONS (PMD)

Definition (**Positive matchings**¹ of graphs)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Definition (**Positive matchings**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHINGS

Definition (**Positive matchings**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- M is a matching in Γ

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHINGS

Definition (Positive matchings¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- M is a matching in Γ

M is **positive** if $M = \{e \in E: \omega(e) > 0\}$

for some weight function $\omega: V \rightarrow \mathbb{R}$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHINGS

Definition (Positive matchings¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- M is a matching in Γ

M is **positive** if $M = \{e \in E : \omega(e) > 0\}$
for some weight function $\omega: V \rightarrow \mathbb{R}$

Remark

\mathbb{R} can be replaced with \mathbb{Z}

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- $\{M_1, \dots, M_p\}$ is an edge partition of E into matchings

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- $\{M_1, \dots, M_p\}$ is an edge partition of E into matchings
- $M_0 = \emptyset$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- $\{M_1, \dots, M_p\}$ is an edge partition of E into matchings
- $M_0 = \emptyset$

M_1, \dots, M_p is a **pmd** of Γ if

M_i is a **positive matching** in $\Gamma \setminus M_1 \cup \dots \cup M_{i-1}$,
for all i with $1 \leq i \leq p$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition (**Positive matching decompositions**¹ of graphs)

- $\Gamma = (V, E)$ is a simple graph
- $\{M_1, \dots, M_p\}$ is an edge partition of E into matchings
- $M_0 = \emptyset$

M_1, \dots, M_p is a **pmd** of Γ if

M_i is a **positive matching** in $\Gamma \setminus M_1 \cup \dots \cup M_{i-1}$,
for all i with $1 \leq i \leq p$

$$\text{pmd}(\Gamma) = \min(p: M_1, \dots, M_p \text{ is a pmd of } \Gamma)$$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Example

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}.$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$
- $\text{pmd}(C_n) = 3$ for all $n \geq 3$

Example

- $\text{pmd}(\Gamma) \leq |E(\Gamma)|$
- $\text{pmd}(\Gamma) = |E(\Gamma)|$ iff $\Gamma = K_{1,s} \cup tK_1$
- $\text{pmd}(\Gamma) = \max\{\text{pmd}(C) : C \text{ is a connected component of } \Gamma\}$.
- $\text{pmd}(\Gamma) = 1$ iff $\Gamma = sK_2 \cup tK_1$
- $\text{pmd}(\Gamma) = 2$ iff $\Gamma = P_{n_1} \cup \dots \cup P_{n_k}$ with $\max_i(n_i) > 1$
- $\text{pmd}(C_n) = 3$ for all $n \geq 3$

Question

Any characterization of graphs Γ with $\text{pmd}(\Gamma) = 3$?

Theorem (A. Conca and V. Welker, 2019¹)

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

- Γ is a simple graph

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

- Γ is a simple graph
- $d \geq \text{pmd}(\Gamma)$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019¹)

- Γ is a simple graph
- $d \geq \text{pmd}(\Gamma)$

$L_{\Gamma}^{\mathbb{K}}(d)$ is a *radical* and *complete intersection*

$L_{\Gamma}^{\mathbb{K}}(d)$ is *prime* if $d \geq \text{pmd}(\Gamma) + 1$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

Theorem (A. Conca and V. Welker, 2019¹)

If Γ is a simple graph of order $n > 1$, then

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019¹)

If Γ is a simple graph of order $n > 1$, then

$$(1) \text{ pmd}(\Gamma) \leq 2n - 3,$$

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019¹)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Theorem (A. Conca and V. Welker, 2019¹)

If Γ is a simple graph of order $n > 1$, then

- (1) $\text{pmd}(\Gamma) \leq 2n - 3$,
- (2) $\text{pmd}(\Gamma) \leq n - 1$ if Γ is a *bipartite graph*,
- (3) $\text{pmd}(\Gamma) \geq \Delta(\Gamma)$ with equality if Γ is a *forest*.

¹A. Conca and V. Welker, Lovász-Saks-Schrijver ideals and coordinate sections of determinantal varieties, *Algebra Number Theory* **13**(2) (2019), 455–484.

POSITIVE MATCHING DECOMPOSITIONS

Definition

Definition

- Γ is a simple graph

Definition

- Γ is a simple graph
- M is a matching of Γ

Definition

- Γ is a simple graph
- M is a matching of Γ
- $M^c := E(\Gamma) \setminus M$

Definition

- Γ is a simple graph
- M is a matching of Γ
- $M^c := E(\Gamma) \setminus M$
- W is a walk in Γ

POSITIVE MATCHING DECOMPOSITIONS

Definition

- Γ is a simple graph
- M is a matching of Γ
- $M^c := E(\Gamma) \setminus M$
- W is a walk in Γ

W is **alternating** w.r.t. M if

the edges of W **alternate** between M and M^c

Theorem

*The following conditions are equivalent for a **matching** M of a **graph** Γ :*

Theorem

*The following conditions are equivalent for a **matching** M of a **graph** Γ :*

- (1) M is **positive matching**,

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :

- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is *pendant* in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a *matching* M of a *graph* Γ :


- (1) M is *positive matching*,
- (2) $\Gamma[M]$ has no *alternating closed walk* w.r.t. M ,
- (3) $\Gamma[M']$ has a *pendant edge* belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is *pendant* in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.

POSITIVE MATCHING DECOMPOSITIONS

Theorem

The following conditions are equivalent for a **matching** M of a **graph** Γ :

- (1) M is **positive matching**,
- (2) $\Gamma[M]$ has no **alternating closed walk** w.r.t. M ,
- (3) $\Gamma[M']$ has a **pendant edge** belonging to M' for any $M' \subseteq M$,
- (4) $M = \{e_1, \dots, e_n\}$ s.t. e_i is **pendant** in $\Gamma[\{e_1, \dots, e_i\}]$, for all i with $1 \leq i \leq n$.



TREE-LIKE GRAPHS

Definition

Definition

- Γ is a simple graph

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

- Γ is a graph

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

- Γ is a graph
- Γ' is an antler graph of Γ

Definition

- Γ is a simple graph
- A **corona graph** of Γ : Attach pendants to vertices of Γ .
- An **antler graph** (or **hartshorne graph**) of Γ : Attach trees to vertices of Γ or
a corona of corona of ... corona of Γ .

Theorem

- Γ is a graph
- Γ' is an antler graph of Γ

$$\text{pmd}(\Gamma') = \max\{\text{pmd}(\Gamma), \Delta(\Gamma')\}.$$

Definition (Cactus graph)

A connected graph with any two cycles having at most one vertex in common.

CACTUS GRAPHS

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ is a cactus

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ is a cactus

$$\Delta(\Gamma) \leq \text{pmd}(\Gamma) \leq \Delta(\Gamma) + 1.$$

Definition (Cactus graph)

A connected graph with any **two cycles** having at most **one vertex** in common.

Theorem

- Γ is a cactus


$$\Delta(\Gamma) \leq \text{pmd}(\Gamma) \leq \Delta(\Gamma) + 1.$$

$\text{pmd}(\Gamma) = \Delta(\Gamma)$ if Γ is *triangle-free* and *non-cycle*

CACTUS GRAPHS

Problem

Any characterization of cacti with given pmd?



A cactus with $\text{pmd} = 4$

MULTIPARTITE GRAPHS

Theorem

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

$$\Delta(\Gamma) = n \text{ and } \text{pmd}(\Gamma) - \Delta(\Gamma) = m.$$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\text{pmd}(K_n) = 2n - 3$ for all $n \geq 2$.
- $\text{pmd}(K_{m,n}) = m + n - 1$ for all $m, n \geq 1$.

Remark

- $\Gamma = K_{m+1,n}$ ($m < n$)

$$\Delta(\Gamma) = n \text{ and } \text{pmd}(\Gamma) - \Delta(\Gamma) = m.$$

Conjecture

$\text{pmd}(\Gamma) \leq 2\Delta(\Gamma) - 1$ for any graph Γ .

Theorem

Theorem

- $\Gamma = K_{n_1, \dots, n_m} (n_1 \leq \dots \leq n_m)$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)
- $\text{pmd}(K_{2,2,2}) = 6$

COMPLETE MULTIPARTITE GRAPHS

Theorem

- $\Gamma = K_{n_1, \dots, n_m}$ ($n_1 \leq \dots \leq n_m$)

$$\max \left\{ \frac{3}{2}|\Gamma| - n_m - 1, |\Gamma| + \frac{m}{2} - 2 \right\} \leq \text{pmd}(\Gamma) \leq 2|\Gamma| - n_{m-1} - n_m - 1.$$

Corollary

- $\text{pmd}(K_{1,m,n}) = m + n + 1$
- $\text{pmd}(K_{2,m,n}) = m + n + 2 + \varepsilon$ ($\varepsilon \in \{0, 1\}$)
- $\text{pmd}(K_{2,2,2}) = 6$
- $\text{pmd}(K_{2,2,3}) = 7$

Theorem

Theorem

- Γ is a bipartite graph with $\Delta(\Gamma) \geq 3$

Theorem

- Γ is a bipartite graph with $\Delta(\Gamma) \geq 3$
- M is a perfect matching of Γ

Theorem

- Γ is a bipartite graph with $\Delta(\Gamma) \geq 3$
- M is a perfect matching of Γ

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma - M) + \Delta(\Gamma) - 1.$$

Corollary

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ *with Γ_1, Γ_2 spanning subgraphs*

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r -regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r -regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- Γ is an *r-regular* bipartite graph

Corollary

- $\Gamma = \Gamma_1 \cup \Gamma_2$ with Γ_1, Γ_2 spanning subgraphs
- Γ_1 is *r-regular*
- Γ_2 is non-empty and $\Delta(\Gamma_2) = s$

$$\text{pmd}(\Gamma) \leq \text{pmd}(\Gamma_2) + \binom{r}{2} + rs.$$

Corollary

- Γ is an *r-regular* bipartite graph

$$\text{pmd}(\Gamma) \leq \binom{r}{2} + 2.$$

CARTESIAN PRODUCT OF GRAPHS

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)

- $\Gamma_1 \square \Gamma_2 = (V, E)$,

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)

- $\Gamma_1 \square \Gamma_2 = (V, E)$,
- $V = V_1 \times V_2$,

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)

- $\Gamma_1 \square \Gamma_2 = (V, E)$,
- $V = V_1 \times V_2$,
- $E = (E(\Gamma_1) \times V_2) \cup (V_1 \times E(\Gamma_2))$,

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)


- $\Gamma_1 \square \Gamma_2 = (V, E)$,
- $V = V_1 \times V_2$,
- $E = (E(\Gamma_1) \times V_2) \cup (V_1 \times E(\Gamma_2))$,

CARTESIAN PRODUCTS

- $\Gamma_1 = (V_1, E_1)$,
- $\Gamma_2 = (V_2, E_2)$.

Definition (**Cartesian Product** of Γ_1 and Γ_2)

- $\Gamma_1 \square \Gamma_2 = (V, E)$,
- $V = V_1 \times V_2$,
- $E = (E(\Gamma_1) \times V_2) \cup (V_1 \times E(\Gamma_2))$,



CARTESIAN PRODUCTS

Theorem

CARTESIAN PRODUCTS

Theorem

- Γ_1, Γ_2 *are graphs*

CARTESIAN PRODUCTS

Theorem

- Γ_1, Γ_2 *are graphs*

Theorem

- Γ_1, Γ_2 are graphs

$$\text{pmd}(\Gamma_1 \square \Gamma_2) \leq \min\{\text{pmd}(\Gamma_i)\chi(\Gamma_j) + \text{pmd}(\Gamma_j) : \{i, j\} = \{1, 2\}\}.$$

Corollary

Corollary

- Γ is a *graph*

Corollary

- Γ is a graph
- $n \geq 1$

Corollary

- Γ is a graph
- $n \geq 1$

Corollary

- Γ is a graph
- $n \geq 1$

$$\text{pmd}(\square^n \Gamma) \leq ((n - 1)\chi(\Gamma) + 1) \cdot \text{pmd}(\Gamma).$$

Corollary

- Γ is a graph
- $n \geq 1$

$$\text{pmd}(\square^n \Gamma) \leq ((n - 1)\chi(\Gamma) + 1) \cdot \text{pmd}(\Gamma).$$

Corollary

$$\text{pmd}(Q_n) \leq 2n - 1.$$

Definition (**Forest decomposition** of graphs)

Definition (Forest decomposition of graphs)

- Γ is a graph

Definition (Forest decomposition of graphs)

- Γ is a graph
- $\{F_1, \dots, F_n\}$ is an edge-partition of Γ

Definition (Forest decomposition of graphs)

- Γ is a graph
- $\{F_1, \dots, F_n\}$ is an edge-partition of Γ

Definition (Forest decomposition of graphs)

- Γ is a graph
- $\{F_1, \dots, F_n\}$ is an edge-partition of Γ

F_1, \dots, F_n is a **forest decomposition** of Γ

$(\Gamma - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ is a **forest**,
for all $i = 1, \dots, n$.

CARTESIAN PRODUCTS

Theorem

Theorem

- Γ_1, Γ_2 *are graphs*

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a forest decomposition of Γ_1

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a forest decomposition of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a forest decomposition of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a forest decomposition of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$

$$\text{pmd}(\Gamma_1 \square \Gamma_2) \leq n \cdot \text{pmd}(\Gamma_2) + \sum_{i=1}^n \max\{\Delta(F'_i), \chi(\Gamma_2)\}.$$

Corollary

Corollary

- T_1, \dots, T_n are trees

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

CARTESIAN PRODUCTS

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

$$\text{pmd}(T_1 \square \dots \square T_n) \leq \Delta(T_1 \square \dots \square T_n) + m - \delta_{m \neq 0},$$

Corollary

CARTESIAN PRODUCTS

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

$$\text{pmd}(T_1 \square \dots \square T_n) \leq \Delta(T_1 \square \dots \square T_n) + m - \delta_{m \neq 0},$$

Corollary

- T_1, \dots, T_n are trees

CARTESIAN PRODUCTS

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

$$\text{pmd}(T_1 \square \dots \square T_n) \leq \Delta(T_1 \square \dots \square T_n) + m - \delta_{m \neq 0},$$

Corollary

- T_1, \dots, T_n are trees
- $T_1, \dots, T_n \not\cong K_2$

CARTESIAN PRODUCTS

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

$$\text{pmd}(T_1 \square \dots \square T_n) \leq \Delta(T_1 \square \dots \square T_n) + m - \delta_{m \neq 0},$$

Corollary

- T_1, \dots, T_n are trees
- $T_1, \dots, T_n \not\cong K_2$

CARTESIAN PRODUCTS

Corollary

- T_1, \dots, T_n are trees
- $m := \#\{i: T_i \cong K_2\}$

$$\text{pmd}(T_1 \square \dots \square T_n) \leq \Delta(T_1 \square \dots \square T_n) + m - \delta_{m \neq 0},$$

Corollary

- T_1, \dots, T_n are trees
- $T_1, \dots, T_n \not\cong K_2$

$$\begin{aligned}\text{pmd}(T_1 \square \dots \square T_n) &= \Delta(T_1 \square \dots \square T_n), \\ \text{pmd}(T_1 \square \dots \square T_n \square K_2) &= \Delta(T_1 \square \dots \square T_n \square K_2),\end{aligned}$$

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}
- \mathcal{P} is a **set-theoretical property** on subsets of $2^{\mathcal{S}}$

CARTESIAN PRODUCTS

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}
- \mathcal{P} is a **set-theoretical property** on subsets of $2^{\mathcal{S}}$
- $\mathcal{C} = \{\mathcal{C}_i\}_{i \in I}$ a family of **subsets** of \mathcal{F}

CARTESIAN PRODUCTS

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}
- \mathcal{P} is a **set-theoretical property** on subsets of $2^{\mathcal{S}}$
- $\mathcal{C} = \{\mathcal{C}_i\}_{i \in I}$ a family of **subsets** of \mathcal{F}

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}
- \mathcal{P} is a **set-theoretical property** on subsets of $2^{\mathcal{S}}$
- $\mathcal{C} = \{\mathcal{C}_i\}_{i \in I}$ a family of **subsets** of \mathcal{F}

\mathcal{C} is a \mathcal{P} -cover of \mathcal{S} by \mathcal{F} if

$$\mathcal{S} = \cup_{i \in I} (\cup \mathcal{C}_i) \text{ and } \mathcal{C} \text{ is a } \mathcal{P}\text{-subset of } 2^{\mathcal{S}}.$$

CARTESIAN PRODUCTS

Definition (\mathcal{P} -covers and \mathcal{P} -covering numbers)

- \mathcal{S} is a **multiset**
- $\mathcal{F} = \{\mathcal{S}_1, \dots, \mathcal{S}_k\}$ is a family of **submultisets** of \mathcal{S}
- \mathcal{P} is a **set-theoretical property** on subsets of $2^{\mathcal{S}}$
- $\mathcal{C} = \{\mathcal{C}_i\}_{i \in I}$ a family of **subsets** of \mathcal{F}

\mathcal{C} is a \mathcal{P} -cover of \mathcal{S} by \mathcal{F} if

$$\mathcal{S} = \bigcup_{i \in I} (\bigcup \mathcal{C}_i) \text{ and } \mathcal{C} \text{ is a } \mathcal{P}\text{-subset of } 2^{\mathcal{S}}.$$

\mathcal{P} -covering number of \mathcal{S} by \mathcal{F} is

$$\text{cov}_{\mathcal{P}}(\mathcal{S}, \mathcal{F}) = \min\{|\mathcal{C}| : \mathcal{C} \text{ is a } \mathcal{P}\text{-cover of } \mathcal{S} \text{ by } \mathcal{F}\}$$

CARTESIAN PRODUCTS

Definition

Definition

- Γ is a graph

Definition

- Γ is a graph
- \mathcal{E} is a family of subsets of $E(\Gamma)$

Definition

- Γ is a graph
- \mathcal{E} is a family of subsets of $E(\Gamma)$

Definition

- Γ is a graph
- \mathcal{E} is a family of subsets of $E(\Gamma)$

\mathcal{E} is an **acyclic family of edge-sets** (AFE) if

$\bigcup_{E \in \mathcal{E}} \Gamma[E]$ is acyclic.

Theorem

Theorem

- Γ_1, Γ_2 *are graphs*

Theorem

- Γ_1, Γ_2 are graphs
- $\mathcal{M}_i : M_1^i, \dots, M_{p_i}^i$ is a *pmd* of Γ_i

Theorem

- Γ_1, Γ_2 are graphs
- $\mathcal{M}_i : M_1^i, \dots, M_{p_i}^i$ is a *pmd* of Γ_i
- $C_1^i, \dots, C_{\chi_i}^i$ are *color classes* of a proper coloring of Γ_i

Theorem

- Γ_1, Γ_2 are graphs
- $\mathcal{M}_i : M_1^i, \dots, M_{p_i}^i$ is a *pmd* of Γ_i
- $C_1^i, \dots, C_{\chi_i}^i$ are *color classes* of a proper coloring of Γ_i
- $p_i \leq \chi_j$ ($i \neq j$)

Theorem

- Γ_1, Γ_2 are graphs
- $\mathcal{M}_i : M_1^i, \dots, M_{p_i}^i$ is a *pmd* of Γ_i
- $C_1^i, \dots, C_{\chi_i}^i$ are *color classes* of a proper coloring of Γ_i
- $p_i \leq \chi_j$ ($i \neq j$)

Theorem

- Γ_1, Γ_2 are graphs
- $\mathcal{M}_i : M_1^i, \dots, M_{p_i}^i$ is a *pmd* of Γ_i
- $C_1^i, \dots, C_{\chi_i}^i$ are *color classes* of a proper coloring of Γ_i
- $p_i \leq \chi_j$ ($i \neq j$)

$$\text{pmd}(\Gamma_1 \square \Gamma_2) \leq \text{cov}_{\text{AFE}}(\chi_j * \Gamma_i, \mathcal{M}_i) + p_j.$$

Theorem

Theorem

- Γ_1, Γ_2 *are graphs*

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a *forest decomposition* of Γ_1

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a *forest decomposition* of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a *forest decomposition* of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$
- $\sum_{i=1}^m \Delta(F'_i) \leq |\Gamma_2|$.

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a *forest decomposition* of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$
- $\sum_{i=1}^m \Delta(F'_i) \leq |\Gamma_2|$.

Theorem

- Γ_1, Γ_2 are graphs
- F_1, \dots, F_n is a *forest decomposition* of Γ_1
- $F'_i := (\Gamma_1 - E(F_1) \cup \dots \cup E(F_{i-1}))[V(F_i)]$ for $i = 1, \dots, n$
- $\sum_{i=1}^m \Delta(F'_i) \leq |\Gamma_2|$.

$$\text{pmd}(\Gamma_1 \square \Gamma_2) \leq \text{pmd}(\Gamma_2) + n \cdot \max \left\{ \sum_{i=1}^n \Delta(F'_i), \chi(\Gamma_2) \right\}.$$

Corollary

Corollary

- *T is a tree*

Corollary

- T is a tree
- Γ is a graph satisfying $\Delta(T) \leq |\Gamma|$

Corollary

- T is a tree
- Γ is a graph satisfying $\Delta(T) \leq |\Gamma|$

Corollary

- T is a tree
- Γ is a graph satisfying $\Delta(T) \leq |\Gamma|$

$$\text{pmd}(\Gamma \square T) \leq \text{pmd}(\Gamma) + \max\{\Delta(T), \chi(\Gamma)\}.$$

Corollary

- T is a tree
- Γ is a graph satisfying $\Delta(T) \leq |\Gamma|$

$$\text{pmd}(\Gamma \square T) \leq \text{pmd}(\Gamma) + \max\{\Delta(T), \chi(\Gamma)\}.$$

Conjecture

There exists a constant $c > 0$ such that

$$\text{pmd}(\Gamma_1) + \text{pmd}(\Gamma_2) \leq \text{pmd}(\Gamma_1 \square \Gamma_2) + c$$

for all graphs Γ_1 and Γ_2 .

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

Conjecture

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

Conjecture

- $m, n \geq 3$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

Conjecture

- $m, n \geq 3$
- n is even and $n < 4(m - 1)$ or n is odd and $n < 2m$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$

In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.

Conjecture

- $m, n \geq 3$
- n is even and $n < 4(m - 1)$ or n is odd and $n < 2m$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$4 \leq \text{pmd}(P_m \square C_n) \leq 5.$$


In addition, $\text{pmd}(P_m \square C_n) = 4$ if

- n is even and $n \geq 4(m - 1)$, or
- n is odd and $n > 2m$.


Conjecture

- $m, n \geq 3$
 - n is even and $n < 4(m - 1)$ or n is odd and $n < 2m$
- $$\text{pmd}(P_m \square C_n) = 5.$$

CARTESIAN PRODUCTS




n is odd




n is even

CARTESIAN PRODUCTS



$n > 2m$ is odd



$n \geq 4(m - 1)$ is even

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- *$m + n$ is odd, and*

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

Conjecture

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

Conjecture

- $m, n \geq 3$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

Conjecture

- $m, n \geq 3$
- $m + n$ is even or $(m, n) = (3, 4), (3, 6), (5, 6)$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

Conjecture

- $m, n \geq 3$
- $m + n$ is even or $(m, n) = (3, 4), (3, 6), (5, 6)$

CARTESIAN PRODUCTS

Theorem

If $m, n \geq 3$, then

$$5 \leq \text{pmd}(C_m \square C_n) \leq 6.$$

In addition, $\text{pmd}(C_m \square C_n) = 5$ if

- $m + n$ is odd, and
- $\{m, n\} \neq \{3, 4\}, \{3, 6\}, \{5, 6\}$.

Conjecture


- $m, n \geq 3$
- $m + n$ is even or $(m, n) = (3, 4), (3, 6), (5, 6)$

$$\text{pmd}(C_m \square C_n) = 6.$$

CARTESIAN PRODUCTS

Theorem

$$\text{pmd}(C_3 \square C_3) = \text{pmd}(C_4 \square C_4) = 6.$$



CAYLEY GRAPHS

Definition (Cayley Graphs)

Definition (Cayley Graphs)

- G is a group

Definition (Cayley Graphs)


- G is a group
- C is an inversed closed subset of G

Definition (Cayley Graphs)

- G is a group
- C is an inversed closed subset of G

$$\text{Cay}(G, C) = (G, \{\text{edges } \{g, gc\}\})$$

CAYLEY GRAPHS: $\text{Cay}(A_4, \{\alpha^{\pm 1}, \beta\})$



$\alpha = (1\ 2\ 3)$ and $\beta = (1\ 2)(3\ 4)$
 $a = ()$, $b = (1\ 4)(2\ 3)$, $c = (1\ 2)(3\ 4)$, and $d = (1\ 3)(2\ 4)$

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

(1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

- (1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.
- (2) $H \cdot e$ is a *p.m.* iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

- (1) $H \cdot e$ is a **matching** iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.
- (2) $H \cdot e$ is a **p.m.** iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

- (3) $H \cdot e$ is a **p.m.** iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

Lemma

If $\Gamma = \text{Cay}(G, C)$, $c \in C$, $e = \{x, xc\}$, and $H = H_c \leq G$, then

- (1) $H \cdot e$ is a *matching* iff $c \notin H^x$. In this case, $|H \cdot e| = |H|$.
- (2) $H \cdot e$ is a *p.m.* iff $H = \{h_1, \dots, h_n\}$ s.t. for any $j > 1$, either

$$(h_i^{-1}h_j)^x \notin C \cup cC \quad \text{or} \quad (h_i^{-1}h_j)^{xc} \notin C \cup c^{-1}C$$

for all $i < j$.

- (3) $H \cdot e$ is a *p.m.* iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

- (4) $\Gamma[H \cdot e]$ is a *union of edges* iff

$$H^x \cap (C \cup cC) = \{1\} \quad \text{and} \quad H^{xc} \cap (C \cup c^{-1}C) = \{1\}.$$

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

(1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

- (1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.
- (2) the edge sets $H_c \cdot \{x, xc\}$ give rise to a *pmd* of Γ .

Theorem

If $\Gamma = \text{Cay}(G, C)$ and $\{H_c\}_{c \in C}$ is a family of subgroups of G satisfying $H_c = H_{c^{-1}}$, and

$$H_c^x \cap (C \cup cC) = \{1\} \quad \text{or} \quad H_c^{xc} \cap (C \cup c^{-1}C) = \{1\}$$

for all $c \in C$ and $x \in G$, then

- (1) the edge sets $H_c \cdot \{x, xc\}$ *partition* $E(\Gamma)$.
- (2) the edge sets $H_c \cdot \{x, xc\}$ give rise to a *pmd* of Γ .
- (3) if $I(C) = \text{Involutions}(C)$ and $H \in \{H_c\}_{c \in C}$ has *min. order*, then

$$\text{pmd}(\Gamma) \leq \left(|C| - \frac{1}{2} |I(C)| \right) [G : H].$$

Theorem

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
- $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
 - $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$
- $$\text{pmd}(Q_n) \leq 2n.$$

Theorem

- $Q_n = \text{Cay}(\mathbb{Z}_2^n, \{e_1, \dots, e_n\})$ the hypercube of dimension n
- $H_{e_i} = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j = x_i = 0 \right\}$ for all i with $1 \leq i \leq n$

$$\text{pmd}(Q_n) \leq 2n.$$

- Merging the last two (suitably chosen) matchings yields

$$\text{pmd}(Q_n) \leq 2n - 1.$$

THANKS!