

On the probability of being a 2-Engel group

A. Erfanian and M. Farrokhi D. G.

Ferdowsi University of Mashhad

Johor Bahru

14-18 February 2011

Definition

If G is a finite group, then the *commutativity degree* of G , denoted by $d(G)$, is the probability that two randomly chosen elements $x, y \in G$ commute, i.e.,

$$d(G) = \frac{|\{(x, y) \in G \times G : xy = yx\}|}{|G|^2}.$$

Definition

If G is a finite group, then the *commutativity degree* of G , denoted by $d(G)$, is the probability that two randomly chosen elements $x, y \in G$ commute, i.e.,

$$d(G) = \frac{|\{(x, y) \in G \times G : xy = yx\}|}{|G|^2}.$$

Theorem

If G is a finite group, then

$$d(G) = \frac{k(G)}{|G|},$$

where $k(G)$ is the number of conjugacy classes of G .

Theorem

Let G be a finite group. Then

Theorem

Let G be a finite group. Then

- 1 if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);

Theorem

Let G be a finite group. Then

- ① if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);
- ② if $d(G) > \frac{1}{2}$, then G is nilpotent (P. Lescot, 1987);

Theorem

Let G be a finite group. Then

- ① if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);
- ② if $d(G) > \frac{1}{2}$, then G is nilpotent (P. Lescot, 1987);
- ③ if $d(G) = \frac{1}{2}$ and G is not nilpotent, then $\frac{G}{Z(G)} \cong S_3$ (P. Lescot, 1988);

Theorem

Let G be a finite group. Then

- ① if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);
- ② if $d(G) > \frac{1}{2}$, then G is nilpotent (P. Lescot, 1987);
- ③ if $d(G) = \frac{1}{2}$ and G is not nilpotent, then $\frac{G}{Z(G)} \cong S_3$ (P. Lescot, 1988);
- ④ if $d(G) > \frac{1}{3}$, the G is supersolvable (F. Barry, D. MacHale and Á. NíShé, 2006);

Theorem

Let G be a finite group. Then

- ① if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);
- ② if $d(G) > \frac{1}{2}$, then G is nilpotent (P. Lescot, 1987);
- ③ if $d(G) = \frac{1}{2}$ and G is not nilpotent, then $\frac{G}{Z(G)} \cong S_3$ (P. Lescot, 1988);
- ④ if $d(G) > \frac{1}{3}$, then G is supersolvable (F. Barry, D. MacHale and Á. NíShé, 2006);
- ⑤ if $d(G) > \frac{11}{75}$ and $|G|$ is odd, then G is supersolvable (F. Barry, D. MacHale and Á. NíShé, 2006);

Theorem

Let G be a finite group. Then

- ① if G is a non-abelian group, then $d(G) \leq \frac{5}{8}$ and $d(G) = \frac{5}{8}$ if and only if $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ (Gustafson, 1973);
- ② if $d(G) > \frac{1}{2}$, then G is nilpotent (P. Lescot, 1987);
- ③ if $d(G) = \frac{1}{2}$ and G is not nilpotent, then $\frac{G}{Z(G)} \cong S_3$ (P. Lescot, 1988);
- ④ if $d(G) > \frac{1}{3}$, then G is supersolvable (F. Barry, D. MacHale and Á. NíShé, 2006);
- ⑤ if $d(G) > \frac{11}{75}$ and $|G|$ is odd, then G is supersolvable (F. Barry, D. MacHale and Á. NíShé, 2006);
- ⑥ if $d(G) > \frac{3}{40}$, then G is solvable and $d(G) = \frac{1}{12}$ if and only if $G \cong A_5 \times B$, where B is abelian (R. M. Guralnick and G. R. Robinson, 2006)

Definition

If G is a finite group, then the n -Engel degree of G , denoted by $E_n(G)$, is the probability that two randomly chosen elements $x, y \in G$ satisfy the equation $[y, {}_n x] = 1$, i.e.,

$$E_n(G) = \frac{|\{(x, y) \in G \times G : [y, {}_n x] = 1\}|}{|G|^2}.$$

Definition

If G is a finite group, then the n -Engel degree of G , denoted by $E_n(G)$, is the probability that two randomly chosen elements $x, y \in G$ satisfy the equation $[y, {}_n x] = 1$, i.e.,

$$E_n(G) = \frac{|\{(x, y) \in G \times G : [y, {}_n x] = 1\}|}{|G|^2}.$$

Definition

Let G be a group and $x \in G$. Then

$$E_G(x) = \{y \in G : [y, x, x] = 1\}.$$

Lemma

Let G be a group and $x \in G$. Then the following statements are equivalent:

Lemma

Let G be a group and $x \in G$. Then the following statements are equivalent:

① $E_G(x) \leq G$;

Lemma

Let G be a group and $x \in G$. Then the following statements are equivalent:

- ① $E_G(x) \leq G$;
- ② $[E_G(x), x, E_G(x), x] = 1$;

Lemma

Let G be a group and $x \in G$. Then the following statements are equivalent:

- ① $E_G(x) \leq G$;
- ② $[E_G(x), x, E_G(x), x] = 1$;
- ③ $[[E_G(x), x], [E_G(x), x]] = 1$.

Lemma

Let G be a group and $x \in G$. Then the following statements are equivalent:

- ① $E_G(x) \leq G$;
- ② $[E_G(x), x, E_G(x), x] = 1$;
- ③ $[[E_G(x), x], [E_G(x), x]] = 1$.

Definition

A finite group G is called 3-metabelian if every subgroup of G that is generated by three elements is metabelian. In other words, G is 3-metabelian if and only if $[[x, y], [x, z]] = 1$ for all elements $x, y, z \in G$.

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.
Then

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.
Then

① $G'^4 = 1$;

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.

Then

- ① $G'^4 = 1$;
- ② $G''^2 = 1$;

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.

Then

- ① $G'^4 = 1$;
- ② $G''^2 = 1$;
- ③ $[\gamma_3(G), \gamma_2(G)] = 1$;

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.

Then

- ① $G'^4 = 1$;
- ② $G''^2 = 1$;
- ③ $[\gamma_3(G), \gamma_2(G)] = 1$;
- ④ $[\gamma_2(G), \gamma_2(G), G] = 1$;

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.

Then

- ① $G'^4 = 1$;
- ② $G''^2 = 1$;
- ③ $[\gamma_3(G), \gamma_2(G)] = 1$;
- ④ $[\gamma_2(G), \gamma_2(G), G] = 1$;
- ⑤ $[[x_1, x_2], [x_3, x_4]] = [[x_{\pi_1}, x_{\pi_2}], [x_{\pi_3}, x_{\pi_4}]]$;

Known results

Let G be a 3-metabelian group, $x_1, x_2, x_3, x_4 \in G$ and $\pi \in S_4$.

Then

- ① $G'^4 = 1$;
- ② $G''^2 = 1$;
- ③ $[\gamma_3(G), \gamma_2(G)] = 1$;
- ④ $[\gamma_2(G), \gamma_2(G), G] = 1$;
- ⑤ $[[x_1, x_2], [x_3, x_4]] = [[x_{\pi_1}, x_{\pi_2}], [x_{\pi_3}, x_{\pi_4}]]$;
- ⑥ $[[x_1, x_2], [x_3, x_4]] = [x_1, x_2, x_3, x_4][x_1, x_2, x_4, x_3]^{-1}$.

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

① $E_G(x) \leq G$;

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;
- ③ $[C_G(x)x^G : C_G(x)] = [G : E_G(x)]$;

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;
- ③ $[C_G(x)x^G : C_G(x)] = [G : E_G(x)]$;
- ④ $|C_G(x)x^G| = [G : C_G(x) \cap x^G]$ divides $|G|$.

Elementary Properties

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;
- ③ $[C_G(x)x^G : C_G(x)] = [G : E_G(x)]$;
- ④ $|C_G(x)x^G| = [G : C_G(x) \cap x^G]$ divides $|G|$.

Lemma

Let G be a finite 3-metabelian group with an element x such that $C_G(x)x^G = G$. Then

Elementary Properties

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;
- ③ $[C_G(x)x^G : C_G(x)] = [G : E_G(x)]$;
- ④ $|C_G(x)x^G| = [G : C_G(x) \cap x^G]$ divides $|G|$.

Lemma

Let G be a finite 3-metabelian group with an element x such that $C_G(x)x^G = G$. Then

- ① $G = [x, G] \rtimes C_G(x)$;

Elementary Properties

Lemma

Let G be a finite 3-metabelian group and $x \in G$. Then

- ① $E_G(x) \leq G$;
- ② $|E_G(x)| = |C_G(x)| |G_G(x) \cap x^G|$;
- ③ $[C_G(x)x^G : C_G(x)] = [G : E_G(x)]$;
- ④ $|C_G(x)x^G| = [G : C_G(x) \cap x^G]$ divides $|G|$.

Lemma

Let G be a finite 3-metabelian group with an element x such that $C_G(x)x^G = G$. Then

- ① $G = [x, G] \rtimes C_G(x)$;
- ② if $x \in L(G)$, then $x \in Z(G)$.

Theorem

Let G be a finite 3-metablian group, which is not 2-Engel group and $p = \min \pi(G)$. Then

$$E_2(G) \leq \frac{1}{p} + \left(1 - \frac{1}{p}\right) \frac{|L_2(G)|}{|G|}$$

and if $L_2(G) \leq G$, then

$$E_2(G) \leq \frac{2p-1}{p^2}.$$

Moreover, both of the upper bounds are sharp at any prime p .

Theorem

Let G be a finite 3-metablian group, which is not 2-Engel group and $p = \min \pi(G)$. Then

$$E_2(G) \geq d(G) - (p-1) \frac{|Z(G)|}{|G|} + (p-1) \frac{k_G(L(G))}{|G|}$$

and if either G is a p -group, or G' is a cyclic 2-group or a generalized quaternion 2-group, then

$$E_2(G) \geq pd(G) - (p-1) \frac{|Z(G)|}{|G|}.$$

Moreover, both of the lower bounds are sharp at any prime p .

Example

Let $G = D_{2n} = \langle a, b : a^n = b^2 = 1, a^b = a^{-1} \rangle$ be a dihedral group of order $2n$. Then

$$E_2(G) = \begin{cases} \frac{n+1}{2n}, & n \text{ odd,} \\ \frac{n+2}{2n}, & n = 2m, m \text{ odd,} \\ \frac{n+4}{2n}, & n = 4m. \end{cases}$$

Examples

Example

Let $G = D_{2n} = \langle a, b : a^n = b^2 = 1, a^b = a^{-1} \rangle$ be a dihedral group of order $2n$. Then

$$E_2(G) = \begin{cases} \frac{n+1}{2n}, & n \text{ odd,} \\ \frac{n+2}{2n}, & n = 2m, m \text{ odd,} \\ \frac{n+4}{2n}, & n = 4m. \end{cases}$$

Example

Let $G = Q_{4n} = \langle a, b : a^{2n} = 1, a^n = b^2, a^b = a^{-1} \rangle$ be a generalized quaternion group of order $4n$. Then

$$E_2(G) = \begin{cases} \frac{n+1}{2n}, & n \text{ odd,} \\ \frac{n+2}{2n}, & n \text{ even.} \end{cases}$$

Example






Let $p > 2$ be a prime and

$$G = \langle a, b, c : a^{p^2} = b^p = c^p = 1, [a, b] = a^p, [a, c] = b, [b, c] = 1 \rangle.$$







Then

$$E_2(G) = \frac{2p-1}{p^2}.$$





References I

-  S. Bachmuth and J. Lewin, *The Jacobi identity in groups*, Math. Z. **83** (1964), 170-176.
-  F. Barry, D. MacHale and Á. NíShé, Some supersolvability conditions for finite groups, *Math. Proc. R. Ir. Acad.*, **106A**(2) (2006), 163-177.
-  P. Erdős and P. Turan, *On some problems of statistical group theory*, Acta Math. Acad. Sci. Hung. **19** (1968), 413-435.
-  M. Farrokhi D. G. and M. R. R. Moghaddam, *On groups satisfying a symmetric Engel word*, Submitted.
-  W. H. Gustafson, *What is the probability that two groups elements commute?*, Amer. Math. Monthly **80** (1973), 1031-1304.

References II

-  W. P. Kappe, *Die A-norm einer gruppe*, Illinois J. Math. **5** (1961), 187-197.
-  P. Lescot, Isoclinism classes and commutativity degrees of finite groups, *J. Algebra*, **177** (1995), 847-869.
-  P. Lescot, *Sur certains groupes finis*, Rev. Math. Spécials, **8** (1987), 276-277.
-  P. Lescot, *Degré de commutativité et structure d'un groupe fini*, Rev. Math. Spécials, **8** (1988), 276-277.
-  I. D. Macdonald, *On certain varieties of groups*, Math. Z. **76** (1961), 270-282.
-  I. D. Macdonald, *Another law for the 3-metabelian groups*, J. Aust. Math. Soc. **6**(4) (1964), 452-453.

References III

-  B. H. Neumann, *On a conjecture of Hanna Neumann*, Proc. Glasgow Math. Assoc. **3** (1956), 13-17.
-  R. M. Guralnick and G. R. Robinson, *On the commuting probability in finite groups*, J. Algebra, **300** (2006), 509-528.
-  G. J. Sherman, *What is the probability an automorphism fixes a group element?*, Amer. Math. Monthly **82** (1975), 261-264.
-  The GAP Group, *GAP-Groups, Algorithms and Programming, Version 4.4.12, 2008*
(<http://www.gap-system.org/>).