

Finite groups with a given number of elements of each order

M. Farrokhi D. G.
(join work with F. Saeedi)

Department of Pure Mathematics, Ferdowsi University of Mashhad

6th Group Theory Conference of Iran
Golestan University
March 12-13, 2014

$$\omega_d(G)$$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d(G)$ ► The number of elements of order d

$$\omega_d^*(G)$$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$ ► $\{\omega_d(G) : d \in \omega(G)\}$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$ ► $\{\omega_d(G) : d \in \omega(G)\}$

$\omega(d)$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$ ► $\{\omega_d(G) : d \in \omega(G)\}$

$\omega(d)$ ► $\{\omega_d(G) : G \text{ is a finite group}\}$

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$ ► $\{\omega_d(G) : d \in \omega(G)\}$

$\omega(d)$ ► $\{\omega_d(G) : G \text{ is a finite group}\}$

ω

$\omega_d(G)$ ► The number of elements of order d

$\omega_d^*(G)$ ► The number of solutions of $x^d = 1$

Spectrum $\omega(G)$ ► The set of orders of elements of G

Co-spectrum $\omega^*(G)$ ► $\{\omega_d(G) : d \in \omega(G)\}$

$\omega(d)$ ► $\{\omega_d(G) : G \text{ is a finite group}\}$

ω ► $\bigcup_{d \in \mathbb{N}} \omega(d)$

Examples

G	$\omega(G)$	$\omega^*(G)$
-----	-------------	---------------

Examples

G	$\omega(G)$	$\omega^*(G)$
-----	-------------	---------------

Examples

G	$\omega(G)$	$\omega^*(G)$
Q_8	$\{1, 2, 4\}$	$\{1, 6\}$

Examples

G	$\omega(G)$	$\omega^*(G)$
Q_8	$\{1, 2, 4\}$	$\{1, 6\}$
S_3	$\{1, 2, 3\}$	$\{1, 2, 3\}$

Examples

G	$\omega(G)$	$\omega^*(G)$
Q_8	$\{1, 2, 4\}$	$\{1, 6\}$
S_3	$\{1, 2, 3\}$	$\{1, 2, 3\}$
\mathbb{Z}_p^n	$\{1, p\}$	$\{1, p^n - 1\}$

Examples

G	$\omega(G)$	$\omega^*(G)$
Q_8	$\{1, 2, 4\}$	$\{1, 6\}$
S_3	$\{1, 2, 3\}$	$\{1, 2, 3\}$
\mathbb{Z}_p^n	$\{1, p\}$	$\{1, p^n - 1\}$
$\mathbb{Z}_p^n \times \mathbb{Z}_2$	$\{1, 2, p, 2p\}$	$\{1, p^n - 1\}$

Examples

G	$\omega(G)$	$\omega^*(G)$
Q_8	$\{1, 2, 4\}$	$\{1, 6\}$
S_3	$\{1, 2, 3\}$	$\{1, 2, 3\}$
\mathbb{Z}_p^n	$\{1, p\}$	$\{1, p^n - 1\}$
$\mathbb{Z}_p^n \times \mathbb{Z}_2$	$\{1, 2, p, 2p\}$	$\{1, p^n - 1\}$
$\mathbb{Z}_p \rtimes \mathbb{Z}_q$	$\{1, p, q\}$	$\{1, p - 1, p(q - 1)\}$

Frobenius Theorem¹, 1903

Let G be a finite group whose order is divisible by a number n .
Then $\sum_{d|n} w_d$ is divisible by n .

¹G. Frobenius, Über einen Fundamentalsatz der Gruppentheorie, *Sitz. Ber. Königl. Preuss. Akad. Wiss. Berlin*, 1903, 987–991.

Frobenius Theorem¹, 1903

Let G be a finite group whose order is divisible by a number n .
Then $\sum_{d|n} w_d$ is divisible by n .

Corollary

¹G. Frobenius, Über einen Fundamentalsatz der Gruppentheorie, *Sitz. Ber. Königl. Preuss. Akad. Wiss. Berlin*, 1903, 987–991.

Frobenius Theorem¹, 1903

Let G be a finite group whose order is divisible by a number n .
Then $\sum_{d|n} w_d$ is divisible by n .

Corollary

- If $p \in \omega(G)$ is prime, then $w_p \equiv -1 \pmod{p}$.

¹G. Frobenius, Über einen Fundamentalsatz der Gruppentheorie, *Sitz. Ber. Königl. Preuss. Akad. Wiss. Berlin*, 1903, 987–991.

Frobenius Theorem¹, 1903

Let G be a finite group whose order is divisible by a number n .
Then $\sum_{d|n} w_d$ is divisible by n .

Corollary

- If $p \in \omega(G)$ is prime, then $w_p \equiv -1 \pmod{p}$.
- If $d \in \omega(G) \setminus \{1\}$, then w_d is odd if and only if $d = 2$.

¹G. Frobenius, Über einen Fundamentalsatz der Gruppentheorie, *Sitz. Ber. Königl. Preuss. Akad. Wiss. Berlin*, 1903, 987–991.

Some bounds

G

Upper bound for $w_p^*(G)$

Some bounds

G	Upper bound for $w_p^*(G)$
Non- p -group	$\frac{p}{p+1} G $

Some bounds

G	Upper bound for $w_p^*(G)$
Non- p -group	$\frac{p}{p+1} G $
3-group of exponent $\neq 3$	$\frac{7}{9} G $

Some bounds

G	Upper bound for $w_p^*(G)$
Non- p -group	$\frac{p}{p+1} G $
3-group of exponent $\neq 3$	$\frac{7}{9} G $
2-group of exponent $\neq 2$	$\frac{3}{4} G $.

Some divisibility properties

Name	Year	Conditions	$w_{p^k}^*(G)$ is divisible by
------	------	------------	--------------------------------

Some divisibility properties

Name	Year	Conditions	$w_{p^k}^*(G)$ is divisible by
Miller	1919	$p^k > p > 2$	p^k

Some divisibility properties

Name	Year	Conditions	$w_{p^k}^*(G)$ is divisible by
Miller	1919	$p^k > p > 2$	p^k
Kulakof	1931	$p^k > p > 2$	p^{k+1}

Some divisibility properties

Name	Year	Conditions	$w_{p^k}^*(G)$ is divisible by
Miller	1919	$p^k > p > 2$	p^k
Kulakof	1931	$p^k > p > 2$	p^{k+1}
Berkovich	1991	$p^k > p > 2$	p^{k+p}

Freud and Pálffy, 1996

Set

Range

Freud and Pálffy, 1996

Set

Range

ω

\mathbb{N}^0

Freud and Pálffy, 1996

Set

Range

ω

\mathbb{N}^0

$\omega(2)$

$2\mathbb{N}^0 + 1$

Freud and Pálffy, 1996

Set	Range
ω	\mathbb{N}^0
$\omega(2)$	$2\mathbb{N}^0 + 1$
$\omega(3)$	Not known but contains $54\mathbb{N}^0 + 44$

Freud and Pálffy, 1996

Set	Range
ω	\mathbb{N}^0
$\omega(2)$	$2\mathbb{N}^0 + 1$
$\omega(3)$	Not known but contains $54\mathbb{N}^0 + 44$
$\omega(6)$	$2\mathbb{N}^0 \setminus \{4, 16, 28\}$

Freud and Pálffy, 1996

Set	Range
ω	\mathbb{N}^0
$\omega(2)$	$2\mathbb{N}^0 + 1$
$\omega(3)$	Not known but contains $54\mathbb{N}^0 + 44$
$\omega(6)$	$2\mathbb{N}^0 \setminus \{4, 16, 28\}$
$\omega(4k)$	$\varphi(k)\mathbb{N}^0$

Freud and Pálffy, 1996

Set	Range
ω	\mathbb{N}^0
$\omega(2)$	$2\mathbb{N}^0 + 1$
$\omega(3)$	Not known but contains $54\mathbb{N}^0 + 44$
$\omega(6)$	$2\mathbb{N}^0 \setminus \{4, 16, 28\}$
$\omega(4k)$	$\varphi(k)\mathbb{N}^0$
$\omega(p)$	Not known but contains $\{(p-1)p_1^{\alpha_1} \cdots p_n^{\alpha_n} : p_i^{\alpha_i} \equiv 1 \pmod{p}, i = 1, \dots, n\}$

A simple fact

If G is a group with $|w^*(G)| = 1$, then $|G| = 1$ or 2 .

Theorem

Let G be a finite group. Then $|w^(G)| = 2$ if and only if G is isomorphic to one of the following groups:*

Theorem

Let G be a finite group. Then $|w^(G)| = 2$ if and only if G is isomorphic to one of the following groups:*

(1) \mathbb{Z}_4 ,

Theorem

Let G be a finite group. Then $|w^(G)| = 2$ if and only if G is isomorphic to one of the following groups:*

- (1) \mathbb{Z}_4 ,
- (2) Q_8 ,

Theorem

Let G be a finite group. Then $|w^(G)| = 2$ if and only if G is isomorphic to one of the following groups:*

- (1) \mathbb{Z}_4 ,
- (2) Q_8 ,
- (3) a p -group of exponent p different from \mathbb{Z}_2 ,

Theorem

Let G be a finite group. Then $|w^(G)| = 2$ if and only if G is isomorphic to one of the following groups:*

- (1) \mathbb{Z}_4 ,
- (2) Q_8 ,
- (3) a p -group of exponent p different from \mathbb{Z}_2 ,
- (4) $H \times \mathbb{Z}_2$, where H is a p -group of exponent $p > 2$.

Preliminary results

Lemma

Let G be a finite group. Then $w(G) = \{1, p, q\}$ if and only if G is a Frobenius group whose kernel is a p -group of exponent p and complements are cyclic q -groups of order q , where p and q are distinct primes.

Preliminary results

Lemma

Let G be a finite group. Then $w(G) = \{1, p, q\}$ if and only if G is a Frobenius group whose kernel is a p -group of exponent p and complements are cyclic q -groups of order q , where p and q are distinct primes.

Lemma

Let G be a finite group. Then $w(G) = \{1, p, q, pq\}$ and $w^(G) = \{1, m, n\}$ if and only if $G/Z(G)$ is a Frobenius group, $Z(G) \cong \mathbb{Z}_2$ and either $G \cong \mathbb{Z}_2 \times (\mathbb{Z}_p^k \rtimes \mathbb{Z}_2)$ or $G \cong \mathbb{Z}_2^k \rtimes \mathbb{Z}_p$.*

Main results

Theorem

Let G be a finite group which is not a p -group. Then $|w^(G)| = 3$ if and only if there exists distinct primes p and q such that*

Main results

Theorem

Let G be a finite group which is not a p -group. Then $|w^(G)| = 3$ if and only if there exists distinct primes p and q such that*

- (1) G is a Frobenius group whose kernel is a p -group of exponent p and complements are cyclic q -groups of order q ,*

Main results

Theorem

Let G be a finite group which is not a p -group. Then $|w^(G)| = 3$ if and only if there exists distinct primes p and q such that*

- (1) G is a Frobenius group whose kernel is a p -group of exponent p and complements are cyclic q -groups of order q ,*
- (2) $G = O_{pqp}(G)$ is a 3-step group, $O_{pq}(G) = O_p(G) \rtimes \mathbb{Z}_q$ is a Frobenius group, $G/O_p(G) \cong \mathbb{Z}_q \rtimes \mathbb{Z}_p$ is a Frobenius group, $\exp(P) = p$ and $Q \cong \mathbb{Z}_q$,*

Main results

Theorem

Let G be a finite group which is not a p -group. Then $|w^(G)| = 3$ if and only if there exists distinct primes p and q such that*

- (1) G is a Frobenius group whose kernel is a p -group of exponent p and complements are cyclic q -groups of order q ,*
- (2) $G = O_{pqp}(G)$ is a 3-step group, $O_{pq}(G) = O_p(G) \rtimes \mathbb{Z}_q$ is a Frobenius group, $G/O_p(G) \cong \mathbb{Z}_q \rtimes \mathbb{Z}_p$ is a Frobenius group, $\exp(P) = p$ and $Q \cong \mathbb{Z}_q$,*
- (3) $G/Z(G)$ is a Frobenius group, $Z(G) \cong \mathbb{Z}_2$ and either $G \cong \mathbb{Z}_2 \times (\mathbb{Z}_p^k \rtimes \mathbb{Z}_2)$ or $G \cong \mathbb{Z}_2^k \rtimes \mathbb{Z}_p$,*

p -groups

Proposition

Let G be a p -group satisfying $|w^*(G)| = 3$. Then $\exp(G) \leq p^2(G)$.

p -groups

Proposition

Let G be a p -group satisfying $|w^*(G)| = 3$. Then $\exp(G) \leq p^2(G)$.

Example

A p -groups G of exponent p^2 satisfies $|w^*(G)| = 3$.

Thank You for Your Attention!