

# Induced cycles in circulant graphs

M. Farrokhi D. G.

Institute for Advanced Studies in Basic Sciences (IASBS)

May 16, 2017

- $G = \text{group}, \quad 1 \notin S = S^{-1} \subseteq G$   
 $\Gamma := \text{Cay}(G, S)$  is a graph with

$$V(\Gamma) = G \quad \text{and} \quad E(\Gamma) = \{\{g, gs\} : g \in G, s \in S\}.$$

- $G = \text{group}, \quad 1 \notin S = S^{-1} \subseteq G$   
 $\Gamma := \text{Cay}(G, S)$  is a graph with

$$V(\Gamma) = G \quad \text{and} \quad E(\Gamma) = \{\{g, gs\} : g \in G, s \in S\}.$$

- A graph is  $\Gamma$ -free if it does not have any induced subgraph isomorphic to  $\Gamma$ .

## Babai, 1976<sup>1</sup>

There is no minimal Cayley graph containing  $K_4 \setminus e$  or  $K_{3,5}$  as a subgraph.

---

<sup>1</sup>L. Babai, Chromatic number and subgraphs of Cayley graphs, *Theory and applications of graphs (Proc. Internat. Conf., Western Mich. Univ., Kalamazoo, Mich., 1976)*, pp. 10–22.

<sup>2</sup>J. Spencer, What's not inside a Cayley graph, *Combinatorica* **3**(2) (1983), 239–241.

## Babai, 1976<sup>1</sup>

There is no minimal Cayley graph containing  $K_4 \setminus e$  or  $K_{3,5}$  as a subgraph.

## Spencer, 1983<sup>2</sup>

There exists a class of graphs of bounded degree and arbitrary girth which cannot be embedded into minimal Cayley graphs as induced subgraphs.

---

<sup>1</sup>L. Babai, Chromatic number and subgraphs of Cayley graphs, *Theory and applications of graphs (Proc. Internat. Conf., Western Mich. Univ., Kalamazoo, Mich., 1976)*, pp. 10–22.

<sup>2</sup>J. Spencer, What's not inside a Cayley graph, *Combinatorica* **3**(2) (1983), 239–241.

## Definition

A graph is **perfect** if the chromatic and clique numbers of its induced subgraphs are equal.

---

<sup>1</sup>M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, *Ann. of Math. (2)* **164**(1) (2006), 51–229.

## Definition

A graph is **perfect** if the chromatic and clique numbers of its induced subgraphs are equal.

Chudnovsky, Robertson, Seymour, and Thomas, 2006<sup>1</sup>

A graph  $\Gamma$  is **perfect** iff neither  $\Gamma$  nor  $\Gamma^c$  has an induced odd cycle of length  $\geq 5$ .

---

<sup>1</sup>M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, *Ann. of Math. (2)* **164**(1) (2006), 51–229.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **bipartite** if and only if  $G$  is a **2-group**.

---

<sup>1</sup>M. Farrokhi D. G. and A. Mohammadian, Groups whose all (minimal) Cayley graphs have a given forbidden structure, Submitted.



MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **bipartite** if and only if  $G$  is a **2-group**.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **perfect** if and only if

---

<sup>1</sup>M. Farrokhi D. G. and A. Mohammadian, Groups whose all (minimal) Cayley graphs have a given forbidden structure, Submitted.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **bipartite** if and only if  $G$  is a **2-group**.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **perfect** if and only if

- $G$  is a **2-group**; or

---

<sup>1</sup>M. Farrokhi D. G. and A. Mohammadian, Groups whose all (minimal) Cayley graphs have a given forbidden structure, Submitted.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **bipartite** if and only if  $G$  is a **2-group**.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all minimal Cayley graphs of  $G$  are **perfect** if and only if

- $G$  is a **2-group**; or
- $G$  is isomorphic to  **$C_3$** ,  **$C_6$** ,  **$S_3$** ,  **$C_3 \times C_3$** ,  **$A_4$** , or  **$E$** , where

$$E = \langle a, b : a^3 = b^3 = [b, a, b] = [b, a, a] = 1 \rangle$$

is a group of order 27.

---

<sup>1</sup>M. Farrokhi D. G. and A. Mohammadian, Groups whose all (minimal) Cayley graphs have a given forbidden structure, Submitted.

MFDG and Mohammadian, 2015<sup>1</sup>

Let  $G$  be a nontrivial finite group. Then all Cayley graphs of  $G$  are **perfect** if and only if  $G$  is isomorphic to one of the groups  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_2 \times C_2$ ,  $S_3$ ,  $C_6$ ,  $C_2 \times C_2 \times C_2$ ,  $C_2 \times C_4$ ,  $D_8$ ,  $Q_8$ ,  $C_3 \times C_3$ .

---

<sup>1</sup>M. Farrokhi D. G. and A. Mohammadian, Groups whose all (minimal) Cayley graphs have a given forbidden structure, Submitted .

Beeler and Wooten, 2008<sup>1</sup>

The size of induced cycles inside  $\text{Cay}(C_m, S)$  is bounded above by  $m/2 + 1$ .

---

<sup>1</sup>R. A. Beeler and T. M. Wooten, Finding induced cycles within circulant graphs, *Congr. Numer.* **191** (2008), 33–49.

Beeler and Wooten, 2008<sup>1</sup>

The size of induced cycles inside  $\text{Cay}(C_m, S)$  is bounded above by  $m/2 + 1$ .

Beeler and Wooten, 2008<sup>1</sup>

The circulant graph  $\text{Cay}(C_m, S)$  contains an induced  $n$ -cycle iff there exists  $l_i \in \{0, 1\}$  and  $x_i \in S$  ( $i = 1, \dots, n$ ) such that

---

<sup>1</sup>R. A. Beeler and T. M. Wooten, Finding induced cycles within circulant graphs, *Congr. Numer.* **191** (2008), 33–49.

Beeler and Wooten, 2008<sup>1</sup>

The size of induced cycles inside  $\text{Cay}(C_m, S)$  is bounded above by  $m/2 + 1$ .

Beeler and Wooten, 2008<sup>1</sup>

The circulant graph  $\text{Cay}(C_m, S)$  contains an induced  $n$ -cycle iff there exists  $l_i \in \{0, 1\}$  and  $x_i \in S$  ( $i = 1, \dots, n$ ) such that

$$(1) \sum_{i=1}^n (-1)^{l_i} x_i \equiv 0 \pmod{n};$$

---

<sup>1</sup>R. A. Beeler and T. M. Wooten, Finding induced cycles within circulant graphs, *Congr. Numer.* **191** (2008), 33–49.

Beeler and Wooten, 2008<sup>1</sup>

The size of induced cycles inside  $\text{Cay}(C_m, S)$  is bounded above by  $m/2 + 1$ .

Beeler and Wooten, 2008<sup>1</sup>

The circulant graph  $\text{Cay}(C_m, S)$  contains an induced  $n$ -cycle iff there exists  $l_i \in \{0, 1\}$  and  $x_i \in S$  ( $i = 1, \dots, n$ ) such that

- (1)  $\sum_{i=1}^n (-1)^{l_i} x_i \equiv 0 \pmod{n}$ ;
- (2)  $\sum_{i=1}^j (-1)^{l_i} x_i \not\equiv \sum_{i=1}^k (-1)^{l_i} x_i \pmod{n}$  for  $j \neq k$ ;

---

<sup>1</sup>R. A. Beeler and T. M. Wooten, Finding induced cycles within circulant graphs, *Congr. Numer.* **191** (2008), 33–49.



Beeler and Wooten, 2008<sup>1</sup>

The size of induced cycles inside  $\text{Cay}(C_m, S)$  is bounded above by  $m/2 + 1$ .

Beeler and Wooten, 2008<sup>1</sup>

The circulant graph  $\text{Cay}(C_m, S)$  contains an induced  $n$ -cycle iff there exists  $l_i \in \{0, 1\}$  and  $x_i \in S$  ( $i = 1, \dots, n$ ) such that

- (1)  $\sum_{i=1}^n (-1)^{l_i} x_i \equiv 0 \pmod{n}$ ;
- (2)  $\sum_{i=1}^j (-1)^{l_i} x_i \not\equiv \sum_{i=1}^k (-1)^{l_i} x_i \pmod{n}$  for  $j \neq k$ ;
- (3)  $\left| \sum_{i=1}^j (-1)^{l_i} x_i - \sum_{i=1}^k (-1)^{l_i} x_i \right|_n \notin S$  for  $|j - k|_n \geq 2$ ;

---

<sup>1</sup>R. A. Beeler and T. M. Wooten, Finding induced cycles within circulant graphs, *Congr. Numer.* **191** (2008), 33–49.


$\mathfrak{F}(n) :=$  maximum non-negative integer  $m$  such that  
 all circulant graphs on  $C_m$  are  $C_n$ -free.

## First lemma

Assume  $m = 6k + i$  with  $i \in \{0, 1, 2, 3, 4, 5\}$ . Then the length of induced cycles in  $\Gamma = \text{Cay}(C_m, \{\pm 1, \pm 2\})$  are exactly

$$3, 3k + \left\lceil \frac{i}{2} \right\rceil, \dots, 4k + \left\lceil \frac{i}{2} \right\rceil - \delta_{1i}.$$

# Proof of the first lemma




There is a one-to-one correspondence between induced cycles of length  $k > 3$  and circular sequences of 1's and 2's of length  $k$  with no two consecutive 1 whose sum equal  $m$ .

## Second lemma

Assume  $m \neq 2$  is even. Then the length of induced cycles in  $\Gamma = \text{Cay}(C_m, \{\pm 1, \frac{m}{2}\})$  are exactly

$$4, 1 + \frac{m}{2}, 3 + \frac{m}{2}, \dots, 2 \left\lfloor \frac{m+4}{8} \right\rfloor - 1 + \frac{m}{2}.$$

# Proof of the second lemma




$\Gamma$  has squares with vertex sets of the form

$$\left\{x, x+1, x+1+\frac{m}{2}, x+\frac{m}{2}, x\right\}$$

for some  $x = 0, \dots, m-1$ .

# Proof of the second lemma



An induced cycle with  $t = 3$

An induced cycle  $C$  of  $\Gamma$  of length  $> 4$  is the union of some spokes and paths


$$P(x_1, x_2), \dots, P(x_{2t-1}, x_{2t})$$

in which  $x_{2s-1} < x_{2s}$  and

$$|x_{2s+1} - x_{2s}| = \frac{m}{2}$$

for all  $s = 1, \dots, t$  (with  $x_{2t+1} := x_1$ ).

# Proof of the second lemma



An induced cycle with  $t = 3$

The length of  $C$  equals


$$\begin{aligned} n &:= \sum_{s=1}^t (x_{2s} - x_{2s-1} + 1) \\ &= t + \sum_{s=1}^t (x_{2s} - x_{2s+1}). \end{aligned}$$

It follows that

$$n = t + (t \bmod 2) \frac{m}{2}.$$



# Proof of the second lemma



An induced cycle with  $t = 3$

Every vertex of  $C_m \setminus C$  is a counterpoint to some vertex of  $C \setminus \{x_1, \dots, x_{2t}\}$ . Hence


$$m = 2n - 2t.$$

So,  $t$  is odd. Also,  $t \leq \frac{m}{4}$  for

$$|V(P(x_{2s-1}, x_{2s}))| \geq 3$$

for all  $s = 1, \dots, t$ .

# Proof of the second lemma



An induced cycle with  $t = 5$

For any odd integer  $t \leq \frac{m}{4}$

$$0, 1, 2, 2 + \frac{m}{2}, 3 + \frac{m}{2}, 4 + \frac{m}{2}, \dots, 2t - 5, 2t - 4, 2t - 3, 2t - 3 + \frac{m}{2}, 2t - 2 + \frac{m}{2}, 2t - 1 + \frac{m}{2}, 2t - 1, \dots, \frac{m}{2}, 0.$$

is a cycle of length  $t + \frac{m}{2}$ .

## Theorem

For every  $n > 2$  we have

$$\mathfrak{F}(n) = 12k + \left\lceil \frac{3i}{2} \right\rceil - \delta_{1, \lfloor \frac{i}{2} \rfloor} - 1,$$

where  $n = 8k + i$  with  $i \in \{0, \dots, 7\}$ .

# Proof of the theorem

Let  $m := \mathfrak{F}(n) > n$ . We claim that

$$m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - \delta_{1, \lfloor \frac{i}{2} \rfloor}.$$

# Proof of the theorem

Let  $m := \mathfrak{F}(n) > n$ . We claim that

$$m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - \delta_{1, \lfloor \frac{i}{2} \rfloor}.$$

- There exists a monomorphism

$$\rho : C_n \hookrightarrow \Gamma := \text{Cay}(C_m, S)$$

for some  $1 \in S = -S \subseteq C_m$ .

# Proof of the theorem

Let  $m := \mathfrak{F}(n) > n$ . We claim that

$$m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - \delta_{1, \lfloor \frac{i}{2} \rfloor}.$$

- There exists a monomorphism

$$\rho : C_n \hookrightarrow \Gamma := \text{Cay}(C_m, S)$$

for some  $1 \in S = -S \subseteq C_m$ .

- Every vertex in  $\rho(C_n)$  is adjacent to  $|S| - 2$  vertices of  $C_m \setminus \rho(C_n)$ . Thus

$$(m - n)|S| \geq n(|S| - 2)$$

so that

$$m \geq 2n(1 - |S|^{-1}).$$

## Proof of the theorem

- Clearly,  $|S| > 2$ .

# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$



# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

- If  $|S| = 3$  then  $m$  is even and  $S = \{\pm 1, \frac{m}{2}\}$ .

# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

- If  $|S| = 3$  then  $m$  is even and  $S = \{\pm 1, \frac{m}{2}\}$ .
  - $C_n$  is an induced subgraph of a circulant graph on  $C_{m+1}$ . Thus

$$m + 1 \geq 12k + \left\lceil \frac{3i}{2} \right\rceil \Rightarrow m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

- If  $|S| = 3$  then  $m$  is even and  $S = \{\pm 1, \frac{m}{2}\}$ .
  - $C_n$  is an induced subgraph of a circulant graph on  $C_{m+1}$ . Thus

$$m + 1 \geq 12k + \left\lceil \frac{3i}{2} \right\rceil \Rightarrow m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

Suppose  $m = 12k + \left\lceil \frac{3i}{2} \right\rceil - 1$ .

# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

- If  $|S| = 3$  then  $m$  is even and  $S = \{\pm 1, \frac{m}{2}\}$ .
  - $C_n$  is an induced subgraph of a circulant graph on  $C_{m+1}$ . Thus

$$m + 1 \geq 12k + \left\lceil \frac{3i}{2} \right\rceil \Rightarrow m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

Suppose  $m = 12k + \left\lceil \frac{3i}{2} \right\rceil - 1$ .

- $i = 2, 3, 6, \text{ or } 7$ .

# Proof of the theorem

- Clearly,  $|S| > 2$ .
- If  $|S| \geq 4$  then

$$m \geq \frac{3}{2}n > 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

- If  $|S| = 3$  then  $m$  is even and  $S = \{\pm 1, \frac{m}{2}\}$ .
  - $C_n$  is an induced subgraph of a circulant graph on  $C_{m+1}$ . Thus

$$m + 1 \geq 12k + \left\lceil \frac{3i}{2} \right\rceil \Rightarrow m \geq 12k + \left\lceil \frac{3i}{2} \right\rceil - 1.$$

Suppose  $m = 12k + \left\lceil \frac{3i}{2} \right\rceil - 1$ .

- $i = 2, 3, 6$ , or  $7$ .
- $n - \frac{m}{2}$  is odd by the second Lemma. Thus  $i = 2$  or  $3$ .

## Corollary

*Let  $G$  be a finite group whose all Cayley graphs are  $C_n$ -free. Then order of elements of  $G$  are bounded above by  $\mathfrak{F}(n)$ .*

Thank You for Your Attention!